

Jan Mařík

Derivatives and convexity

Real Anal. Exchange 17 (2) (1991/92), 745–747

Persistent URL: <http://dml.cz/dmlcz/502145>

Terms of use:

© Michigan State University Press, 1991

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Jan Mařík, Department of Mathematics, Michigan State University, East Lansing, Michigan 48824.

DERIVATIVES AND CONVEXITY

Lemma 4.4 in [MW] says that if the composition of a function F strictly convex on an open interval containing the range of a derivative f is also a derivative, then both functions f and $F \circ f$ are Lebesgue functions. Theorem 4 of this note generalizes that result; f is there an n -tuple of derivatives and F is strictly convex on an open convex set containing the range of f . Theorems 5 and 8 deal with the one-dimensional case without the assumption that the domain of definition of F is open.

1. Notation. The symbols D, D^+, C_{ap}, L mean the systems of all derivatives, nonnegative derivatives, approximately continuous functions and Lebesgue functions on the interval $I = [0, 1]$, respectively. Symbols like $\int_a^b f$ or $\int_S f$ mean the corresponding Lebesgue integrals. The letter n denotes a natural number and R^n the n -dimensional Euclidean space. For $z = (z_1, \dots, z_n) \in R^n$ we write $|z| = (z_1^2 + \dots + z_n^2)^{1/2}$.

2. Lemma. Let $f_1, \dots, f_n \in D$, $x \in I$. Set $f = (f_1, \dots, f_n)$, $b = f(x)$, $S = f(I)$. Let H be a function on S such that $H(b) = 0$, $H \circ f \in D$ and that for each $\varepsilon \in (0, \infty)$ we have

$$(1) \quad \inf\{H(z)/|z - b|; z \in S, |z - b| > \varepsilon\} > 0.$$

Then

$$(2) \quad \frac{1}{y-x} \int_x^y |f - b| \rightarrow 0 \quad (y \rightarrow x, y \in I).$$

Proof. Let $\varepsilon \in (0, \infty)$ and let α be the infimum in (1). Set $\varphi = H \circ f$. Then $\varphi \geq \alpha(|f - b| - \varepsilon)$ whence $|f - b| \leq \alpha^{-1}\varphi + \varepsilon$ on I . Because $\varphi \in D$ and $\varphi(x) = 0$, we have $\limsup \frac{1}{y-x} \int_x^y |f - b| \leq \varepsilon$ ($y \rightarrow x, y \in I$) which proves (2).

3. Lemma. Let $f, g \in C_{ap}$, $f \in D$ and $|g| \leq f$. Then $g \in L$. (See [M], 1.8.)

4. Theorem. Let F be a strictly convex function on an open convex set G in R^n . Let $f_1, \dots, f_n \in D$, $f = (f_1, \dots, f_n)$. Suppose that $f(I) \subset G$ and that $F \circ f \in D$. Then $f_1, \dots, f_n, F \circ f \in L$.

Proof. Let $b \in G$. It is well-known (see, e.g., [Mt], V, 1, Korollar 4) that there is a linear function λ such that $\lambda(b) = F(b)$ and $\lambda \leq F$ on G . Since F is strictly convex, we have $\lambda < F$ on $G \setminus \{b\}$.

Now let ε be a positive number such that the set $A = \{z \in R^n; |z - b| = \varepsilon\}$ is a part of G . Set $H = F - \lambda$. Since F is continuous (see, e.g., [Mt], X, 1, Satz 2) and $H > 0$ on A , there is a $\beta \in (0, \infty)$ such that $H > \beta$ on A .

Let $z \in G$ and $|z - b| > \varepsilon$. Set $v = (z - b)/|z - b|$, $J = \{t \in (-\infty, \infty); b + tv \in G\}$, $h(t) = H(b + tv)$ ($t \in J$). It is easy to see that h is (strictly) convex. Clearly $h(\varepsilon) > \beta$. Thus $h(t) \geq th(\varepsilon)/\varepsilon > t\beta/\varepsilon$ for each $t \in J \cap (\varepsilon, \infty)$. It follows that $H(z) = h(|z - b|) > |z - b|\beta/\varepsilon$ so that, by 2, $f_1, \dots, f_n \in L$. Thus $H \circ f \in C_{ap} \cap D^+$. By 3 we have $H \circ f \in L$ whence $F \circ f \in L$.

5. Theorem. Let $f \in D$. Let F be a strictly convex function on $f(I)$ such that $F \circ f \in D$. Then $f \in L$. If, moreover, F is continuous, then also $F \circ f \in L$.

Proof. Let $x \in I$, $b = f(x)$. Set $S = f(I)$. If $b \in \text{int}S$, we get (2) as in the preceding proof. If, e.g., $b = \min S$, then (2) is obvious (since $f \geq b$ and $f \in D$). Thus $f \in L$.

Now suppose that F is continuous. There is a linear function λ such that $F \geq \lambda$ on S . Then $(F - \lambda) \circ f \in C_{ap} \cap D^+$ whence, by 3, $(F - \lambda) \circ f \in L$. Thus $F \circ f \in L$.

Remark. The example in 7 shows that the relation $F \circ f \in L$ may be false, if F is not continuous (even if $f, F \circ f \in D$ etc.).

We need first a lemma.

6. Lemma. Let f be a nonnegative (Lebesgue) measurable function on I . Let $\frac{1}{x} \int_0^x f^2 \rightarrow 1$, $\lim_{x \rightarrow 0+} f(x) = 0$. Then $\frac{1}{x} \int_0^x f \rightarrow 0$ ($x \rightarrow 0+$).

Proof. Let $\varepsilon \in (0, \infty)$. Set $z_k = 2^{-k}$, $J_k = [z_k, 2z_k]$, $S_k = \{x \in J_k; f(x) > \varepsilon\}$ ($k = 1, 2, \dots$). Let β_k be the measure of S_k . Then $\frac{1}{z_k} \int_{J_k} f \leq \frac{1}{z_k} \int_{S_k} f + \varepsilon$ and $\frac{1}{z_k} \int_{S_k} f \leq (\frac{1}{z_k} \int_{J_k} f^2 \cdot \beta_k/z_k)^{\frac{1}{2}} \rightarrow 0$ ($k \rightarrow \infty$) which easily implies our assertion.

7. Example. Let $F(z) = z^2$ ($z \in (0, \infty)$), $F(0) = 1$. Let f be a function such that $f \in C_{ap}$, f is positive and continuous on $(0, 1]$, $f(0) = 0$ and $\frac{1}{x} \int_0^x f^2 \rightarrow 1$ ($x \rightarrow 0+$). (It is easy to construct such a function.) Then $F \circ f \in D \setminus C_{ap}$ and, by 6, $f \in D$.

8. Theorem. Let $f \in D$. Suppose that f is not constant. Let F be a strictly convex function on $f(I)$ and let $F \circ f \in D \cap C_{ap}$. Then F is continuous.

Proof. By 5 we have $f \in C_{ap}$. Define a function F_0 on $S = f(I)$ setting $F_0 = F$ on $\text{int } S$ and $F_0(x) = \lim F(z)$ ($z \rightarrow x, z \in \text{int } S$) for $x \in S \setminus \text{int } S$. Then F_0 is continuous on S so that $F_0 \circ f \in C_{ap}$. Set $\varphi = (F - F_0) \circ f$. Then $\varphi \in C_{ap}$. We see that φ is a Darboux function that has at most three values. Thus φ is constant, $\varphi = 0$ on I , $F = F_0$ on S , F is continuous.

Remark. Our proof of 8 would fail in more dimensions, because then the limit used there need not exist. (Take, e.g., $G = (0, \infty) \times (0, \infty) \cup (0, 0)$, $F(x, y) = x^2/y + x^2 + y^2$ for $x, y > 0$, $F(0, 0) = 0$.)

References

- [M] Jan Mařík, *Transformation and multiplication of derivatives*, Contemporary Mathematics, Vol. 42 (1985), 119-134.
- [Mt] Jürg T. Marti, *Konvexe Analysis*, Birkhäuser Verlag, 1977.
- [MW] Jan Mařík and Clifford E. Weil, *Sums of powers of derivatives*, Proc. AMS, Vol. 112, No. 3, 807-817.

Received December 12, 1991