

EQUADIFF 10

Pavol Quittner

A priori bounds for solutions of parabolic problems and applications

In: Miroslav Krbeč and Jaromír Kuben (eds.): Proceedings of Equadiff 10, Czechoslovak International Conference on Differential Equations and Their Applications, Prague, August 27-31, 2001, [Part 1] Invited Lectures. Masaryk University, Brno, 2002. CD-ROM issued as a complement to the journal edition *Mathematica Bohemica* 2002/2. pp. 185--196.

Persistent URL: <http://dml.cz/dmlcz/700322>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

A priori bounds for solutions of parabolic problems and applications

Pavol Quittner

Institute of Applied Mathematics
Comenius University
Mlynská dolina
84 248 Bratislava, Slovakia
Email: quittner@fmph.uniba.sk

Abstract. We review some recent results concerning a priori bounds for solutions of superlinear parabolic problems and their applications.

MSC 2000. 35B45, 35K60, 35J65

Keywords. a priori estimate, blow-up rate, periodic solution, multiplicity

1 Introduction

In this paper we study mainly parabolic problems of the form

$$\begin{cases} u_t - \Delta u = f(x, u), & x \in \Omega, t > 0, \\ u = 0, & x \in \Gamma, t > 0, \\ u(x, 0) = u_0(x), & x \in \Omega, \end{cases} \quad (1.1)$$

where Ω is a domain in \mathbb{R}^n with a smooth compact boundary Γ and f is a Carathéodory function which is superlinear in u . Some generalizations and modifications of (1.1) are also considered.

It is well known that under suitable assumptions on f the problem (1.1) is well posed in an appropriate Banach space X ($X = L_\infty(\Omega)$, for example). Denote by $u(t, u_0)$ the solution of this problem and let $T_{\max}(u_0)$ be its maximal existence time. Assume $\delta > 0$. Our main aim is to show that for a large class of nonlinearities, the norm of $u(t, u_0)$, $t \in [0, T_{\max}(u_0) - \delta)$, can be bounded by a constant which

depends only on δ and on the norm of the initial condition u_0 . In other words, we are interested in the estimate

$$\|u(t, u_0)\|_X \leq C(\delta, c_0) \begin{cases} \text{for any } u_0 \in X \text{ with } \|u_0\|_X \leq c_0, \\ \text{and any } t < T_{\max}(u_0) - \delta, \end{cases} \tag{1.2}$$

where $T_{\max}(u_0) - \delta = \infty$ and $C(\delta, c_0)$ does not depend on δ if $T_{\max}(u_0) = \infty$. Note that under some circumstances global solutions are bounded even if the estimate (1.2) does not hold for these solutions, see V. Galaktionov and J.L. Vázquez [16] or M. Fila and P. Poláčik [13]. For a survey on the boundedness of global solutions we refer to [12].

We shall also mention some results on universal bounds of the form

$$\|u(t, u_0)\|_X \leq C(\delta_1, \delta_2) \quad \text{for any } t \in (\delta_1, T_{\max}(u_0) - \delta_2), \tag{1.3}$$

where the constant $C(\delta_1, \delta_2)$ does not depend on u_0 at all.

The bound (1.2) has several important consequences. It implies the continuity of the maximal existence time $T_{\max} : X \rightarrow (0, \infty]$, it plays a crucial role in establishing the blow-up rate of blowing-up solutions, in the study of domains of attraction of stable equilibria and connecting orbits between various equilibria. It can also be used for the proof of existence of multiple stationary and periodic solutions.

Let us first discuss the model case $f(x, u) = |u|^{p-1}u$, $p > 1$, $\Omega \subset \mathbb{R}^n$ bounded. Set

$$p_S := (n + 2)/(n - 2) \text{ if } n \geq 3, \quad p_S := \infty \text{ otherwise.}$$

The bounds (1.2) and (1.3) and their proofs are strongly related to the a priori estimates for positive stationary solutions of (1.1) which were proved in the subcritical case $p < p_S$ by D.G. de Figueiredo, P.-L. Lions and R.D. Nussbaum [11] and B. Gidas and J. Spruck [17] (partial results were obtained before by R.E.L. Turner [36], R.D. Nussbaum [26], H. Brézis and R.E.L. Turner [5]). Due to the result of S.I. Pohozaev [27], the condition $p < p_S$ is optimal in these estimates (at least if Ω is starshaped). The bound (1.2) for the time-dependent solutions of this model problem was derived for any $p < p_S$ by the author in [28] under the assumption $T_{\max}(u_0) = \infty$. Partial results requiring a stronger condition on p and/or nonnegativity of u were previously obtained by W.-M. Ni, P.E. Sacks and J. Tavantzis [25], T. Cazenave and P.-L. Lions [6] and Y. Giga [18]. The condition $p < p_S$ is optimal again.

Considering a general superlinear function f , the results on a priori estimates for positive stationary solutions mentioned above are far from satisfactory: they require either Ω to be convex or various technical conditions on f (either monotonicity of $u \mapsto f(x, u)u^{-p_S}$ in [11] or a precise asymptotic behavior of $f(x, u)$ as $u \rightarrow +\infty$ in [17]). From this point of view it is interesting to know to what extent can one generalize the results of [28] concerning the estimate (1.2) for the time-dependent solutions. The approach in [28] is based on a bootstrap argument, interpolation, energy and maximal regularity estimates. It turns out that the assumption $T_{\max}(u_0) = \infty$ and the precise asymptotic behavior of the nonlinearity f

as $|u| \rightarrow \infty$ are not important for this approach. Moreover, the results remain true for more general differential operators, boundary conditions and nonlinearities.

In Section 2 we discuss the estimate (1.2) for (1.1) and some of its consequences (including continuity of T_{\max} and existence of nontrivial equilibria) in the case of a bounded spatial domain Ω . In Section 3 we study the unbounded domain case. Section 4 is devoted to time-dependent nonlinearities and the existence of periodic solutions. In Section 5 we briefly mention some results on the universal bound (1.3) and initial and final blow-up rates. In Sections 6,7 and 8 we deal with nonlinear boundary conditions, nonlocal problems and problems involving measures, respectively. For one-dimensional problems we refer to [31, Section 6] and [7, Section 5].

2 Bounded domains

Denote $F(x, u) := \int_0^u f(x, v) dv$ and assume that there exist positive constants $p_1, p_2, d_1, d_2, d_3, d_4, \beta, r$ and nonnegative functions

$$a_1 \in L_{(p_1+1)/p_1}(\Omega), \quad a_2 \in L_{(p_2+1)/p_2}(\Omega), \quad a_3 \in L_1(\Omega), \quad a_4 \in L_\beta(\Omega) \quad (2.1)$$

such that

$$1 < p_1 \leq p_2 < p_S, \quad d_3 > 2, \quad \beta > n/2, \quad r < p_S, \quad (2.2)$$

$$|f(x, u)| \leq d_2|u|^{p_2} + a_2(x), \quad (2.3)$$

$$f(x, u) \operatorname{sign}(u) \geq d_1|u|^{p_1} - a_1(x), \quad (2.4)$$

$$f(x, u)u \geq d_3F(x, u) - a_3(x), \quad (2.5)$$

$$|f(x, u) - f(x, v)| \leq d_4(a_4(x) + |u|^{r-1} + |v|^{r-1})|u - v|. \quad (2.6)$$

Assume also that either $p_2 < p_{CL}$ or

$$p_2 - p_1 < \kappa_1(p_2), \quad (2.7)$$

where $\kappa_1 : (1, p_S) \rightarrow (0, \infty)$ is defined in [31] (cf. Figures 1 and 2 below) and

$$p_{CL} := (3n + 8)/(3n - 4) \text{ if } n \geq 2, \quad p_{CL} := \infty \text{ if } n = 1.$$

Set

$$E(u) := \frac{1}{2} \int_\Omega |\nabla u|^2 dx - \int_\Omega F(x, u) dx.$$

Then we have the following theorem (see [31] and [32]).

Theorem 2.1. *Consider the problem (1.1). Let Ω be a smoothly bounded domain in \mathbb{R}^n . Assume (2.1)-(2.6) and either $p_2 < p_{CL}$ or (2.7). Set $X := H_0^1(\Omega)$. Then the estimate (1.2) is true, $T_{\max} : X \rightarrow (0, \infty]$ is continuous and*

$$E(u(t, u_0)) \rightarrow -\infty \quad \text{as } t \rightarrow T_{\max}(u_0) - \quad \text{if } T_{\max}(u_0) < \infty. \quad (2.8)$$

If, in addition, $\beta > n$, $f(\cdot, 0) \in L_\beta(\Omega)$, u_s is an asymptotically stable equilibrium of (1.1) in X and D_A denotes its domain of attraction,

$$D_A = \{u_0 \in X \mid u(t, u_0) \text{ exists globally, } u(t, u_0) \rightarrow u_s \text{ in } X \text{ as } t \rightarrow \infty\},$$

then there exist stationary solutions $u_+, u_-, \tilde{u} \in \partial D_A$ of (1.1) such that $u_+ > u_s > u_-$ and $\tilde{u} - u_s, \tilde{u} - u_+, \tilde{u} - u_-$ change sign.

Remarks 2.1. (i) The condition (2.7) in Theorem 2.1 seems to be of technical nature. In fact, if

$$f(x, u)u \leq d_5 F(x, u) + a_5(x), \quad d_5 > 0, \quad a_5 \in L_1(\Omega),$$

then this assumption can be replaced by

$$p_2 - p_1 < \kappa_2(p_2), \tag{2.9}$$

where $\kappa_2 : (1, p_S) \rightarrow (0, \infty)$ is defined in [31], $\kappa_2 > \kappa_1$ (see Figures 1 and 2). The same is true for all assertions in the subsequent sections.

In Figures 1 and 2 we set $p(n) := 1 + 4/n$,

$$p^* := \begin{cases} \frac{9n^2 - 4n + 16\sqrt{n(n-1)}}{(3n-4)^2} & \text{if } n \geq 2, \\ +\infty & \text{if } n = 1. \end{cases}$$

Note that the condition (2.7) or (2.9) is superfluous if $p \leq p(n)$ or $p < p^*$, respectively.




Figure 1. Functions κ_1, κ_2 for $n = 2$




Figure 2. Functions κ_1, κ_2 for $n = 3$:
 $p(n) = 2 + 1/3, p_{CL} = 3.4, p^* \doteq 4.3, p_S = 5, \kappa^* \doteq 0.27$

(ii) The property (2.8) plays an important role in the proof of complete blow-up, see [2]. This property was proved before by H. Zaag [37] for the model case $f(x, u) = |u|^{p-1}u$ under additional assumptions $p(3n - 4) < 3n + 8$ or $u \geq 0$.

(iii) Continuity of T_{\max} for nonnegative solutions, bounded domains Ω and convex functions $f = f(u)$ with subcritical growth was previously proved by P. Baras and L. Cohen [2]. Note that the function T_{\max} need not be continuous in the supercritical case, due to a result of V. Galaktionov and J.L. Vázquez [16]. More precisely, the set $\{u_0 \mid T_{\max}(u_0) = \infty\}$ need not be closed.

(iv) If $u_s = 0$ in Theorem 2.1 then this theorem guarantees the existence of a sign-changing equilibrium \tilde{u} of (1.1) lying on ∂D_A . Similar assertions (without the information $\tilde{u} \in \partial D_A$) were proved by variational and topological methods by many authors: see the discussion in [32], for example.

3 Unbounded domains

Let F and E be the same as in Section 2. Assume that there exist positive constants $p_1, p_2, d_1, d_2, d_3, d_4, \beta, r$ satisfying (2.2) and nonnegative constants e_1, C_1 such that

$$|f(x, u)| \leq d_2(|u|^{p_2} + |u|) + a_2(x), \tag{3.1}$$

$$f(x, u) \operatorname{sign}(u) \geq d_1|u|^{p_1} - e_1|u| - a_1(x), \tag{3.2}$$

$$f(x, u)u \geq d_3F(x, u) + C_1u^2 - a_3(x), \tag{3.3}$$

$$|f(x, u) - f(x, v)| \leq (a_4(x) + d_4(1 + |u|^{r-1} + |v|^{r-1}))|u - v|, \tag{3.4}$$

$$f(\cdot, 0) \in L_\beta(\Omega), \tag{3.5}$$

where a_1, a_2, a_3, a_4 satisfy (2.1). Notice that the assumptions (3.1)–(3.4) are equivalent to (2.3)–(2.6) if Ω is bounded. The conditions above guarantee, in particular, that the problem (1.1) is well posed in $H_0^1(\Omega)$. Denote by $T_{\max}(u_0)$ the maximal existence time of the solution in $H_0^1(\Omega)$. Then we have the following theorem (see [31]).

Theorem 3.1. *Let $\Omega \subset \mathbb{R}^n$ have a smooth compact boundary (or Ω be a half-space). Assume (2.1)–(2.2), (3.1)–(3.5) and (2.7). Set*

$$X := H_0^1(\Omega) \cap L_{(p_2+1)/p_2}(\Omega) \cap L_\infty(\Omega),$$

assume $u_0 \in X$ and let

$$T_{\max}^X(u_0) := \sup\{t \in [0, T_{\max}(u_0)) \mid u(\tau) \in X \text{ for } \tau \leq t\}.$$

Then the following holds:

- (i) $T_{\max}^X(u_0) = T_{\max}(u_0)$, $T_{\max} : X \rightarrow (0, \infty]$ is continuous and (2.8) is true.
- (ii) Let $C_1 > 0$ in (3.3) and let there exist constants $d_6, \lambda > 0$, $\alpha \in (1, p_2)$, a nonnegative function $a_6 \in L_{(p_2+1)/p_2}(\Omega)$ and a bounded measurable function $V : \Omega \rightarrow [\lambda, \infty)$ such that

$$|f(x, v) + V(x)v| \leq d_6(|v|^{p_2} + |v|^\alpha) + a_6(x).$$

Let $u_0 \in X$ and $T_{\max}(u_0) = \infty$. Then there exists a constant $C = C(\|u_0\|_X)$ such that

$$\|u(t)\|_X \leq C \quad \text{for any } t \geq 0. \tag{3.6}$$

Remarks 3.1. (i) We are not able to show the bound (1.2) if $T_{\max}(u_0) < \infty$. Consequently, the proof of the continuity of T_{\max} requires some additional arguments (using a refinement of the concavity method due to H.A. Levine [23]). Note that all previous results concerning the estimate (3.6) and the continuity of T_{\max} required a stronger assumption on the growth of f or they were restricted to nonnegative solutions and nonlinearities with a precise asymptotic behavior (see C. Fermanian Kammerer, F. Merle and H. Zaag [10], for example).

(ii) If $\lambda > 0$ and $1 < p < p_S$ then $f(x, u) := |u|^{p-1}u - \lambda u$ satisfies all assumptions of Theorem 3.1(ii).

4 Periodic solutions

In this section we study a priori estimates of solutions and existence of positive periodic solutions of the problem

$$\begin{cases} u_t - \Delta u = m(t)f(u), & x \in \Omega, t > 0, \\ u = 0, & x \in \Gamma, t > 0, \\ u(x, 0) = u_0(x), & x \in \Omega, \end{cases} \tag{4.1}$$

where Ω is a smoothly bounded domain in \mathbb{R}^n , $m > 0$ is T -periodic and $f(u) = |u|^{p-1}u$, $1 < p < p_S$. We refer to [32] for the case of a general superlinear function $f = f(u)$ and to [22] for the case $f = f(x, u)$.

Theorem 4.1 (see [32]). *Let $\Omega \subset \mathbb{R}^n$ be smoothly bounded, $m \in W_\infty^1(\mathbb{R})$ be T -periodic, $m(t) \geq m_0 > 0$ for any t , $f(u) = |u|^{p-1}u$, $1 < p < p_S$. Set $X := H_0^1(\Omega)$.*

(i) *Let u be the solution of (4.1), $T_{\max}(u_0) \geq T + \delta$, $\delta > 0$. Then there exists a constant $C = C(\|u_0\|_X, \delta, T)$ such that*

$$\|u(t)\|_X \leq C \quad \text{for any } t \in [0, T].$$

(ii) *Assume*

$$\frac{(m'(t))^-}{m(t)} < \frac{2n - (n - 2)(p + 1)}{r^2(\Omega)} \quad \text{for a.a. } t \in (0, T), \tag{4.2}$$

where $r(\Omega)$ is the radius of the smallest ball containing Ω and $a^- := \max(0, -a)$. Then there exists at least one positive T -periodic solution of (4.1) and there exists $C > 0$ such that any positive T -periodic solution of (4.1) satisfies

$$\|u(t)\|_X \leq C \quad \text{for any } t \in [0, T].$$

Remarks 4.1. (i) The technical assumption (4.2) is superfluous if $p(n - 2) < n$.

(ii) Existence of positive periodic solutions of (4.1) with $f(u) = |u|^{p-1}u$ (and more general nonlinearities) was obtained before by M.J. Esteban in [8] and [9] under the additional assumptions $(3n - 4)p < 3n + 8$ and $p(n - 2) < n$, respectively.

(iii) Assertion (i) in Theorem 4.1 is based on the fact that the functional

$$V(u(t)) = \frac{1}{2} \int_\Omega |\nabla u(t)|^2 dx - m(t) \int_\Omega F(u(t)) dx$$

is “almost” a Lyapunov functional for (4.1). A Pohozaev’s type identity plays a significant role in the proof of (ii) (cf. [11] in the elliptic case).

(iv) A different approach to problems without variational structure can be found in [33].

5 Universal bounds and blow-up rates

In this section we are interested in the universal bound (1.3) for positive solutions of (1.1) (note that this bound cannot be true for all solutions, in general). The following theorem follows from the results in [35].

Theorem 5.1. *Consider the problem (1.1) with $\Omega \subset \mathbb{R}^n$ being (smoothly) bounded and convex, $f(x, u) = |u|^{p-1}u$, $1 < p < p_S$, $u_0 \geq 0$. Let $p(n - 3) < n - 1$ if $n \geq 5$ and $T_{\max}(u_0) \geq T_0 > 0$. Set $X := L_\infty(\Omega)$. Then there exist $C(p, \Omega, T_0) > 0$ and $\alpha = \alpha(n, p) > 0$ such that*

$$\|u(t)\|_X \leq C(p, \Omega, T_0)(1 + t^{-\alpha} + (T_{\max}(u_0) - t)^{-1/(p-1)})$$

for any $t \in (0, T_{\max}(u_0))$, where $(T_{\max}(u_0) - t)^{-1/(p-1)} := 0$ if $T_{\max}(u_0) = \infty$.

Remarks 5.1. (i) The convexity of Ω is needed only for the estimate of $u(t)$ close to $T_{\max}(u_0)$. The assumption $p < (n - 1)/(n - 3)$ for $n \geq 5$ seems to be of technical nature.

(ii) If $p < 1 + 2/(n + 1)$ then one can choose $\alpha = (n + 1)/2$ in Theorem 5.1 and this choice is optimal. Note that this initial blow-up rate exponent is different from the corresponding exponent for the homogeneous Neumann problem (see [35]).

(iii) Due to the result of M.-F. Bidaut-Véron in [4] concerning the Cauchy problem, one can conjecture that the choice $\alpha = 1/(p - 1)$ should be possible (and optimal) for $p \geq 1 + 2/(n + 1)$ but this seems to be an open problem.

(iii) The (final) blow-up rate estimate

$$\|u(t)\|_X \leq C(p, \Omega, u_0)(T_{\max}(u_0) - t)^{-1/(p-1)} \tag{5.1}$$

(where C depends on u_0 !) is true also for sign-changing solutions and any $p \in (1, p_S)$ if $\Omega = \mathbb{R}^n$. This follows from a very recent result of Y. Giga, S. Matsui and S. Sasayama based on the approach in [28]. If $p(3n - 4) < 3n + 8$ or $u_0 \geq 0$ and $p < p_S$ then (5.1) was proved by Y. Giga and R.V. Kohn [19] for both unbounded and bounded convex domains. On the other hand, it is known that such an estimate fails, in general, for $p \geq p_S$, see the results of S. Filippas, M.A. Herrero and J.J.L. Velázquez in [15], [20] and [21]. Concerning universal blow-up rate estimates for positive solutions in unbounded domains we refer to J. Matos and Ph. Souplet [24].

(iv) First results concerning universal bounds for global positive solutions of (1.1) with $f(x, u) = |u|^{p-1}u$ and Ω bounded were obtained by M. Fila, Ph. Souplet, F. Weissler in [14] and the author in [30].

6 Nonlinear boundary conditions

In this section we study a priori estimates for global solutions of the problem

$$\begin{cases} u_t = \Delta u - au, & x \in \Omega, t \in (0, \infty), \\ u_\nu = |u|^{q-1}u, & x \in \Gamma, t \in (0, \infty), \\ u(x, 0) = u_0(x), & x \in \Omega, \end{cases} \tag{6.1}$$

where $a > 0$, $q > 1$, Ω is a smoothly bounded domain in \mathbb{R}^n and ν denotes the outer unit normal on the boundary Γ . Since we study only global solutions, the bounds (1.2) and (1.3) have the form

$$\|u(t)\|_X \leq C(\|u_0\|_X), \quad \text{for any } t > 0, \tag{6.2}$$

$$\|u(t)\|_X \leq C(\delta), \quad \text{for any } t > \delta. \tag{6.3}$$

The following result is proved in [34].

Theorem 6.1. *Consider the problem (6.1). Let $X := H^1(\Omega)$ and $q(n - 2) < n$.*

(i) Let $T_{\max}(u_0) = \infty$. If $u_0 \geq 0$ or $q < q^*$, where

$$q^* = \begin{cases} +\infty & \text{if } n = 1, \\ (9n^2 - 22n + 24 + 8\sqrt{4n^2 - 10n + 8}) / (3n - 4)^2 & \text{if } n > 1, \end{cases}$$

then the bound (6.2) is true.

(ii) Assume $q(n - 4) < n - 3$ if $n \geq 7$. Then the bound (6.3) is true for all global nonnegative solutions of (6.1).

Remarks 6.1. (i) The value $q_S := n/(n - 2)$ is the limiting exponent for which the trace operator maps $H^1(\Omega)$ into $L_{q+1}(\Gamma)$. Unlike the case of the homogeneous Dirichlet boundary condition, it is not clear whether the subcriticality condition $q < q_S$ is necessary for the a priori bounds mentioned above.

(ii) The assumptions $q < q^*$ and $q < (n - 3)/(n - 4)$ for $n \geq 7$ seem to be of technical nature.

(iii) The validity of (1.2) or (1.3) for non-global solutions is open.

7 Nonlocal problems

As already mentioned in the introduction, the estimate (1.2) can be derived for more general problems than (1.1). For example, in [31] we considered two nonlocal problems, which were frequently studied from the point of view of blow-up and global existence in the past decade (see the references in [31]). For both of these problems we derived the estimate (1.2) and the continuity of the blow-up time.

The first problem has the form

$$\begin{aligned} u_t - \Delta u &= f(x, u(x, t)) - \frac{1}{|\Omega|} \int_{\Omega} f(x, u(x, t)) \, dx, & x \in \Omega, \, t > 0, \\ u_\nu &= 0, & x \in \Gamma, \, t > 0, \\ u(x, 0) &= u_0(x), & x \in \Omega, \end{aligned}$$

where Ω is a smoothly bounded domain in \mathbb{R}^n and $f(x, \cdot)$ is a superlinear function (in particular, one can choose $f(x, u) = |u|^{p-1}u$, $p_S > p > 1$).

The second nonlocal problem has the form

$$\begin{aligned} u_t - \Delta u &= \phi \left(\int_{\Omega} F(u) \, dx \right) f(u), & x \in \Omega, \, t > 0, \\ u &= 0, & x \in \Gamma, \, t > 0, \\ u(x, 0) &= u_0(x), & x \in \Omega, \end{aligned}$$

where $f = F'$, Ω is a smoothly bounded domain in \mathbb{R}^n and either

$$F(u) = \frac{1}{p+1} |u|^{p+1}, \quad \phi(s) = (s+1)^{-\alpha}, \quad 1 < p < p_S, \quad 0 \leq \alpha < \frac{p-1}{p+1},$$

or

$$F(u) = e^u, \quad \phi(s) = s^{-q}, \quad 0 < q < 1, \quad n = 1.$$

8 Problems involving measures

Notice that the assumption (2.3) in Section 2 requires $f(\cdot, 0) \in L_{(p_2+1)/p_2}(\Omega)$ and that even a stronger assumption on the integrability of $f(\cdot, 0)$ is required in the second part of Theorem 2.1. If $f(\cdot, 0)$ is less regular then we can still expect similar results as in Theorem 2.1 provided we restrict the range for the exponent p_2 . Consider, for example, the model problem

$$u_t - \Delta u = |u|^{p-1}u + a\mu, \quad x \in \Omega, \quad t > 0, \quad (8.1)$$

$$u = 0, \quad x \in \Gamma, \quad t > 0, \quad (8.2)$$

$$u(x, 0) = u_0(x), \quad x \in \Omega, \quad (8.3)$$

where Ω is a bounded domain in \mathbb{R}^n , $n \geq 2$, μ is a positive bounded Radon measure on Ω , $a > 0$ and $1 < p$, $p(n-2) < n$. The restriction on p is necessary for the local solvability of (8.1).

It is known (see [3] or [1]) that

$$a^* := \sup\{a > 0 \text{ (8.1) has a positive equilibrium}\} > 0.$$

Set $X := \{u \in W_q^z(\Omega) \mid u = 0 \text{ on } \Gamma\}$, where

$$-\frac{n}{p} \leq z - \frac{n}{q} < 2 - n, \quad q > 1, \quad z \geq 0, \quad z \neq 1/q.$$

The following result from [29] is restricted to global solutions of (8.1), but we believe that a complete analogon to Theorem 2.1 can be proved.

Theorem 8.1. *Let $\Omega, n, p, \mu, a^*, X$ be as above and let $0 < a < a^*$. Let u be a global solution of (8.1). Then $\|u(t)\|_X \leq C(\|u_0\|_X)$.*

Let u_s be the minimal positive stationary solution of (8.1). Then there exist stationary solutions u_+, u_-, \tilde{u} of (8.1) such that $u_+ > u_s > u_-$ and the function $\tilde{u} - u_s$ changes sign.

Acknowledgement. The author was supported by VEGA Grant 1/7677/20.

References

1. H. Amann and P. Quittner, *Elliptic boundary value problems involving measures: existence, regularity, and multiplicity*, Advances in Diff. Equations **3** (1998), 753-813.
2. P. Baras and L. Cohen, *Complete blow-up after T_{max} for the solution of a semilinear heat equation*, J. Funct. Anal. **71** (1987), 142-174.
3. P. Baras and M. Pierre, *Critère d'existence de solutions positives pour des équations semi-linéaires non monotones*, Analyse Non Linéaire, Ann. Inst. H. Poincaré **2** (1985), 185-212.
4. M.-F. Bidaut-Véron, *Initial blow-up for the solutions of a semilinear parabolic equation with source term*, In: Equations aux dérivées partielles et applications, articles dédiés à Jacques-Louis Lions, Gauthier-Villars, Paris (1998), pp. 189-198.

5. H. Brézis and R.E.L. Turner, *On a class of superlinear elliptic problems*, Commun. Partial Differ. Equations **2** (1977), 601-614.
6. T. Cazenave and P.-L. Lions, *Solutions globales d'équations de la chaleur semi-linéaires*, Commun. Partial Differ. Equations **9** (1984), 955-978.
7. M. Chipot, M. Fila and P. Quittner, *Stationary solutions, blow up and convergence to stationary solutions for semilinear parabolic equations with nonlinear boundary conditions*, Acta Math. Univ. Comenianae **60** (1991), 35-103.
8. M.J. Esteban, *On periodic solutions of superlinear parabolic problems*, Trans. Amer. Math. Soc. **293** (1986), 171-189.
9. M.J. Esteban, *A remark on the existence of positive periodic solutions of superlinear parabolic problems*, Proc. Amer. Math. Soc. **102** (1988), 131-136.
10. C. Fermanian Kammerer, F. Merle, H. Zaag, *Stability of the blow-up profile of nonlinear heat equations from the dynamical system point of view*, Math. Ann. **317** (2000), 347-387.
11. D.G. de Figueiredo, P.-L. Lions and R.D. Nussbaum, *A priori estimates and existence of positive solutions of semilinear elliptic equations*, J. Math. pures et appl. **61** (1982), 41-63.
12. M. Fila, *Boundedness of global solutions of nonlinear parabolic problems*, Proc. of the 4th European Conf. on Elliptic and Parabolic Problems, Rolduc 2001 (to appear).
13. M. Fila and P. Poláčik, *Global solutions of a semilinear parabolic equation*, Adv. Differ. Equations **4** (1999), 163-196.
14. M. Fila, P. Souplet and F. Weissler, *Linear and nonlinear heat equations in L^q_δ spaces and universal bounds for global solutions*, Math. Ann. **320** (2001), 87-113.
15. S. Filippas, M.A. Herrero and J.J.L. Velázquez, *Fast blow-up mechanism for sign-changing solutions of a semilinear parabolic equation with critical nonlinearity*, R. Soc. Lond. Proc. Ser. A **456** (2000), 2957-2982.
16. V. Galaktionov and J.L. Vázquez, *Continuation of blow-up solutions of nonlinear heat equations in several space dimensions*, Commun. Pure Applied Math. **50** (1997), 1-67.
17. B. Gidas and J. Spruck, *A priori bounds for positive solutions of nonlinear elliptic equations*, Commun. Partial Differ. Equations **6** (1981), 883-901.
18. Y. Giga, *A bound for global solutions of semilinear heat equations* Commun. Math. Phys. **103** (1986), 415-421.
19. Y. Giga and R.V. Kohn, *Characterizing blowup using similarity variables*, Indiana Univ. Math. J. **36** (1987), 1-40.
20. M.A. Herrero and J.J.L. Velázquez, *Explosion de solutions d'équations paraboliques semilinéaires supercritiques*, C. R. Acad. Sci. Paris Sér. I Math. **319** (1994), 141-145.
21. M.A. Herrero and J.J.L. Velázquez, *A blow up result for semilinear heat equations in the supercritical case*, Preprint.
22. J. Hůska, *Periodic solutions in superlinear parabolic problems*, Acta Math. Univ. Comenianae (to appear).
23. H.A. Levine, *Some nonexistence and instability theorems for solutions of formally parabolic equations of the form $Pu_t = -Au + F(u)$* , Arch. Rational Mech. Anal. **51** (1973), 371-386.
24. J. Matos, Ph. Souplet, *Universal blow-up estimates and decay rates for a semilinear heat equation*, Preprint.
25. W.-M. Ni, P.E. Sacks and J. Tavantzis, *On the asymptotic behavior of solutions of certain quasilinear parabolic equations*, J. Differ. Equations **54** (1984), 97-120.
26. R.D. Nussbaum, *Positive solutions of nonlinear elliptic boundary value problems*, J. Math. Anal. Appl. **51** (1975), 461-482.

27. S.I. Pohozaev *Eigenfunctions of the equation $\Delta u + \lambda f(u) = 0$* , Soviet Math. Dokl. **5** (1965), 1408-1411.
28. P. Quittner, *A priori bounds for global solutions of a semilinear parabolic problem*, Acta Math. Univ. Comenianae **68** (1999), 195-203.
29. P. Quittner, *A priori estimates of global solutions and multiple equilibria of a superlinear parabolic problem involving measure*, Electronic J. Differ. Equations **29** (2001), 1-17.
30. P. Quittner, *Universal bound for global positive solutions of a superlinear parabolic problem*, Math. Ann. **320** (2001), 299-305.
31. P. Quittner, *Continuity of the blow-up time and a priori bounds for solutions in superlinear parabolic problems*, Houston J. Math. (to appear).
32. P. Quittner, *Multiple equilibria, periodic solutions and a priori bounds for solutions in superlinear parabolic problems*, NoDEA, Nonlinear Differ. Equations Appl. (to appear).
33. P. Quittner and Ph. Souplet, *A priori estimates of global solutions of superlinear parabolic problems without variational structure*, Discrete Contin. Dyn. Systems (to appear).
34. P. Quittner and Ph. Souplet, *Bounds of solutions of parabolic problems with nonlinear boundary conditions*, In preparation.
35. P. Quittner, Ph. Souplet and M. Winkler, *Initial blow-up rates and universal bounds for nonlinear heat equations*, Preprint.
36. R.E.L. Turner, *A priori bounds for positive solutions of nonlinear elliptic equations in two variables*, Duke Math. J. **41** (1974), 759-774.
37. H. Zaag, *A remark on the energy blow-up behavior for nonlinear heat equations*, Duke Math. J. **103** (2000), 545-556.