

EQUADIFF 10

Arina A. Arkhipova

Solvability problem for strong-nonlinear nondiagonal parabolic system

In: Miroslav Krbeč and Jaromír Kuben (eds.): Proceedings of Equadiff 10, Czechoslovak International Conference on Differential Equations and Their Applications, Prague, August 27-31, 2001, [Part 1] Invited Lectures. Masaryk University, Brno, 2002. CD-ROM issued as a complement to the journal edition *Mathematica Bohemica* 2002/2. pp. 7--14.

Persistent URL: <http://dml.cz/dmlcz/700327>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Solvability Problem for Strong-Nonlinear Nondiagonal Parabolic System

Arina A. Arkhipova

St.Petersburg State University
Department of Mathematics and Mechanics
Bibliotechnaya pl., 2, Stary Petergof
198504, St. Petersburg, Russia
Email: arina@AA1101.spb.edu

Abstract. A class of q -nonlinear parabolic systems with nondiagonal principal matrix and strong nonlinearities in the gradient is considered. We discuss the global in time solvability results of the classical initial boundary value problems in the case of two spatial variables. The systems with nonlinearities $q \in (1, 2)$, $q = 2$, $q > 2$, are analyzed.

MSC 2000. 35K50, 35K45, 35K60

Keywords. Boundary value problems, nonlinear parabolic systems, solvability

Let Ω be a bounded domain in R^n , $n \geq 2$, with sufficiently smooth boundary. For a fixed $T > 0$ and $Q = \Omega \times (0, T)$, we consider a solution $u: Q \rightarrow R^N$, $u = (u^1, \dots, u^N)$, $N > 1$, of the parabolic system

$$u_t^k - \frac{d}{dx_\alpha} a_\alpha^k(z, u, u_x) + b^k(z, u, u_x) = 0, \quad z = (x, t) \in Q, \quad k = 1, \dots, N. \quad (1)$$

We define the set $D = \overline{Q} \times R^N \times R^{nN}$ and assume that

- the functions $a = \{a_\alpha^k\}_{\alpha \leq n}^{k \leq N}$ and $b = \{b^k\}^{k \leq N}$ are sufficiently smooth on D ;
- for a fixed $q > 1$ $a(\cdot, \cdot, p) \sim |p|^{q-1}$, $b(\cdot, \cdot, p) \sim |p|^q$, $|p| \gg 1$, all derivatives of a and b that we need have the natural growth with respect to the gradient;

- c) nondiagonal principal matrix $\left\{ \frac{\partial a_\alpha^k}{\partial p_\beta^l} \right\}_{k,l \leq N}^{\alpha,\beta \leq n}$ satisfies the following assumptions on D :

$$\frac{\partial a_\alpha^k(z, u, p)}{\partial p_\beta^l} \xi_\alpha^k \xi_\beta^l \geq \nu(1 + |p|)^{q-2} |\xi|^2, \quad \left| \frac{\partial a(\dots)}{\partial p} \right| \leq \mu(1 + |p|)^{q-2}, \quad \forall \xi \in R^{nN}; \quad (2)$$

- d) strong-nonlinear term b satisfies the condition

$$|b(z, u, p)| \leq b_0(1 + |p|)^q, \quad (z, u, p) \in D. \quad (3)$$

Here $\nu, \mu, b_0 = \text{const} > 0$.

We investigate solvability of the Cauchy-Dirichlet problem

$$u|_{\partial'Q} = \varphi, \quad (4)$$

where $\partial'Q$ is the parabolic boundary of Q , and φ is a given smooth function.

First of all, we recall some known results.

We fix the class

$$V = L_q((0, T), W_q^1(\Omega)) \cap L_\infty(Q)$$

and note that the global solvability of (1), (4) in V was stated for the scalar situation ($N = 1$) in the following sense. Assume that for a fixed $M > 0$ an a priori estimate

$$\|u\|_{\infty, Q} \leq M \quad (5)$$

can be derived. Then there exists a solution $u \in V \cap C^\alpha(\bar{Q})$ with some $\alpha \in (0, 1)$. Further regularity of the solution follows provided that all the data are smooth enough [1].

In some sense this result is also valid for a class of quasilinear *diagonal* systems ($N > 1, q = 2$). More precisely, if estimate (5) and the ‘‘smallness’’ condition $b_0 M < \nu$ hold then a solution u of (1), (4) exists in $V \cap C^\alpha(\bar{Q})$.

It should be remarked that due to the maximum principle we are able to formulate sufficient conditions, which provides estimate (5) in the cases mentioned above.

Now, let us consider the parabolic (elliptic) system with *nondiagonal* principal matrix. In this situation, the following questions arise: i) how to guarantee estimate (5)? ii) is the class V suitable for proving global solvability of (1), (4)?

Under conditions a)–d), the global solvability problem for (1), (4) has not been solved yet.

Certainly, we can not expect classical global solvability of this problem. As is known, there are counterexamples of the regularity for quasilinear nondiagonal systems even if $b \equiv 0$ ($q = 2, n > 2$) [2]. From the other hand, for systems (1) whose main part is the heat operator, but term $b(z, u, p)$ is non-zero and satisfies (3), singularities can appear in Q in some time. The heat flow of harmonic maps gives us an example of such a situation (see, for example, [3], [4]).

From the above, it follows that there are two reasons that cause nonsmoothness of solutions of the problem under consideration.

In recent years, the author investigated the global solvability for (1), (4) under assumptions a)–d) in the following particular case.

To put the problem, we define the functional

$$E[u] = \int_{\Omega} f(x, u, u_x) dx, \quad u = (u^1, \dots, u^N), \quad N > 1, \quad (6)$$

and denote by $L = \{L^k\}^{k \leq N}$ the Euler operator of E :

$$L^k u = -\frac{d}{dx_{\alpha}} f_{p_{\alpha}^k} + f_{u^k}.$$

Then system (1) is the gradient flow for the functional E . Consider the problem

$$\begin{aligned} u_t^k - \frac{d}{dx_{\alpha}} f_{p_{\alpha}^k}(x, u, u_x) + f_{u^k}(x, u, u_x) &= 0, \quad (x, t) \in Q, \quad k \leq N, \\ u|_{\Gamma} &= 0, \quad u|_{t=0} = \varphi_0(x), \end{aligned} \quad (7)$$

where $\Gamma = \partial\Omega \times (0, T)$.

The variational structure of system (7) provides an apriori estimate of solution u :

$$\|u_t\|_{2, Q}^2 + \sup_{(0, T)} \|u_x(\cdot, t)\|_{q, \Omega}^q \leq \epsilon_0, \quad (8)$$

$\epsilon_0 = \text{const}$ depends on the data only.

Moreover, this structure also ensures monotonicity of the global energy

$$E[u(\cdot, t_1)] \leq E[u(\cdot, t_2)], \quad \forall t_1 > t_2,$$

and a local energy estimate

$$\|u_t\|_{2, \mathcal{P}_R(z^0)}^2 + \sup_{\lambda_R(t^0)} \|u_x(\cdot, t)\|_{q, \Omega_R(x^0)}^q \leq \frac{c}{R^q} \int_{\mathcal{P}_{2R}(z^0)} (1 + |u_x|)^q dP. \quad (9)$$

In (9) and below, we denote

$$\begin{aligned} \mathcal{P}_R(z^0) &= Q_R(z^0) \cap Q, \quad Q_R(z^0) = B_R(x^0) \times \lambda_R(t^0), \\ B_R(x^0) &= \{x \in \mathbb{R}^n \mid |x - x^0| < R\}, \quad \lambda_R(t^0) = (t^0 - R^q, t^0 + R^q), \\ \Omega_R(x^0) &= B_R(x^0) \cap \Omega. \end{aligned}$$

We say that $Q_R(z^0)$ is a q -parabolic cylinder and denote by

$$\delta_q(z^1, z^2) = \sup \{|x^1 - x^2|, |t^1 - t^2|^{1/q}\}, \quad \forall z^1, z^2 \in \mathbb{R}^{n+1}, \quad (10)$$

q -parabolic distance in \mathbb{R}^{n+1} .

To introduce an example of system (7), we put

$$f(x, u, p) = \langle A(x, u)p, p \rangle (l + |p|)^{q-2}, \quad q > 1, \quad (11)$$

in the definition (6) of E . We assume that $A(\cdot, \cdot)$ is a nondiagonal positive definite and smooth matrix on $\bar{\Omega} \times \mathbb{R}^N$, and, in addition, $A_{kl}^{\alpha\beta} = A_{lk}^{\beta\alpha}$. Generated by function (11) system (7) satisfies conditions a)–d), in the particular, $f_u(\cdot, \cdot, p) \sim |p|^q$, $|p| \gg 1$.

Let us now proceed to discussing some solvability results recently proved by the author.

We stated some solvability results for problem (7) in the case of two spatial variables.

First, we considered problem (7) with $n = q = 2$. We analyzed it with quasilinear and nonlinear operators under Dirichlet or Neumann type conditions ([5]–[8]). For all these situations the following result was proved.

Theorem 1. *For a fixed number $T > 0$, there exists a global solution of (7), which is almost everywhere smooth in \bar{Q} . The singular set consists of at most finitely many points. The solution u has finite norms (6), and it is a weak solution in the sense of distributions.*

This result was proved with the help of the continuability theorem of smooth solutions from a semiclosed time interval. We essentially exploited the imbedding theorems for two dimensional domains and the fact that “local normalized energy”

$$\frac{1}{R^{n-q}} \int_{B_R(x^0)} |u_x(x, t)|^q dx$$

is a monotone function of R if $n = q = 2$.

For the case $n = 2$ and $q > 2$, we prove the following result.

Theorem 2. *Let $q > 2$, $n = 2$, and T be a positive fixed number. There exists a smooth solution of problem (7) in \bar{Q} if all the data are sufficiently smooth.*

Now, we give a sketch of the proof of this result. We start from the derivation of some apriori estimates for solutions u of (7) smooth on time interval $[0, T)$.

First of all, from (8) one can deduce an apriori estimate

$$\|u\|_{C^\gamma(\bar{Q}, \delta_q)} \leq \text{const} \quad (12)$$

where γ is some number in $(0, 1)$.

Estimate (12) allows us to derive apriori estimates of stronger norms of u in \bar{Q} .

From this point, we study problem (7) in a local setting. Let $v(y, t) = u(x(y), t)$ be a solution of the problem

$$v_t - \frac{d}{dy_\alpha} A_\alpha^k(y, v, v_y) + \mathbb{B}^k(y, v, v_y) = 0, \quad (y, t) \in Q_2^+, \quad k, \dots, N, \quad (13)$$

$$v|_{t=0} = \varphi_0(x(y)), \quad v|_{\Gamma_2^+} = 0,$$

where $Q_R^+ = B_R^+(0) \times (0, T)$, $B_R^+(0) = B_R(0) \cap \{y_2 > 0\}$, $\Gamma_R^+ = \gamma_R(0) \times (0, T)$, and $\gamma_R(0) = B_R(0) \cap \{y_2 = 0\}$. On the set $D = Q_2^+ \times \mathbb{R}^N \times \mathbb{R}^{2N}$, functions A_α^k and \mathbb{B}^k satisfy conditions (2) and (3) with some other constants.

With the help of (12), one can derive the inequality

$$\begin{aligned} & \sup_{\lambda_R(t^0)} \int_{\Omega_R(y^0)} |v_y(y, t)|^2 dy + \int_{\mathcal{P}_R(z^0)} [(1 + |v_y|)^{q-2} |v_{yy}|^2 + \\ & (1 + |v_y|)^{q+2}] d\mathcal{P} \leq cR^{2\alpha}, \quad \forall R \leq R_0, \quad z^0 = (y^0, t^0) \in Q_{3/2}^+, \end{aligned} \quad (14)$$

with some $\alpha \in (0, 1)$ and $R_0 > 0$.

Next, we state that for some $s \in (0, 1)$

$$\sup_{(0, T)} \|v_t(\cdot, t)\|_{2+2s, B_1^+(0)} \leq c_1. \quad (15)$$

Here and below, $c_i, i = 1 \dots, 4$, are positive constants depending on the data only. To derive (15), we use (14) estimating strong-nonlinear terms generated by functions \mathbb{B}^k . After that, we are able to look at our problem as at the elliptic one for a fixed $t \in (0, T)$.

The reverse Hölder inequalities hold for

$$V(\cdot, t) = (1 + |v_y(\cdot, t)|)^{\frac{q-2}{2}} |v_{yy}(\cdot, t)|$$

in $B_{\frac{1}{2}}^+(0)$.

Due to the Gehring Lemma, we have the estimate of $\|V(\cdot, t)\|_{p, B_{\frac{1}{2}}^+(0)}$ with some $p > 2$. As a consequence, we come to the estimates

$$\sup_{(0, T)} \|v_{yy}(\cdot, t)\|_{p, B_{\frac{1}{2}}^+(0)} \leq c_2, \quad \sup_{(0, T)} \|v_y(\cdot, t)\|_{C^\beta(\overline{B_{\frac{1}{2}}^+(0)})} \leq c_3, \quad \beta = 1 - 2/p > 0. \quad (16)$$

Estimates (12), (16) guarantee us that

$$\|u_x\|_{C^{\beta_0}(\overline{Q}; \delta_q)} \leq c_4$$

with some $\beta_0 > 0$.

A priori estimates of the stronger norms of u up to $t = T$ follows from the linear theory. It means that u can be extended as a smooth function up to $t = T$.

Due to the known solvability results, there exists a smooth solution u of (7) on some time interval $[0, T_0)$. Let T_0 defines the maximal interval of the existence of smooth solution u . Suppose that $T_0 < T$. As it was explained above, u can be extended as a smooth function up to $t = T_0$. Thus, one comes to the contradiction with the definition of T_0 . From all said above, it follows that $T_0 \geq T$. Theorem 2 is proved.

A more complicated case is $n = 2, q \in (1, 2)$. To prove the solvability of (7), we introduce approximate problems

$$\begin{aligned} & u_t + Lu - \varepsilon \Delta u = 0 \quad \text{in } Q, \\ & u|_{\Gamma} = 0, \quad u|_{t=0} = \varphi^\varepsilon, \quad \varepsilon \in (0, 1]. \end{aligned} \quad (17)$$

Here φ^ε is an approximation of φ , φ^ε satisfies the compatibility conditions for system (17), φ^ε goes to φ in the strong sense if $\varepsilon \rightarrow 0$.

For a fixed $\varepsilon > 0$, we prove global classical solvability of (17) in the space $\mathcal{H}^{2+\alpha, 1+\alpha/2}(\bar{Q})$ with Hölder exponent $\alpha \in (0, 1)$ (the definition of this space see in [1, Ch. I, § 1]). Certainly, the norm of the solutions u^ε of (17) in this space goes to infinity if $\varepsilon \rightarrow 0$. We are able to estimate different norms of u^ε due to the fact that for a fixed $\varepsilon > 0$ the Laplace operator forms the main part of the elliptic operator of system (17), and functions $f_u(x, u, u_x)$ are not the strong-nonlinear terms with respect to the Laplace operator. Also, it is worth noting, that all estimates of u^ε we derive in the standard cylinders ($q = 2$ in the definition of Q_R).

The sequence u^ε goes in some sense to the limit function u if $\varepsilon \rightarrow 0$, and u is a solution of (7). More exactly, the following fact was proved.

Theorem 3. *Let $q \in (1, 2)$ and T be a fixed positive number. There exists a solution u of problem (7), which is almost everywhere smooth in \bar{Q} ; $u_t \in L_2(Q)$, and $u \in L_\infty((0, T); \dot{W}_q^1(\Omega))$. The closed singular set Σ of u has $\dim_{q-\mathcal{H}} \Sigma \leq 2$ ($\dim_{\mathcal{H}} \Sigma \leq 4 - q$). Moreover, $\dim_H \Sigma^\tau \leq 2 - q$, $\forall \tau > 0$, where $\Sigma^\tau = \Sigma \cap \{t = \tau\}$.*

In the statement of Theorem 3, the estimate $\dim_{q-\mathcal{H}} \Sigma \leq 2$ means that for all $\eta > 0$, $\mathcal{H}_{2-q+\eta}(\Sigma; \delta_q) = 0$, where δ_q -parabolic metric is defined in (10).

Now, we explain the main steps of the proof of Theorem 3.

Lemma 4. *There exists a number ω_0 depending on the data only, such that if*

$$\omega_{R_0}^{\varepsilon_j}(z^0) \equiv \rightarrow_{\mathcal{P}_{R_0}(z^0)} \text{osc } u^{\varepsilon_j} \leq \omega_0 \quad (18)$$

for a point $z^0 \in \bar{Q}$ with some $R_0 > 0$ and a sequence $\varepsilon_j \rightarrow 0$, then

$$\|u^{\varepsilon_j}\|_{C^{\gamma_1}(\overline{\mathcal{P}_{R_*}(z^0)}; \delta_q)} + \|u_x^{\varepsilon_j}\|_{C^{\gamma_2}(\overline{\mathcal{P}_{R_*}(z^0)}; \delta_q)} \leq c_0, \quad (19)$$

with some $\gamma_1, \gamma_2 \in (0, 1)$ and $R_* = R_*(R_0, \omega_0) < R_0$, u^{ε_j} is a solution of (17).

It should be remarked that, in general, (in the case of nondiagonal matrix and condition (3)) the smallness of the oscillation of a solution does not guarantee an estimate of the Hölder norm of the solution.

It is evident that condition (18) provides smoothness of the solution u at the point z_0 .

Next step is to introduce an integral description of a regular point of u .

Lemma 5. *Suppose that for a point $z^0 \in \bar{Q}$ there exist numbers $\mathcal{K} > 0$, $\beta > 1$, $R_0 > 0$ and a sequence $\varepsilon_j \rightarrow 0$ such that*

$$\sup_{\hat{z} \in \mathcal{P}_{R_0}(z^0)} \sup_{R \leq R_0} \frac{(\log_2 \frac{2R_0}{R})^{\frac{\beta q^2}{2(q-1)}}}{R^2} \int_{\mathcal{P}_R(\hat{z})} H_{\varepsilon_j}^2 dz \leq \mathcal{K}, \quad (20)$$

where $H_\varepsilon^2 = (1 + |u_x^\varepsilon|)^q + \varepsilon |u_x^\varepsilon|^2$. Then estimate (18) holds in $\mathcal{P}_{R_1}(z^0)$ with some $R_1 < R_0$.

To derive (18) from (20), we exploit a local energy estimate for solutions u^ε of (17). We also use a certain condition for functions from Sobolev space $W_q^1(\Omega)$, $n = 2$, that makes it possible to estimate their oscillation. To prove (18), we analyze both cases $\varepsilon_j < \frac{R^{2-q}}{\chi(R)}$ and $\varepsilon_j \geq \frac{R^{2-q}}{\chi(R)}$, where $\chi(R) = \frac{\mathcal{K} \frac{2-q}{q}}{(\log_2 \frac{2R_0}{R})^\gamma}$, $\gamma = \frac{q(2-q)\beta}{2(q-1)}$, and $R \leq R_0$. Next, we denote by \mathcal{R} the set of all points z_0 in \bar{Q} , where (20) holds with some parameters \mathcal{K} , β , R_0 , and $\{\varepsilon_j\}_{j \in N}$, $\varepsilon_j \rightarrow 0$, and put $\Sigma = \bar{Q} \setminus \mathcal{R}$.

We have the following description of Σ :

A point z^0 belongs to Σ if for all $M_k \rightarrow \infty$ and $R_k \rightarrow 0$, there exist sequences of points $\xi_k \in \mathcal{P}_{R_k}(z^0)$ and numbers $\rho_k \leq R_k$ such that

$$\liminf_{\varepsilon \rightarrow 0} \frac{\left(\log_2 \frac{2R_k}{\rho_k}\right)^\gamma}{\rho_k^2} \int_{\mathcal{P}_{\rho_k}(\xi_k)} H_\varepsilon^2 dz > M_k, \quad (21)$$

where $\gamma = \frac{\beta q^2}{2(q-1)}$.

The relation (21) does not allow us to estimate the Hausdorff measure of Σ and, therefore, instead of (21), we prove that one can exploit the following description of $z^0 \in \Sigma$:

for arbitrary sequences $M_k \rightarrow \infty$ and $R_k \rightarrow 0$, there exist a sequence of numbers $r_k \leq 2R_k$ and an absolute number $c_* > 1$ such that

$$\liminf_{\varepsilon \rightarrow 0} \frac{\left(\log_2 \frac{4R_k}{r_k}\right)^\gamma}{r_k^2} \int_{\mathcal{P}_{r_k}(z^0)} H_\varepsilon^2 dz > \frac{M_k}{c_*}. \quad (22)$$

From (22), we deduce an estimate of the Hausdorff dimension of the set Σ as it was pointed in Theorem 3.

It should be noted that the presence of the logarithm multiplier in (22) does not allow us to assert that

$$\mathcal{H}_2(\Sigma; \delta_q) < +\infty \quad (\mathcal{H}_{4-q}(\Sigma; \delta) < +\infty, \delta = \delta_2) \quad \text{and} \quad H_{2-q}(\Sigma^\tau) < +\infty, \forall \tau > 0. \quad (23)$$

If (23) would be proved, then we could pass to the limit in the integral identity corresponding to problem (17) and state that the limit function u is a weak solution of (22) in the sense of distributions.

References

1. Ladyzhenskaya O.A., Solonnikov V.A., Uraltseva N.N. *Linear and Quasilinear Equations of Parabolic Type*. Amer. Math. Soc., Providence, 1968.
2. Stará J., John O. *Some (new) counterexamples of parabolic systems*. Comment. Math. Univ. Carolinae, **36**, 3, (1995), 503–510.
3. Chen Y., Struwe M. *Existence and Partial Regularity Results for the Heatflow for Harmonic Maps*. Math. Z., **201**, (1989), 83–103.
4. Chang K.-C. *Heat Flow and Boundary Value Problem for Harmonic Maps*. Ann. Inst. Henri Poincaré, **6**, 5 (1989), 363–395.

5. Arkhipova A. *Global Solvability of the Cauchy-Dirichlet Problem for Nondiagonal Parabolic Systems with Variational Structure in the Case of Two Spatial Variables*. Probl. Mat. Anal. **16**, St. Petersburg Univ., (1997), 3–40; English transl. J. Math Sci., **92**, 6 (1998), 4231–4255.
6. Arkhipova A. *Local and Global Solvability of the Cauchy-Dirichlet Problem for a Class of Nonlinear Nondiagonal Parabolic Systems*. St. Petersburg Math. J., **11**, 6 (2000), 989–1017.
7. Arkhipova A. *Cauchy-Neumann Problem for a Class of Nondiagonal Parabolic Systems with Quadratic Growth Nonlinearities. I. On the Continuability of Smooth Solutions*. Comment. Math Univ. Carolinae, **41**, 4 (2000), 693–718.
8. Arkhipova A. *Cauchy-Neumann Problem for a Class of Nondiagonal Parabolic Systems with Quadratic Growth Nonlinearities. II. Local and Global Solvability Results*. Comment. Math. Univ. Carolinae, **42**, 1 (2001), 53–76.