

EQUADIFF 10

Ondřej Došlý

Qualitative theory of half-linear second order differential equations

In: Miroslav Krbeč and Jaromír Kuben (eds.): Proceedings of Equadiff 10, Czechoslovak International Conference on Differential Equations and Their Applications, Prague, August 27-31, 2001, [Part 1] Invited Lectures. Masaryk University, Brno, 2002. CD-ROM issued as a complement to the journal edition *Mathematica Bohemica* 2002/2. pp. 37--50.

Persistent URL: <http://dml.cz/dmlcz/700330>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Qualitative theory of half-linear second order differential equations

Ondřej Došlý*

Mathematical Institute
Academy of Sciences of the Czech Republic
Žižkova 22, 616 62 Brno
Czech Republic
Email: dosly@math.muni.cz

Abstract. Some recent results concerning properties of solutions of the half-linear second order differential equation

$$(r(t)\Phi(x'))' + c(t)\Phi(x) = 0, \quad \Phi(x) := |x|^{p-2}x, \quad p > 1, \quad (*)$$

are presented. A particular attention is devoted to the oscillation theory of (*). Related problems are also discussed.

MSC 2000. 345C10

Keywords. Half-linear equation, Picone's identity, scalar p -Laplacian, variational method, Riccati technique, principal solution

Dedicated to the memory of Prof. Árpád Elbert.

1 Introduction

In this contribution we deal with the oscillatory properties and related problems concerning the half-linear second order differential equation

$$(r(t)\Phi(x'))' + c(t)\Phi(x) = 0, \quad \Phi(x) := |x|^{p-2}x, \quad p > 1, \quad (1)$$

* Research supported by the Grant A1019902 of the Grant Agency of the Czech Academy of Sciences.

where r, c are continuous functions and $r(t) > 0$. In the recent years it was shown that solutions of (1) behave in many aspects like those of the Sturm-Liouville equation

$$(r(t)x')' + c(t)x = 0 \quad (2)$$

which is the special case $p = 2$ of (1). The aim of this paper is to present some results of this investigation and also to point out situations where properties of (1) and (2) (considerably) differ. Note that terminology half-linear equations is motivated by the fact that the solution space of (1) has just one half of the properties which characterize linearity, namely homogeneity (but not additivity).

The investigation of qualitative properties of nonlinear second order differential equations has a long history. Recall here only the papers of Emden [26], Fowler [27], Thomas [40], and the book of Sansone [39] containing the survey of the results achieved in the first half of the last century. In the fifties and later decades the number of papers devoted to nonlinear second order differential equations increased rapidly, so we mention here only treatments directly associated with (1). Even if some ideas concerning the properties of solutions of (1) can be already found in the papers of Bihari [4,5], as pioneers of the qualitative theory of (1) are usually regarded Elbert and Mirzov with their papers [21,36]. In the following years, in particular in the nineties, the striking similarity between oscillatory properties of (1) and (2) was revealed. On the other hand, in some aspects, e.g. the Fredholm-type alternative for solutions of boundary value problems associated with (1), it turned out that the situation is completely different in linear and half-linear case and the absence of the additivity of the solution space of (1) brings completely new phenomena.

The paper is organized as follows. In the next section we present a brief survey of basic properties of solutions of (1). Section 3 is devoted to the oscillation theory of (1) and in the last section we discuss some other problems associated with (1).

2 Basic properties of (1)

Consider a special equation of the form (1)

$$(\Phi(x'))' + (p-1)\Phi(x) = 0$$

and denote by $S = S(t)$ its solution given by the initial condition $S(0) = 0$, $S'(0) = 1$. In [21] it is shown that the behaviour of this function is very similar to that of the classical sine function. In particular, this function is odd, periodic with the half-period $\pi_p := \frac{2\pi}{p \sin(\pi/p)}$, and satisfies the generalized Pythagorean identity $|S(t)|^p + |S'(t)|^p = 1$.

Using this function, one can introduce the generalized Prüfer transformation as follows. Let x be a nontrivial solution of (1). There exist differentiable functions ρ, φ such that

$$x(t) = \rho(t)S(\varphi(t)), \quad r^{q-1}(t)x'(t) = \rho(t)S'(\varphi(t)),$$

where q is the conjugate number of p , i.e., $\frac{1}{p} + \frac{1}{q} = 1$, and the functions ρ, φ satisfy a certain first order system with Lipschitzian right-hand-side, which is, in turn, uniquely solvable. Hence, the same holds for (1): given $t_0 \in \mathbb{R}$ and $A, B \in \mathbb{R}$, there exists a unique solution of (1) satisfying $x(t_0) = A, x'(t_0) = B$ which is extensible over the whole interval where r, c are continuous and $r(t) > 0$.

Other important objects associated with (1) are the p -degree functional

$$\mathcal{F}(y; a, b) := \int_a^b [r(t)|y'|^p - c(t)|y|^p] dt \tag{3}$$

(equation (1) is the Euler-Lagrange equation of \mathcal{F} considered in the class of functions satisfying the zero boundary condition $y(a) = 0 = y(b)$) and the generalized Riccati equation

$$w' + c(t) + (p - 1)r^{1-q}(t)|w|^q = 0, \quad q = \frac{p}{p - 1}, \tag{4}$$

which is related to (1) by the substitution $w = \frac{r\Phi(x')}{\Phi(x)}$.

Functional (3) and equation (4) are related by the half-linear version of Picone's identity

$$\mathcal{F}(y; a, b) = w(t)|y|_a^b + p \int_a^b r^{1-q}(t)P(r^{q-1}(t)y', w(t)\Phi(y)) dt,$$

where

$$P(u, v) := \frac{|u|^p}{p} - uv + \frac{|v|^q}{q} \geq 0$$

for all $u, v \in \mathbb{R}$ with the equality if and only if $v = \Phi(u)$, and w is a solution of (4) defined in the whole interval $[a, b]$. This identity is the main tool in the proof of the next statement which summarizes the basic oscillatory properties of (1). This statement is usually referred as the Roundabout Theorem.

Proposition 1. *The following statements are equivalent:*

- (i) Equation (1) is disconjugate on an interval $I = [a, b]$, i.e., any nontrivial solution of (1) has at most one zero in I .
- (ii) There exists a solution of (1) having no zero in $[a, b]$.
- (iii) There exists a solution w of the generalized Riccati equation (4) which is defined on the whole interval $[a, b]$.
- (iv) The p -degree functional $\mathcal{F}(y; a, b)$ is positive for every $0 \neq y \in W_0^{1,p}(a, b)$.

Observe that Proposition 1 implies that Sturm separation and comparison theorems extend verbatim to (1). Indeed, the separation theorem is essentially the equivalence (i) \iff (ii), while the comparison theorem is hidden in the equivalence (i) \iff (iv). In particular, similarly as in the linear case, equation (1) can be classified as oscillatory or nonoscillatory according to whether any nontrivial solution has or has not infinitely many zeros tending to ∞ .

Finally, let us mention at least two differences in the basic properties of solutions (1) (in addition to the already mentioned absence of the additivity of the solution space). If x_1, x_2 is a pair of linearly independent solutions of (1), we have no analogue of the Wronskian-type identity $r(x_1'x_2 - x_1x_2') = \text{const}$ which holds for (2), see [22]. We have also no half-linear analogue of the linear transformation identity with $x = h(t)y$

$$h(t)[(r(t)x')' + c(t)x] = (r(t)h^2(t)y')' + h(t)[(r(t)h'(t))' + c(y)h(t)]y \quad (5)$$

which is the starting point of the transformation theory of (2), see [6].

3 Oscillation theory

The equivalences given in the Roundabout Theorem (Proposition 1) suggest two main methods of the oscillation theory of (1). The basic ideas and results based on them are briefly explained in this section.

3.1 Variational method

This method is based on the equivalence (i) \iff (iv) in Proposition 1. According to this equivalence, to prove that (1) is oscillatory, it suffices to construct (for any $T \in \mathbb{R}$) a nontrivial function $y \in W_0^{1,p}(T, \infty)$ such that

$$\mathcal{F}(y; T, \infty) := \int_T^\infty [r(t)|y'|^p - c(t)|y|^p] dt \leq 0. \quad (6)$$

On the other hand, for nonoscillation of (1) we need to show the existence of $T \in \mathbb{R}$ such that $\mathcal{F}(y; T, \infty) > 0$ for every $0 \neq y \in W_0^{1,p}(T, \infty)$.

In oscillation criteria, a typical construction of a function y for which (6) holds reads as follows. Let T be arbitrary, $T < t_0 < t_1 < t_2 < t_3$, and let

$$y(t) = \begin{cases} 0, & t \in [T, t_0], \\ f(t), & t \in [t_0, t_1], \\ 1, & t \in [t_1, t_2], \\ g(t), & t \in [t_2, t_3], \\ 0, & t \in [t_3, \infty), \end{cases} \quad (7)$$

where f, g are solutions of the one-term equation

$$(r(t)\Phi(x'))' = 0 \quad (8)$$

satisfying $f(t_0) = 0$, $f(t_1) = 1$, $g(t_2) = 1$, $g(t_3) = 0$, i.e.

$$f(t) = \left(\int_{t_0}^t r^{1-q}(s) ds \right) \left(\int_{t_0}^{t_1} r^{1-q}(s) ds \right)^{-1},$$

$$g(t) = \left(\int_t^{t_3} r^{1-q}(s) ds \right) \left(\int_{t_2}^{t_3} r^{1-q}(s) ds \right)^{-1}.$$

By a direct computation, using the second mean value theorem of integral calculus applied to the integrals $\int_{t_0}^{t_1} r|f|^p$, $\int_{t_2}^{t_3} r|g|^p$ (see e.g. [14]) we have

$$\mathcal{F}(y; T, \infty) = \left(\int_{t_0}^{t_1} r^{1-q}(s) ds \right)^{1-p} + \left(\int_{t_2}^{t_3} r^{1-q}(s) ds \right)^{1-p} + \int_{s_1}^{s_2} c(t) dt, \quad (9)$$

where $s_1 \in (t_0, t_1)$, $s_2 \in (t_2, t_3)$. Using this computation we can now easily prove the following oscillation criteria.

Theorem 2. *Suppose that $\int^\infty r^{1-q}(t) dt = \infty$. Then each of the following conditions is sufficient for oscillation of (1):*

(i) *(Leighton-Wintner criterion [36]).*

$$\lim_{b \rightarrow \infty} \int^b c(t) = \infty. \quad (10)$$

(ii) *(Nehari-type criterion [14]). The integral $\int^\infty c(t) dt$ is convergent and*

$$\liminf_{t \rightarrow \infty} \left(\int^t r^{1-q}(s) ds \right)^{p-1} \left(\int_t^\infty c(s) ds \right) > 1. \quad (11)$$

Proof. A short computation and (9) show that each of conditions (i), (ii) implies that the points t_i , $i = 0, \dots, 3$, can be chosen in such a way that $\mathcal{F}(y; T, \infty) < 0$.

Concerning the nonoscillation criteria proved via the variational method, their proofs are usually based on the Wirtinger-type inequality

$$\int_T^\infty |M'(t)||y|^p dt \leq p^p \int_T^\infty \frac{M^p(t)}{|M'(t)|^{p-1}} |y'|^p dt \quad (12)$$

where M is a differentiable function such that $M'(t) \neq 0$ on $[T, \infty)$, which holds for every $0 \neq y \in W_0^{1,p}(T, \infty)$, see e.g. [12]. Using (12) we can prove the following statement.

Theorem 3. *Suppose that $\int^\infty r^{1-q}(t) dt = \infty$ and*

$$\limsup_{t \rightarrow \infty} \left(\int^t r^{1-q}(s) ds \right)^{p-1} \left(\int_t^\infty c_+(s) ds \right) < \frac{1}{p} \left(\frac{p-1}{p} \right)^{p-1}, \quad (13)$$

where $c_+(t) = \max\{0, c(t)\}$. Then (1) is nonoscillatory.

Proof. Setting $M(t) := \left(\int^t r^{1-q}(s) ds \right)^{1-p}$ and using (13) one can show that $\mathcal{F}(y; T, \infty) < 0$.

The previous two statements are typical examples of the application of the variational method in the oscillation theory of half-linear equations.

3.2 Riccati technique

In this subsection we briefly sketch how the equivalence (i) \iff (iii) can be used to derive (non)oscillation criteria. We illustrate the application of this method in the criteria which are similar to those presented in the previous subsection.

We start with the improvement of the statement given in Theorem 2 (ii).

Theorem 4. *Suppose that $\int^\infty r^{1-q}(t) dt = \infty$, the integral $\int^\infty c(t) dt$ is convergent and \liminf in (11) is $> \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1}$. Then (1) is oscillatory.*

Proof. By contradiction, suppose that (1) is nonoscillatory. Then solutions of the associated Riccati equation (4) satisfy the integral equation

$$w(t) = \int_t^\infty c(s) ds + (p-1) \int_t^\infty r^{1-q}(s) |w|^q ds,$$

see [34]. Multiplying this equation by $\left(\int^t r^{1-q}(s) ds\right)^{p-1}$, using (13) and supposing that $\limsup_{t \rightarrow \infty} \left(\int^t r^{1-q}(s) ds\right)^{p-1} w(t) =: \mu < \infty$ (in case when $\mu = \infty$, to get a contradiction is even easier than in our case $\mu < \infty$) we find an $\varepsilon > 0$ such that μ satisfies the inequality

$$\mu > \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1} + \varepsilon + |\mu|^q.$$

However, since $|t|^q - t + \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1} \geq 0$ for $t \in \mathbb{R}$, we have the required contradiction.

Nonoscillation criteria based on the Riccati technique are usually proved using a slightly different method than just the equivalence mentioned in Proposition 1. We look for a solution of the inequality

$$v' + c(t) + (p-1)r^{1-q}(t)|v|^q \leq 0 \tag{14}$$

instead of (4), since it essentially means that a certain Sturmian majorant of (1) is nonoscillatory and hence (1) is nonoscillatory as well.

Theorem 5. *Suppose that $\int^\infty r^{1-q}(t) dt = \infty$ and $\int^\infty c(t) dt$ converges. If*

$$\limsup_{t \rightarrow \infty} \left(\int^t r^{1-q}(s) ds\right)^{p-1} \left(\int_t^\infty c(s) ds\right) < \frac{1}{p} \left(\frac{p-1}{p}\right)^{p-1}, \tag{15}$$

$$\liminf_{t \rightarrow \infty} \left(\int^t r^{1-q}(s) ds\right)^{p-1} \left(\int_t^\infty c(s) ds\right) > -\frac{2p-1}{p} \left(\frac{p-1}{p}\right)^{p-1}, \tag{16}$$

then (1) is nonoscillatory.

Proof. Let $v(t) = \beta \left(\int^t r^{1-q}(s) ds \right)^{1-p}$, $\beta := \frac{1}{p} \left(\frac{p-1}{p} \right)^{p-1}$. Using (15) and (16) it is not difficult to verify that this function really satisfies (14), see [14].

All criteria presented by Theorems 2-5 contain the assumption $\int^\infty r^{1-q} = \infty$. A slight modification of the proofs of these statements enables us to find their counterparts in case $\int^\infty r^{1-q} < \infty$, see [14].

3.3 Perturbation principle and principal solution

In the previous criteria, equation (1) has been regarded as a perturbation of the (nonoscillatory) one-term differential equation (8). It was shown that if the “perturbation” function c in (1) is “sufficiently positive” (“not too positive”) then (1) becomes oscillatory (remains nonoscillatory). The exact quantitative characterization of these vague concepts is just given in Theorems 2-5.

From this point of view, it is a natural idea to consider (1) not as a perturbation of the one-term equation (8), but as a perturbation of the the general two-term nonoscillatory equation

$$(r(t)\Phi(x'))' + \tilde{c}(t)\Phi(x) = 0, \tag{17}$$

where \tilde{c} is a continuous function, and to formulate (non)oscillation criteria for (1) in terms of the difference $c - \tilde{c}$. Note that this idea applied to *linear* equation (2) does not produce essentially new criteria - due to the transformation identity (5). Indeed, this identity applied to (2) written in the form $(rx')' + cx = (rx')' + \tilde{c}x + (c - \tilde{c})x = 0$ and with the transformation function h which a solution of $(rx')' + \tilde{c}x = 0$ gives

$$h(t)[(r(t)x')' + \tilde{c}(t)x + (c(t) - \tilde{c}(t))x] = (r(t)h^2(t)y')' + h^2(t)[c(t) - \tilde{c}(t)]y,$$

so the resulting equation is again an equation of the form (2). One can then investigate it as the perturbation of the one-term equation $(r(t)h^2(t)y')' = 0$ and to transfer the obtained results “back” to the original equation (2).

Concerning the half-linear equation (1), as mentioned in Section 2, we have no half-linear version of (5), so the idea to investigate (1) as a perturbation of two-terms equation (17) brings really qualitatively new criteria. This approach was (implicitly) used for the first time by Elbert [23], where it is proved that (1) with $r \equiv 1$ is oscillatory provided

$$\lim_{b \rightarrow \infty} \int^b \left(c(t) - \frac{\gamma}{t^p} \right) t^{p-1} dt = \infty, \quad \gamma = \left(\frac{p-1}{p} \right)^p. \tag{18}$$

In this setting equation (1) with $r \equiv 1$ is viewed as a perturbation of the generalized Euler equation with the critical constant

$$(\Phi(x'))' + \frac{\gamma}{t^p}\Phi(x) = 0, \tag{19}$$

whose one solution $x(t) = t^{\frac{p-1}{p}}$ can be computed explicitly and other solutions behave asymptotically as $t^{\frac{p-1}{p}} \lg^{\frac{2}{p}} t$, see [24].

A natural question is why just the power t^{p-1} appears by the difference $c(t) - \tilde{\gamma}t^{-p}$ in (18). In the linear case $p = 2$ the answer is that $h = t^{1/2}$ is the so called *principal solution* of the Euler equation $x'' + \frac{1}{4t^2}x = 0$ (principal solutions of (2) and (1) are discussed in more details later in this section). More precisely, the transformation $x = t^{1/2}y$ transforms (2) with $r \equiv 1$ into the equation $(ty)'' + (c - \frac{1}{4t^2})ty = 0$.

The concept of the principal solution of (1) was introduced by Mirzov [37] and in [13,25] it was shown that this solution has many of the properties of the principal solution of (2). Using this concept (1) we can now prove the following generalization of the Elbert criterion (18), see [14,17].

Theorem 6. *Suppose that (17) is nonoscillatory and h is its (positive) principal solution. If*

$$\lim_{b \rightarrow \infty} \int^b (c(t) - \tilde{c}(t)) h^p(t) dt = \infty \quad (20)$$

then (1) is oscillatory.

Proof. The proof is similar to that of Theorem 2. We modify the test function (7) as follows. We let $y = h$ for $t \in [t_1, t_2]$. Then f, g are solution of (17) (instead of (8)) satisfying $f(t_1) = h(t_1)$, $g(t_2) = h(t_2)$. Using (20) and properties of principal solutions of (1) one can show that the points t_i , $i = 0, \dots, 3$, can be chosen in such a way that $\mathcal{F}(y; T, \infty) < 0$, see [17].

Observe that in case $\tilde{c} \equiv 0$ Theorem 6 reduces to Theorem 2(i). Indeed, if $\int^\infty r^{1-q} = \infty$ then $h \equiv 1$ is the principal solution of the one term equation (8) and (20) is the same as (10). Note also that Theorems 3-5 can be extended along the line treated in this subsection as well, we refer to [12,17] for more details.

We finish this part with some remarks and open problems concerning the principal solution of (1). In the linear case the principal solution \tilde{x} of (2) is (equivalently) defined as a solution for which

$$\lim_{t \rightarrow \infty} \frac{\tilde{x}(t)}{x(t)} = 0 \quad \iff \quad \int^\infty \frac{dt}{r(t)\tilde{x}^2(t)} = \infty, \quad (21)$$

where x is any solution of (2) linearly independent of \tilde{x} . Since both these characterizations are based on the linearity of the solutions space of (2), they do not extend (directly) to (1). Mirzov [37] in his construction of the principal solution of (1) used the fact that if this equation is nonoscillatory, then among all solutions of the associated Riccati equation (4) there exists the minimal one \tilde{w} , in the sense that any other solution w of this equation satisfies $w(t) > \tilde{w}(t)$ eventually. The principal solution of (1) is then defined by

$$\tilde{x}(t) = C \exp \left\{ \int^t r^{1-q}(s) \Phi^{-1}(\tilde{w}(s)) ds \right\},$$

Φ^{-1} being the inverse function of Φ , i.e. the solution \tilde{x} which is determined by $\tilde{w} = \frac{r\Phi(\tilde{x}')}{\Phi(\tilde{x})}$. In the linear case this construction is equivalent to (21), see [28].

In [16] we have tried to find an integral characterization of the principal solution of (1) in such a way that in the linear case it reduces to the second expression in (21). The main results of [16] are summarized in the next statement.

Theorem 7. *Suppose that equation (1) is nonoscillatory and \tilde{x} is its solution such that $\tilde{x}'(t) \neq 0$ for large t .*

(i) *Let $p \in (1, 2)$. If*

$$\int^{\infty} \frac{dt}{r(t)x^2(t)|x'(t)|^{p-2}} = \infty, \tag{22}$$

then \tilde{x} is the principal solution.

(ii) *Let $p > 2$. If \tilde{x} is the principal solution then (22) holds.*

(iii) *Suppose that $\int^{\infty} r^{1-q}(t) dt = \infty$, the function $\gamma(t) := \int_t^{\infty} c(s) ds$ exists and $\gamma(t) \geq 0$, but $\gamma(t) \not\equiv 0$ eventually. Then $\tilde{x}(t)$ is the principal solution if and only if (22) holds.*

This theorem shows that the equivalent integral characterization of the principal solution of (1) is known only in some particular cases. The subject of the present investigation is whether (22) is really a good characterization. Another intensively studied problem is the limit characterization of the principal solution of (1), i.e. the extension of the first relation of (21) to solutions of (1). For some results of this effort see [8,15,19].

4 Related topics

In this section we very briefly present some selected results related to the oscillation theory of (1).

4.1 Asymptotics of nonoscillatory solutions

In this subsection we suppose that the function c in (1) is positive or negative for large t , in the former case we suppose in addition that (1) is nonoscillatory. Since the solution space of (1) is homogeneous, we can restrict our attention to positive solutions. These solutions can be divided into two main classes according to their behaviour for large t

$$\mathbb{M}^+ = \{x : x'(t) > 0\}, \quad \mathbb{M}^- = \{x : x'(t) < 0\}$$

and each class $\mathbb{M}^+, \mathbb{M}^-$ is the union of two subclasses

$$\begin{aligned} \mathbb{M}^+ &= \mathbb{M}^+_{\infty} \cup \mathbb{M}^+_{B}, & \mathbb{M}^+_{\infty} &= \{x : x(t) \rightarrow \infty\}, & \mathbb{M}^+_{B} &= \{x : x(t) \rightarrow L < \infty\}, \\ \mathbb{M}^- &= \mathbb{M}^-_{B} \cup \mathbb{M}^-_{0}, & \mathbb{M}^-_{B} &= \{x : x(t) \rightarrow L > 0\}, & \mathbb{M}^-_{0} &= \{x : x(t) \rightarrow 0\}. \end{aligned}$$

The investigation of the classification of nonoscillatory solutions of linear equations (2) along this line was initiated in [35]. Afterward, several papers extending the results of this paper have appeared, see e.g. in [7,9] and the references given therein. An important role is played by the integrals

$$J_1 = \lim_{T \rightarrow \infty} \int^T r^{1-q}(t) \Phi^{-1} \left(\int^t c(s) ds \right) dt,$$

$$J_2 = \lim_{T \rightarrow \infty} \int^T r^{1-q}(t) \Phi^{-1} \left(\int_t^T c(s) ds \right) dt$$

and using these integrals the following results have been established in [7].

Theorem 8. *Suppose that $c(t) < 0$ for large t , then*

- (a) $J_1 = -\infty, J_2 = -\infty \implies \mathbb{M}^- = \mathbb{M}_0^- \neq \emptyset,$
- (b) $J_1 = -\infty, J_2 > -\infty \implies \mathbb{M}^- = \mathbb{M}_B^- \neq \emptyset,$
- (c) $J_1 > -\infty, J_2 > -\infty \implies \mathbb{M}_B^- \neq \emptyset, \mathbb{M}_0^- \neq \emptyset,$
- (d) $J_1 = -\infty \implies \mathbb{M}^+ = \mathbb{M}_\infty^+,$
- (e) $J_1 > -\infty \implies \mathbb{M}^+ = \mathbb{M}_B^+.$

4.2 BVP's associated with (1)

Consider the Dirichlet boundary value problem

$$(\Phi(x'))' + \lambda \Phi(x) + g(t, x) = f(t), \quad x(0) = 0 = x(\pi_p), \quad (23)$$

where λ is a real-valued parameter, the number π_p is defined in Section 2 and the “nonhalf-linearity” g and the forcing term f satisfy certain additional assumptions. The literature dealing with solvability of (23) (not only with the Dirichlet boundary condition) is very voluminous, see e.g. [10] and the references given therein. The situation is similar to the linear case in some aspects, but in some cases one meets completely different phenomena. Here we mention one of them, the Fredholm-type alternative for solvability of the BVP

$$(\Phi(x'))' + \lambda \Phi(x) = f(t), \quad x(0) = 0 = x(\pi_p). \quad (24)$$

Note that $\lambda_1 = (p-1)$ is the principal eigenvalue of the unforced equation (24).

Using the detailed analysis of the geometry of the functional associated with (24)

$$\mathcal{J}_f(y) = \frac{1}{p} \int_0^{\pi_p} \{|y'|^p - \lambda |y|^p\} dt - \int_0^{\pi_p} f(t)y dt$$

whose critical points over $W_0^{1,p}(0, \pi_p)$ are solutions of (24), one can prove the following statements, see [20] and the references given therein.

Theorem 9. Fredholm’s alternative for p -Laplacian with $p \neq 2$.

- (i) If $\lambda < 0$, the functional J_f has the unique minimum over $W_0^{1,p}(0, \pi_p)$, it is coercive and (24) has the unique solution.
- (ii) If $0 < \lambda < \lambda_1$, the functional J_f is still coercive, but there exists $f \in C[0, \pi_p]$ such that J_f has at least two critical points (one of them is the global minimum over $W_0^{1,p}(0, \pi_p)$, the other one is of saddle type).
- (iii) If $\lambda = \lambda_1$ (and similarly for higher eigenvalues $\lambda_n = (p - 1)^n$), the condition

$$\int_0^{\pi_p} f(t)S(t) dt = 0,$$

where $S(t)$ is the generalized sine function given in Section 2, i.e. the solution of (24) with $f \equiv 0$, is only sufficient but not necessary for solvability of (24). More precisely, there exists an open cone $\mathcal{C} \subset C[0, \pi_p]$ such that (24) has at least two solutions for every $f \in \mathcal{C}$ and

$$\int_0^{\pi_p} f(t)S(t) dt \neq 0.$$

4.3 Half-linear difference equations

In recent years, considerable attention has been paid to the oscillation theory of various *difference equations*. In particular, the oscillation theory of the Sturm-Louville difference equation

$$\Delta(r_k \Delta x_k) + c_k x_{k+1} = 0, \quad \Delta x_k = x_{k+1} - x_k,$$

has been deeply developed, see [1,31]. The discrete counterpart of (1) is the difference equation

$$\Delta(r_k \Phi(\Delta x_k)) + c_k \Phi(x_{k+1}) = 0, \quad r_k \neq 0. \tag{25}$$

The basic oscillatory properties of (25) are established in [38] and the main result of that paper is the discrete version of the Roundabout Theorem (Proposition 1).

Theorem 10. *The following statements are equivalent*

- (i) Equation (25) is disconjugate on the discrete interval $[0, N]$, i.e. the solution x given by the initial condition $x_0 = 0, x_1 \neq 0$ has no generalized zero in $(0, N + 1]$, i.e.

$$r_k x_k x_{k+1} > 0, \quad k = 1, \dots, N.$$

- (ii) There exists a solution x of (25) having no generalized zero in $[0, N + 1]$.
- (iii) There exists a solution $w = \{w_k\}_{k=0}^{N+1}$ of the Riccati-type equation

$$\Delta w_k + c_k + \left(1 - \frac{r_k}{\Phi(\Phi^{-1}(r_k)) + \Phi^{-1}(w_k)}\right) w_k = 0, \quad w = \frac{r\Phi(\Delta x)}{\Phi(x)}$$

such that $r_k + w_k > 0, k = 1, \dots, N$.

(iv) We have

$$\mathcal{F}_d(y; 0, N) = \sum_{k=0}^N \{r_k |\Delta y_k|^p - c_k |y_{k+1}|^p\} > 0$$

for every nontrivial $y = \{y_k\}_{k=0}^{N+1}$ satisfying $y_0 = 0 = y_{N+1}$.

4.4 Scalar methods for p -Laplacian

Several physical phenomena can be described by the partial differential equation with the so-called p -Laplacian

$$\operatorname{div} (|\nabla u|^{p-2} \nabla u) + c(x) \Phi(u) = 0, \quad x = (x_1, \dots, x_n) \in \mathbb{R}^n, \quad (26)$$

see e.g. [11]. It can be shown that the p -degree functional

$$\mathcal{F}(u; \Omega) = \int_{\Omega} \{|\nabla u|^p - c(x)|u|^p\} dx,$$

the Riccati type equation

$$\operatorname{div} w + c(x) + (p-1)|w|^q = 0, \quad w = \frac{\nabla u}{u} \quad (27)$$

and Picone's-type identity

$$\begin{aligned} \mathcal{F}(y; \Omega) &= \int_{\partial\Omega} w|y|^p dS + p \int_{\Omega} P(\nabla y, \Phi(y)w) dx, \\ P(u, v) &= \frac{\|u\|^p}{p} - \langle u, v \rangle + \frac{\|v\|^q}{q} \geq 0, \end{aligned}$$

where w is a solution of (27) defined in the whole domain $\overline{\Omega}$, can be used to establish Roundabout Theorem, Sturmian theory and to a certain extent also the oscillation theory similarly as for ordinary differential equation (1) and for partial differential equations with the "normal" Laplacian, see [2,3,18,29].

References

1. R. P. AGARWAL, *Difference Equations and Inequalities, Theory, Methods and Applications*, Second Edition, M. Dekker, New York-Basel, 2000.
2. W. ALLEGRETTO, Y. X. HUANG, *A Picone's identity for the p -Laplacian and applications*, Nonlin. Anal. **32** (1998), 819-830.
3. W. ALLEGRETTO, Y. X. HUANG, *Principal eigenvalues and Sturm comparison via Picone's identity*, J. Differ. Equations **156** (1999), 427-438.
4. I. BIHARI, *On the second order half-linear differential equation*, Studia Sci. Math. Hungar. **3** (1968), 411-437.
5. I. BIHARI, *An oscillation theorem concerning the half-linear differential equation of the second order*. Publ. Math. Inst. Hungar. Acad. Sci. Ser. A **8** (1963), 275-279.

6. O. BORŮVKA, *Lineare Differentialtransformationen 2. Ordnung*, Deutscher Verlag der Wissenschaften, Berlin 1961.
7. M. CECCHI, Z. DOŠLÁ, M. MARINI, *On nonoscillatory solutions of differential equations with p -Laplacian*, *Advances Math. Sci. Appl.* **11** (2001), 419-436.
8. M. CECCHI, Z. DOŠLÁ, M. MARINI, *Minimal sets for quasilinear differential equations*, submitted.
9. M. CECCHI, M. MARINI, G. VILLARI, *Integral criteria for a classification of solutions of linear differential equations*, *J. Differ. Equations* **99** (1992)
10. M. CUESTA, D. DE FIGUERO, J.-P. GOSSEZ, *The beginning of the Fučík spectrum for the p -Laplacian*, *J. Differ. Equations* **159** (1999), 212-238.
11. J. I. DÍAZ, *Nonlinear Partial Differential Equations and Free Boundaries. Vol I. Elliptic Equations*, Pitman, London 1985.
12. O. DOŠLÝ, *Oscillation criteria for half-linear second order differential equations*, *Hiroshima J. Math.* **28** (1998), 507-521.
13. O. DOŠLÝ, *A remark on conjugacy of half-linear second order differential equations*, *Math. Slovaca* **50** (2000), 67-79.
14. O. DOŠLÝ, *Methods of oscillation theory of half-linear second order differential equations*, *Czech Math. J.* **50** (125) (2000), 657-671.
15. O. DOŠLÝ, *Half-linear oscillation theory*, *Studies Univ. Žilina, Math.-Phys. Ser.* **13** (2001), 65-73.
16. O. DOŠLÝ, Á. ELBERT, *Integral characterization of principal solution of half-linear differential equations*, *Studia Sci. Math. Hungar.* **36** (2000), 455-469.
17. O. DOŠLÝ, A. LOMTATIDZE, *Oscillation and nonoscillation criteria for half-linear second order differential equations*, submitted.
18. O. DOŠLÝ, R. MAŘIK, *Nonexistence of positive solutions of PDE's with p -Laplacian*, *Acta Math. Hungar.* **90** (2001), 89-107,
19. O. DOŠLÝ, J. REZNÍČKOVÁ, *Regular half-linear second order differential equations*, submitted.
20. P. DRÁBEK, *Fredholm alternative for the p -Laplacian: yes or no?* Mustonen, Vesa (ed.) at al, *Function spaces, differential operators and nonlinear analysis: Proceedings of the conference FSDONA-99, Syöte, Finland, June 10-16, 1999*, Prague, Mathematical Institute of the Academy of Sciences of the Czech Republic, 57-64, (2000).
21. Á. ELBERT, *A half-linear second order differential equation*, *Colloq. Math. Soc. János Bolyai* **30** (1979), 158-180.
22. Á. ELBERT, *The Wronskian and the half-linear differential equations*, *Studia Sci. Math. Hungar.* **15** (1980), 101-105.
23. Á. ELBERT, *Oscillation and nonoscillation theorems for some non-linear ordinary differential equations*, *Lectures Notes in Math. No.* **964** (1982), 187-212.
24. Á. ELBERT, *Asymptotic behaviour of autonomous half-linear differential systems on the plane*, *Studia Sci. Math. Hungar.* **19** (1984), 447-464.
25. Á. ELBERT, T. KUSANO, *Principal solutions of nonoscillatory half-linear differential equations*, *Advances Math. Sci. Appl.* **18** (1998), 745-759.
26. R. EMDEN, *Gaskugeln, Anwendungen der mechanischen Warmentheorie auf Kosmologie und meteorologische Probleme*, Leipzig, 1907.
27. R. H. FOWLER, *The solutions of Emdem's and similar differential equations*, *Monthly Notices Roy. Astronom. Soc.* **91** (1930), 63-91.
28. P. HARTMAN, *Ordinary Differential Equations*, John Wiley & Sons, Inc., New York-London-Sydney, 1964.
29. J. JAROŠ, T. KUSANO, *A Picone type identity for half-linear differential equations*, *Acta Math. Univ. Comeniana* **68** (1999), 137-151.

30. J. JAROŠ, T. KUSANO, N. YOSIDA, *A Picone-type identity and Sturmian comparison and oscillation theorems for a class of half-linear partial differential equations of the second order*, Nonlin. Anal. TMA **40** (2000), 381-395.
31. W. G. KELLEY, A. PETERSON, *Difference Equations: An Introduction with Applications*, Acad. Press, San Diego, 1991.
32. T. KUSANO, Y. NAITO, *Oscillation and nonoscillation criteria for second order quasi-linear differential equations*, Acta. Math. Hungar. **76**, (1997), 81-99.
33. T. KUSANO, Y. NAITO, A. OGATA, *Strong oscillation and nonoscillation of quasi-linear differential equations of second order*, Diff. Equations Dyn. Syst. **2** (1994), 1-10.
34. H. J. LI, CH. CH. YEH, *Oscillations of half-linear second order differential equations*, Hiroshima Math. J. **25** (1995), 585-594.
35. M. MARINI, P. ZEZZA, *On the asymptotic behavior of the solutions of second order linear differential equations*, J. Differ. Equations **28** (1978), 1-17.
36. J. D. MIRZOV, *On some analogs of Sturm's and Kneser's theorems for nonlinear systems*, J. Math. Anal. Appl. **53** (1976), 418-425.
37. J. D. MIRZOV, *Principal and nonprincipal solutions of a nonoscillatory system*, Tbiliss. Gos. Univ. Inst. Prikl. Mat. Trudy **31** (1988), 100-117.
38. P. ŘEHÁK, *Oscillatory properties of second order half-linear difference equations*, Czech. Math. J., **51** (126) (2001), 303-321.
39. G. SANSONE, *Equazioni differenziali nel campo reale I,II*, Zanichelli, Bologna, 1949.
40. L. H. THOMAS, *The calculation of atomic fields*, Proc. Cambridge Phil. Soc. **23** (1927), 542-548.