

EQUADIFF 10

Yakov Yakubov

Irregular boundary value problems for ordinary differential equations

In: Jaromír Kuben and Jaromír Vosmanský (eds.): Equadiff 10, Czechoslovak International Conference on Differential Equations and Their Applications, Prague, August 27-31, 2001, [Part 2] Papers. Masaryk University, Brno, 2002. CD-ROM; a limited number of printed issues has been issued. pp. 437--441.

Persistent URL: <http://dml.cz/dmlcz/700374>

Terms of use:

© Institute of Mathematics AS CR, 2002

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Irregular boundary value problems for ordinary differential equations

Yakov Yakubov*

School of Mathematical Sciences, Tel-Aviv University,
69978 Tel-Aviv, Israel

Email: yakubov@post.tau.ac.il

Abstract. Birkhoff-irregular boundary value problems for quadratic ordinary differential pencils of the second order have been considered. The spectral parameter may appear in a boundary condition, the equation contains an abstract linear operator while the boundary conditions contain internal points of an interval and a linear functional. Isomorphism and coerciveness with a defect are proved for such problems. Two-fold completeness of root functions of corresponding spectral problems is also established. As an application of the obtained results, an initial boundary value problem for second order parabolic equations is considered, and the well-posedness and completeness of the elementary solutions are proved. These and some other results have been published in [1].

MSC 2000. 34L10,34B05,35K15

Keywords. Birkhoff-irregular boundary value problems, isomorphism, completeness, well-posedness

1 Isomorphism and Two-fold Completeness

Consider a principally (because of the addition of B , T , and x_{j_i}) boundary value problem for ordinary differential equations

$$L(\lambda)u := \lambda^2 u(x) + \lambda [a_1 u'(x) + b_1 u(x)] + [a_2 u''(x) + b_2 u'(x)] + Bu|_x = f(x), \quad x \in (0, 1), \quad (1)$$

* Research supported in part by the Israel Ministry of Absorption

$$\left\{ \begin{aligned} L_1(\lambda)u &:= \lambda \left[\alpha_{11}u(0) + \beta_{11}u(1) + \sum_{i=1}^{N_0} \eta_{0i}u(x_{0i}) \right] + \\ &\quad + [\alpha_{10}u'(0) + \beta_{10}u'(1) + \gamma_{10}u(0) + \delta_{10}u(1)] + \\ &\quad + \sum_{i=1}^{N_1} \eta_{1i}u'(x_{1i}) + \sum_{i=1}^{N_2} \eta_{2i}u(x_{2i}) + Tu = f_1, \\ L_2u &:= [\alpha_{20}u(0) + \beta_{20}u(1)] + \sum_{i=1}^{N_3} \eta_{3i}u(x_{3i}) = f_2, \end{aligned} \right. \tag{2}$$

where $a_k, b_k, \alpha_{\nu k}, \beta_{\nu k}, \gamma_{10}, \delta_{10}, \eta_{ji}, f_\nu$ are complex numbers, $f(x)$ is a given function; $x_{ji} \in (0, 1)$; B is a linear operator in $L_q(0, 1)$ and T is a linear functional in $L_q(0, 1)$, a real $q \in (1, \infty)$.

We assume that $a_2 \neq 0$ and denote by

$$\omega_1 := \frac{-a_1 + (a_1^2 - 4a_2)^{\frac{1}{2}}}{2a_2}, \quad \omega_2 := \frac{-a_1 - (a_1^2 - 4a_2)^{\frac{1}{2}}}{2a_2}$$

the roots of the equation

$$a_2\omega^2 + a_1\omega + 1 = 0,$$

where $z^{\frac{1}{2}} := |z|^{\frac{1}{2}} e^{i\frac{\arg z}{2}}$, $-\pi < \arg z \leq \pi$. We also assume that $\arg \omega_1 \neq \arg \omega_2$. Further,

$$\underline{\omega} := \min\{\arg \omega_1, \arg \omega_2 + \pi\},$$

$$\overline{\omega} := \max\{\arg \omega_1, \arg \omega_2 + \pi\},$$

and values $\arg \omega_j$ are chosen up to a multiple of 2π , so that $\overline{\omega} - \underline{\omega} < \pi$.

Introduce now the following notations:

$$\theta_0(\omega_1, \omega_2) := \begin{vmatrix} \alpha_{11} + \alpha_{10}\omega_1 & \beta_{11} + \beta_{10}\omega_2 \\ \alpha_{20} & \beta_{20} \end{vmatrix},$$

$$\theta_1(\omega_1, \omega_2) := \begin{vmatrix} -\alpha_{10} \frac{b_1 + b_2\omega_1}{(a_1^2 - 4a_2)^{\frac{1}{2}}} + \gamma_{10} & \beta_{10} \frac{b_1 + b_2\omega_2}{(a_1^2 - 4a_2)^{\frac{1}{2}}} + \delta_{10} \\ \alpha_{20} & \beta_{20} \end{vmatrix}.$$

Definition 1. Problem (1)—(2) is called *regular (regular with defect 1) with respect to the numbers ω_1, ω_2* if:

- (1) $a_2 \neq 0, \arg \omega_1 \neq \arg \omega_2$;
- (2) $x_{ji} \in (0, 1)$, for some real $q \in (1, \infty)$ the operator B from $W_q^1(0, 1)$ into $L_q(0, 1)$ is compact and the functional T is continuous in $L_q(0, 1)$;
- (3) $\theta_0(\omega_1, \omega_2) \neq 0$ ($\theta_0(\omega_1, \omega_2) = 0, \theta_1(\omega_1, \omega_2) \neq 0$).

Theorem 2. *Let problem (1)–(2) be regular with defect 1 with respect to the numbers ω_1, ω_2 .*

Then for any $\varepsilon > 0$ there exists $R_\varepsilon > 0$ such that for all complex numbers λ that satisfy $|\lambda| > R_\varepsilon$ and lying inside the angle

$$\frac{\pi}{2} - \underline{\omega} + \varepsilon < \arg \lambda < \frac{3\pi}{2} - \bar{\omega} - \varepsilon,$$

the operator

$$\mathbb{L}(\lambda) : u \rightarrow \mathbb{L}(\lambda)u := (L(\lambda)u, L_1(\lambda)u, L_2u)$$

from $W_q^2(0, 1)$ onto $L_q(0, 1) \dot{+} \mathbb{C}^2$ is an isomorphism and for these λ the following estimate holds for a solution $u(x)$ of problem (1)–(2)

$$\sum_{k=0}^2 |\lambda|^{1-k} \|u\|_{W_q^k(0,1)} \leq C(\varepsilon) \left(\|f\|_{L_q(0,1)} + \sum_{\nu=1}^2 |\lambda|^{\nu-\frac{1}{q}} |f_\nu| \right).$$

Theorem 3. *Let $|\alpha_{\nu 0}| + |\beta_{\nu 0}| \neq 0, \nu = 1, 2$ and let homogeneous problem (1)–(2) be regular with defect 1 with respect to the numbers ω_1, ω_2 and regular, or regular with defect 1 with respect to the numbers ω_2, ω_1 , for $q = 2$.*

Then the spectrum of homogeneous problem (1)–(2) is discrete and a system of its root functions (eigenfunctions and associated functions) is two-fold complete in the space

$$H := \{v \mid v := (v_1, v_2) \in W_2^1(0, 1) \oplus L_2(0, 1), L_2v_1 = 0\}.$$

2 Well-posedness and Completeness of Elementary Solutions

Consider, in $[0, T] \times [0, 1]$, the following initial boundary value problem

$$u_t(t, x) + [au_{xx}(t, x) + bu_x(t, x)] + Bu(t, \cdot)|_x = f(t, x), \tag{3}$$

$$\left\{ \begin{aligned} L_1u &:= [\alpha_{10}u'_x(t, 0) + \beta_{10}u'_x(t, 1)] + [\gamma_{10}u(t, 0) + \delta_{10}u(t, 1)] + \\ &+ \sum_{i=1}^{N_1} \eta_{1i}u'_x(t, x_{1i}) + \sum_{i=1}^{N_2} \eta_{2i}u(t, x_{2i}) + Qu(t, \cdot) = 0, \end{aligned} \right. \tag{4}$$

$$\left\{ \begin{aligned} L_2u &:= [\alpha_{20}u(t, 0) + \beta_{20}u(t, 1)] + \sum_{i=1}^{N_3} \eta_{3i}u(t, x_{3i}) = 0, \\ u(0, x) &= u_0(x). \end{aligned} \right. \tag{5}$$

Let E, E_1 , and E_2 be Banach spaces. Introduce two Banach spaces

$$C_\mu((0, T], E) := \{f \mid f \in C((0, T], E), \|f\| = \sup_{t \in (0, T]} \|t^\mu f(t)\| < \infty\}, \mu \geq 0,$$

$$C_\mu^\gamma((0, T], E) := \{f \mid f \in C((0, T], E), \|f\| = \sup_{t \in (0, T]} \|t^\mu f(t)\| + \sup_{0 < t < t+h \leq T} \|f(t+h) - f(t)\| h^{-\gamma} t^\mu < \infty\}, \mu \geq 0, \gamma \in (0, 1],$$

and a linear space (in the case $E_1 \subset E_2$)

$$C^1((0, T], E_1, E_2) := \{f \mid f \in C((0, T], E_1) \cap C^1((0, T], E_2)\},$$

where $C((0, T], E)$ and $C^1((0, T], E)$ are spaces of continuous and continuously differentiable, respectively, vector-functions from $(0, T]$ into E .

Theorem 4. *Let the following conditions be satisfied:*

- (1) $a \neq 0, |\arg a| > \frac{\pi}{2}, \alpha_{10}\beta_{20} + \alpha_{20}\beta_{10} = 0, \gamma_{10}\beta_{20} - \delta_{10}\alpha_{20} - \frac{b}{a}\alpha_{10}\beta_{20} \neq 0;$
- (2) *the operator B from $W_q^1(0, 1)$ into $L_q(0, 1)$ is compact;*
- (3) *the functional Q is continuous in $L_q(0, 1)$.*
- (4) $f \in C_\mu^\gamma((0, T], L_q(0, 1))$ for some $\gamma \in (\frac{1}{2}, 1]$ and $\mu \in [0, \frac{1}{2});$
- (5) $u_0 \in W_q^2((0, 1), L_\nu u = 0, \nu = 1, 2)$.

Then problem (3)—(5) has a unique solution

$$u \in C([0, T], L_q(0, 1)) \cap C^1((0, T], W_q^2(0, 1), L_q(0, 1))$$

and for the solution the following estimates hold

$$\|u(t, \cdot)\|_{L_q(0,1)} \leq C(\|u_0\|_{W_q^2(0,1)} + \|f\|_{C_\mu((0,t],L_q(0,1))}),$$

and

$$\|u(t, \cdot)\|_{W_q^2(0,1)} + \|u'(t, \cdot)\|_{L_q(0,1)} \leq Ct^{-1}(\|u_0\|_{W_q^2(0,1)} + \|f\|_{C_\mu^\gamma((0,t],L_q(0,1))}),$$

for $t \in (0, T]$.

Consider now a spectral problem corresponding to the homogeneous (3), (4):

$$\lambda u(x) + [au''(x) + bu'(x)] + Bu|_x = 0, \quad x \in (0, 1), \tag{6}$$

$$\left\{ \begin{aligned} L_1 u &:= [\alpha_{10}u'(0) + \beta_{10}u'(1)] + [\gamma_{10}u(0) + \delta_{10}u(1)] + \\ &+ \sum_{i=1}^{N_1} \eta_{1i}u'(x_{1i}) + \sum_{i=1}^{N_2} \eta_{2i}u(x_{2i}) + Qu = 0, \\ L_2 u &:= [\alpha_{20}u(0) + \beta_{20}u(1)] + \sum_{i=1}^{N_3} \eta_{3i}u(x_{3i}) = 0, \end{aligned} \right. \tag{7}$$

A function of the form

$$u_j(t, x) = e^{\lambda_j t} \left(\frac{t^{k_j}}{k_j!} u_{j0}(x) + \frac{t^{k_j-1}}{(k_j-1)!} u_{j1}(x) + \dots + u_{jk_j}(x) \right) \tag{8}$$

becomes an elementary solution of the homogeneous (3), (4) if and only if a system of the functions $u_{j0}(x), u_{j1}(x), \dots, u_{jk_j}(x)$ is a chain of root functions of problem (6)—(7), corresponding to the eigenvalue λ_j .

Theorem 5. *Let conditions of the previous theorem be satisfied with $q = 2$. Then problem (3)–(5) (with $f(t, x) \equiv 0$) has a unique solution*

$$u \in C([0, T], L_2(0, 1)) \cap C^1((0, T], W_2^2(0, 1), L_2(0, 1))$$

and there exist numbers c_{jn} such that

$$\lim_{n \rightarrow \infty} \sup_{t \in (0, T]} \|u(t, \cdot) - \sum_{j=1}^n c_{jn} u_j(t, \cdot)\|_{L_2(0,1)} = 0,$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \sup_{t \in (0, T]} t(\|u'_t(t, \cdot) - \sum_{j=1}^n c_{jn} u'_{jt}(t, \cdot)\|_{L_2(0,1)} + \\ + \|u(t, \cdot) - \sum_{j=1}^n c_{jn} u_j(t, \cdot)\|_{W_2^2(0,1)}) = 0, \end{aligned}$$

where $u(t, x)$ is a solution to problem (3)–(5) (with $f(t, x) \equiv 0$) and $u_j(t, x)$ is an elementary solution (8) of homogeneous problem (3)–(4).

References

1. Yakubov Ya., *Irregular Boundary Value Problems for Ordinary Differential Equations*, Analysis, **18** 1998, 359–402.

