Erik G. F. Thomas Application of vector integration to spectral theory

In: Zdeněk Frolík (ed.): Abstracta. 4th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1976. pp. 141–144.

Persistent URL: http://dml.cz/dmlcz/701063

Terms of use:

 $\ensuremath{\mathbb C}$ Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1976

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* http://dml.cz

FOURTH WINTER SCHOOL (1976).

APPLICATIONS OF VECTOR INTEGRATION TO SPECTRAL THEORY

py

G.Erik F. THOMAS

1. A generalization of Bochner's integral.

The spectral theory requires a theory of integration of functions with values in locally convex spaces which in general are neither metrisable nor separable. The Pettis integral in its full generality is not suitable because the most useful theorems of classical integration theory e.g. Fubini's thm, Lebesgue's differentiation thm, are not valid for general Pettis integrals.

Let T be a topological Hausdorff space, μ a Radon measure on T, and assume T is a countable union of open sets of finite measure.

Definition: A function f: $T \rightarrow E$, quasi complete l.c.v.s. is totally μ -summable if it is μ -measurable (Lusin property) and if there exists BCE, bounded closed, absolutely convex, such that $\int |f|_{B} d < +\infty$. $(j_{X})_{B} \leq \inf\{A, x \in AB\}$)

It can be shown that a totally μ -summable function is Pettis integrable and that the integrals $\int_A f d_{\mu} \mu$ belong to E_B the Banach space generated by B. Although $f(t) \in E_B$ a.e. $f: T \longrightarrow E_B$ is not in general Bochner integrable. Nevertheless most of the useful properties of Bochner integral carry

141

over, in particular Fubini's thm and the Lebesgue differentiation theorem e.g. $\frac{d}{dt} \int_a^t f(s)ds = f(t)$ a.e.

To extend the definition to general measure spaces requires the solution of a problem: see Problem book .

2. Given a Hilbert space \mathcal{H} and a selfadjoint or normal operator (T,D_T) , with discrete spectrum, there exists a decomposition

(1) $\mathcal{H} = \bigoplus \mathcal{H}_{\lambda}$ \mathcal{H} \mathcal{H} \mathcal{H} be the formula of T $\mathcal{A} \in \mathcal{O}(T)$ with

 $x = \sum_{\lambda} x_{\lambda} \text{ in } \mathcal{H}$ $\|x\|^2 = \sum \|x_{\lambda}\|^2$

when the spectrum is not discrete, one can obtain a decomnosition $T = \int \lambda E(d\lambda)$ with E() the spectral measure associated with T, but a decomposition closer to (1) is also possible as follows:

Let \mathfrak{L} be a quasi-complete locally convex space over \mathfrak{C} . With every Hilbert subspace $\mathfrak{R} \hookrightarrow \mathfrak{L}$ there is associated a kernel $H = \mathfrak{j}\mathfrak{j}^*$ a map from the anti-dual of \mathfrak{L} , \mathfrak{L}^* , to \mathfrak{L} , such that $H(\mathfrak{L}^*) \subset \mathfrak{R}$ and $(x, H\mathfrak{e}^*) =$ $= \langle x, \mathfrak{e}^* \rangle$. The kernel H is linear Hermitian: $\langle H\mathfrak{e}^*, \mathfrak{f}^* \rangle =$ $= \langle H\mathfrak{f}^*, \mathfrak{e}^* \rangle$ and positive $\langle H\mathfrak{e}^*, \mathfrak{e}^* \rangle \geq 0$. It has been shown by L. Schwartz [2] that conversely there corresponds to every positive kernel H: $\mathfrak{L}^* \longrightarrow \mathfrak{L}$ a unique Hilbert subspace $\mathfrak{R} \subset \mathfrak{L}$. The set of positive kernels is a closed convex cone in the real locally convex space of all Hermitian kernels, topologized with the topology of pointwise convergence in \mathscr{C}^* ; the topology is carried over to the set Hilb(\mathscr{C}) of Hilbert subspaces of \mathscr{C} . If we now put $\mathscr{R}_{\underline{A}} =$ = E(A) \mathscr{H} the map A $\longrightarrow \mathscr{R}_{\underline{A}}$ is a Hilb (\mathscr{C}) valued measure which under conditions to be described below has a Radon Ni-

kodym derivative with respect to a basis measure μ .

Theorem. Let \mathscr{C} be a Fréchet space and assume the map $\mathscr{H} \xrightarrow{\mathcal{F}} \mathscr{C}$ is absolutely summing. Then given a spectral measure $E(\)$ there exists a measure $\omega \geq 0$ such that $(\omega(A) = 0 \iff \mathscr{H}_A = 0 \text{ and } a \quad \omega$ -essentially unique family $\mathcal{A} \longrightarrow \mathscr{H}_A = \operatorname{Hilb}(\mathscr{C})$ totally ω -summable such that $\mathscr{H}_A = \int_A \mathscr{H}_A d\mu(\mathcal{A})$

This implies that every $x \in \mathcal{H}$ has a decomposition.

$$x = \int x(\lambda) d\mu(\lambda)$$
 in \mathcal{E}

with

$$\|\mathbf{x}\|^{2} = \int \|\mathbf{x}(\boldsymbol{\lambda})\|_{\mathcal{U}_{\mathcal{X}}}^{2} d\mu(\boldsymbol{\lambda}).$$

$$\mathbf{E}(\mathbf{A})\mathbf{x} = \int_{A} \mathbf{x}(\boldsymbol{\lambda}) d\mu(\boldsymbol{\lambda})$$

and

$$Tx = \int \lambda x (\lambda) d\mu(\lambda) \qquad x \in D_{T}$$

if E () is associated with T.

Furthermore if there exists a continuous linear operator $T: \mathcal{C} \longrightarrow \mathcal{C}$ such that $T = \frac{T}{D_T}$ it follows that $\mathcal{H}_{\mathcal{A}} \subset \operatorname{Ker}(\mathcal{A}\mathcal{J} - \mathcal{J})$ for $(u \ a.e. \mathcal{A})$, i.e. $\mathcal{I} \times = \mathcal{A} \times \quad \forall \times \in \mathcal{H}_{\mathcal{A}}$.

Remark. The condition **j** absolutely summing cannot be improved in this theorem.

This is closely related to and inspired by the works of

- [1] Gel'fand & Shilov: Generalized Functions
- [2] L. Schwartz: Sous-espaces Hilbertiens d'espaces vectoriels topologiques et noyaux reproduisants. Journal d'Analyse Mathématique d'Israel 1964.