

Rastislav Telgársky

Scattered compactifications

In: Zdeněk Frolík (ed.): Abstracta. 5th Winter School on Abstract Analysis. Czechoslovak Academy of Sciences, Praha, 1977. pp. 99–101.

Persistent URL: <http://dml.cz/dmlcz/701101>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic, 1977

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

SCATTERED COMPACTIFICATIONS

R. TELGÁRSKY

Wrocław

A space is said to be scattered if it does not contain a dense in itself subset. E.g., the space $W(\alpha) = \{\xi : \xi < \alpha\}$ is scattered in the open interval topology for any ordinal α .

Z. Semadeni [10] posed the following question: Is there a scattered compactification for any completely regular scattered space? In particular: Is there a scattered compactification for $N \cup \{p\} \subset \beta N$ where $p \in \beta N - N$?

In this contribution we present a survey of results concerning the above questions.

(1) [4] Each metrizable separable scattered space can be embedded in $W(\alpha)$ for some $\alpha < \omega_1$ and thus it admits a scattered compactification.

Example 3.6, p. 17, in [10] is incorrect. We have

(2) [14] Each metrizable scattered space can be embedded in $W(\alpha)$ for some ordinal α , and thus it admits a scattered compactification. Similarly as in [14] one can prove the following.

(3) Each suborderable paracompact scattered space can be embedded in $W(\alpha)$ for some ordinal α , and thus it admits a scattered compactification.

(4) [7] Each suborderable scattered space of countable height is orderable and admits an orderable scattered compactification.

(5) [14] If X is a paracompact scattered space, then $\text{ind } X = \text{Ind } X = \dim X = 0$.

Hence we have

(6) If X admits a scattered compactification, then $\text{ind } X = 0$.

It is easy to prove that

(7) If βX contains a non-degenerate continuum C such that $C \cap X \neq \emptyset$, then X has no scattered compactification.

(8) [8] (CH) Example of a completely regular scattered space X of cardinality $\aleph_1$ with $\text{ind } X > 0$; by (6), X has no scattered compactification.

(9) [12] Example of a completely regular scattered space X of the cardinality $2^{\aleph_0}$ with $\text{ind } X > 0$; by (6), X has no scattered compactification.

(10) [9] (CH) If p is a P -point in $\beta N - N$, then $N \cup \{p\}$ admits a scattered compactification.

A point $p \in \beta\mathbb{N} - \mathbb{N}$ is said to be a P-point of order 2 in $\beta\mathbb{N} - \mathbb{N}$ if there exists a countable set M of P-points in $\beta\mathbb{N} - \mathbb{N}$ such that p is a P-point in $\overline{M} - M$.

(11) [2] (CH) If p is a P-point of order 2 in $\beta\mathbb{N} - \mathbb{N}$, then $\mathbb{N} \cup \{p\}$ admits a scattered compactification.

(12) [3] (CH) If p is a P-point of order α in $\beta\mathbb{N} - \mathbb{N}$, where $\alpha < \omega_1$, then $\mathbb{N} \cup \{p\}$ admits a scattered compactification.

(13) [11] If p belongs to the carrier of a regular Eorel non-atomic measure μ on $\beta\mathbb{N}$ with $\mu(\beta\mathbb{N}) = 1$, then $\mathbb{N} \cup \{p\}$ has no scattered compactification.

Let us note that the carrier of a regular Eorel non-atomic measure μ on $\beta\mathbb{N}$ with $\mu(\beta\mathbb{N}) = 1$ is a perfect subset of $\beta\mathbb{N} - \mathbb{N}$ satisfying the countable chain condition.

(14) [15] If p belongs to a perfect subset of $\beta\mathbb{N} - \mathbb{N}$ satisfying the countable chain condition, then $\mathbb{N} \cup \{p\}$ has no scattered compactification.

(15) [15] The set $\{p \in \beta\mathbb{N} - \mathbb{N} : \mathbb{N} \cup \{p\} \text{ has no scattered compactification}\}$ is $\aleph_0$ -bounded; hence, in particular, it is countably compact.

A point $p \in S$ is said to be a point of extremal disconnectedness of S if given two disjoint open sets U and V in S we have $p \notin \overline{U} \cap \overline{V}$. Let S^{ed} denote the set of all points of extremal disconnectedness of S . Note that each subset of $\beta\mathbb{N} - \mathbb{N}$ satisfying the countable chain condition is extremally disconnected.

(16) [16] If βX contains a dense in itself subset S such that $X \cap S^{\text{ed}} \neq \emptyset$, then X has no scattered compactification.

If X is not pseudocompact, then βX contains a copy of $\beta\mathbb{N}$ and thus βX cannot be scattered. Moreover

(17) [5] If X is not pseudocompact, then there exists a point $p \in \beta X - X$ such that $X \cup \{p\}$ has no scattered compactification.

(18) [6] Let X be a product of uncountably many two-point discrete spaces. Let one point be given a base of neighborhoods as in the product topology; let all other points of X be isolated. Then X has no scattered compactification.

A filter F is called K -regular if it contains a subset F_0 of cardinality K such that the intersection of infinitely many members of F_0 is empty.

(19) [13] If X is a space with the only non-isolated point p , $\text{card } X > \aleph_1$ and the neighborhood base at p restricted to $X - \{p\}$ is K -regular, where $\aleph_1 < K \leq \text{card } X$, then X has no scattered compactification.

(20) [1] Example of a countable space with one non-isolated point all compactifications of which contain $\beta\mathbb{N}$.

References

- [1] E.K.van Douwen and T.Przymusiński, First countable and countable spaces all compactifications of which contain βN , Fund. Math. (to appear).
- [2] M.Jayachandran and M.Rajagopalan, Scattered compactifications for $N \cup \{p\}$, Pacific J.Math. 61 (1975), 161-171.
- [3] M.Jayachandran and M.Rajagopalan, Scattered compactifications of some scattered spaces, preprint 1975.
- [4] B.Knaster et K.Urbaniak, Sur les espaces complets séparables de dimension 0, Fund. Math. 40 (1953), 194-202.
- [5] V.I.Malyhin, On scattered spaces having no scattered compactifications (in Russian), preprint 1976.
- [6] P.J.Nyikos, Not every scattered space has a scattered compactification, Notices A.M.S. 21 (1974), A-570.
- [7] S.Purisch, Scattered compactifications and the orderability of scattered spaces, submitted to General Topology and its Applications.
- [8] M.Rajagopalan, Scattered spaces III, preprint 1976.
- [9] C.Ryll-Nardzewski and R.Telgársky, On the scattered compactification, Bull. Acad. Polon. Sci. 18 (1970), 233-234.
- [10] Z.Semadeni, Sur les ensembles clairsemés, Rozprawy Matematyczne 19, Warszawa 1959.
- [11] R.C.Solomon, A space of the form $N \cup \{p\}$ with no scattered compactification, Bull.Acad.Polon.Sci. 24 (1976), 755-756.
- [12] R.C.Solomon, A scattered space that is not 0-dimensional, Bull. London Math. Soc. 8 (1976), 239-240.
- [13] R.C.Solomon, Scattered spaces and their compactifications, preprint 1975.
- [14] R.Telgársky, Total paracompactness and paracompact dispersed spaces, Bull. Acad. Polon. Sci. 16 (1968), 567-572.
- [15] R.Telgársky, Subspaces $N \cup \{p\}$ of βN with no scattered compactifications, Bull. Acad. Polon. Sci. 25 (1977) (in press).
- [16] R.Telgársky, Scattered compactifications and points of extremal disconnectedness, Bull. Acad. Polon. Sci. 25 (1977), 159-163.

Institute of Mathematics, Technical University, Wrocław, Poland