

Lech Drewnowski

A solution to a problem of De Wilde - Tsirulnikov

In: Zdeněk Frolík (ed.): Abstracta. 9th Winter School on Abstract Analysis.
Czechoslovak Academy of Sciences, Praha, 1981. pp. 19–20.

Persistent URL: <http://dml.cz/dmlcz/701218>

Terms of use:

© Institute of Mathematics of the Academy of Sciences of the Czech Republic,
1981

Institute of Mathematics of the Czech Academy of Sciences provides access to
digitized documents strictly for personal use. Each copy of any part of this
document must contain these *Terms of use*.



This document has been digitized, optimized for electronic
delivery and stamped with digital signature within the project
DML-CZ: The Czech Digital Mathematics Library <http://dml.cz>

NINTH WINTER SCHOOL ON ABSTRACT ANALYSIS (1981)

A solution to a problem of De Wilde - Tsirulnikov

L. Drewnowski

In a recent paper [1], M. De Wilde and B. Tsirulnikov have shown that if E is a dense barrelled subspace of a B -complete locally convex space X , then there is an order-reversing correspondence between the closed subspaces M of X with $M \cap E = \{0\}$ and the weaker barrelled topologies ξ on E . Under this correspondence, the topologies ξ which are assigned to infinite-dimensional subspaces M are nontrivial in the sense that they give rise to duals of E which are of infinite codimension in the original dual of E .

In connection with this, the following question has been asked in [1]:

Suppose E is a dense barrelled subspace of uncountable codimension in a Banach space X . Does there always exist an infinite-dimensional subspace M of X such that $M \cap E = \{0\}$?

The Theorem below answers this question in the negative. Its proof depends heavily upon the results presented at this Winter School in the talk of Z. Lipecki [2].

T H E O R E M. Every infinite-dimensional Banach (or Fréchet) space X has a dense Baire (hence barrelled) subspace E of codimension at least $c = 2^{\aleph_0}$ such that if M is a closed subspace of X with $M \cap E = \{0\}$, then M is of finite dimension.

P r o o f. Let Y be a closed subspace of X such that the dimension of $Z = X/Y$ is c . (E.g., take $Y = \bigcap_n \ker f_n$, where (f_n) is a linearly independent sequence in X^* .) Call a subspace $W \subset Z$ a κ -subspace of Z if every linearly independent sequence (z_n) in Z with $\sum_n z_n$ subseries convergent has a subsequence (w_n) such that $0 \neq \sum_n w_n \in W$. Clearly, every κ -sub-

space W in Z has property (K) ([2]), and it is not hard to see that $\dim W \geq c$. Now, using Theorem 1 and Proposition 1 of [2], we can find two dense κ -subspaces U and V in Z such that $Z = U \oplus V$ (algebraically). As observed above, each of them has property (K) and is of dimension at least c .

Let $Q: X \rightarrow Z$ be the quotient map and define $E = Q^{-1}(U)$. Since U is dense in Z , E is dense in X ; moreover, $\operatorname{codim} E = \dim V \geq c$. Since U has property (K), so does E ; hence, by Theorem 3 in [2], E is Baire.

Finally, suppose that there is an infinite-dimensional closed subspace M in X with $M \cap E = \{0\}$. Choose a linearly independent sequence (x_n) in M for which $\sum_n x_n$ is subseries convergent. Then also the sequence (Qx_n) is linearly independent and the series $\sum_n Qx_n$ is subseries convergent. Since U is a κ -subspace in Z , there is a subsequence (y_n) of (x_n) such that $0 \neq \sum_n Qy_n \in U$. It follows that $0 \neq y = \sum_n y_n \in E$. On the other hand, however, we have $y \in M$ because all y_n are in M and M is closed. Hence $M \cap E \neq \{0\}$; a contradiction.

Thus E has all the required properties.

R e m a r k. The above theorem remains valid for all infinite dimensional complete metric linear spaces X which have a Hausdorff quotient X/Y of dimension c . We do not know if this is always the case.

Let us note also that the main results of [1] can be extended to the non-locally convex setting.

R e f e r e n c e s

- [1] L. De Wilde and B. Tsirlinikov, Barrelled spaces with a B-complete completion, to appear.
- [2] Z. Lipecki, On some nonclosed subspaces of metric linear spaces, this volume, pp. 410 - 413.