

Ivan Kolář; Peter W. Michor

### All natural concomitants of vector valued differential forms

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1987. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 16. pp. [101]–108.

Persistent URL: <http://dml.cz/dmlcz/701412>

### Terms of use:

© Circolo Matematico di Palermo, 1987

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

# ALL NATURAL CONCOMITANTS OF VECTOR VALUED DIFFERENTIAL FORMS

Ivan Kolář, Peter W. Michor

We deduce that in general all natural bilinear concomitants of the type of the Frölicher-Nijenhuis bracket form a 10-dimensional real vector space. - All manifolds and maps are assumed to be infinitely differentiable.

1. Let  $\Omega^p(M, TM)$  denote the space of differential forms of order  $p$  on a manifold  $M$  with values in the tangent bundle  $TM$ , which are called the vector valued  $p$ -forms on  $M$ , [4]. Obviously,  $\Omega^0(M, TM)$  coincides with the set  $\mathcal{X}(M)$  of all vector fields on  $M$ . We consider, for each  $n$ -dimensional manifold  $M$ , a bilinear map  $B_M: \Omega^p(M, TM) \times \Omega^q(M, TM) \rightarrow \Omega^{p+q}(M, TM)$  satisfying  $f^*B_M(P, Q) = B_N(f^*P, f^*Q)$  for each local diffeomorphism  $f: N \rightarrow M$ . Such a family  $B = (B_M)$  is called a natural concomitant of vector valued differential forms.

2. We will use the following notation, [9]. Let  $\Omega^q(M)$  be the space of differential  $q$ -forms on  $M$  and  $\Omega(M) = \bigoplus_q \Omega^q(M)$  be the exterior algebra of  $M$ . Every  $P \in \Omega^p(M, TM)$  determines a graded derivation  $i(P)$  of degree  $p-1$  in  $\Omega(M)$  characterized by  $i(P)\omega = \omega \circ P$  if  $\omega \in \Omega^1(M)$  and  $i(P)f = 0$  for all functions  $f \in \Omega^0(M)$ . The explicit formula for  $i(P)$  is

$$(1) \quad i(P)\omega(X_1, \dots, X_{p+q-1}) = \frac{1}{p!(q-1)!} \sum_{\sigma} (\text{sign } \sigma) \omega(X_{\sigma 1}, \dots, X_{\sigma p}, X_{\sigma(p+1)}, \dots, X_{\sigma(p+q-1)})$$

with  $\omega \in \Omega^q(M)$ ,  $X_1, \dots, X_{p+q-1} \in \mathcal{X}(M)$  and summation with respect to all permutations  $\sigma$  of  $p+q-1$  letters, [9]. This formula makes sense also for a vector valued  $q$ -form  $Q \in \Omega^q(M, TM)$  and defines a vector valued  $(p+q-1)$ -form  $i(P)Q \in \Omega^{p+q-1}(M, TM)$ .

-----  
This paper is in final form and no version of it will be submitted for publication elsewhere.

Let  $\Theta(P) = i(P) \circ d - (-1)^{p-1} d \circ i(P)$  be the graded commutator of  $i(P)$  with the exterior differential  $d$ . For a vector field  $X \in \Omega^0(M, TM)$ ,  $\Theta(X)$  coincides with the Lie derivative with respect to  $X$ . According to [4], for every  $P \in \Omega^p(M, TM)$  and  $Q \in \Omega^q(M, TM)$  there exists a unique element  $[P, Q] \in \Omega^{p+q}(M, TM)$  satisfying  $\Theta([P, Q]) = \Theta(P) \circ \Theta(Q) - (-1)^{pq} \Theta(Q) \circ \Theta(P)$ . This is called the Frölicher-Nijenhuis bracket of  $P$  and  $Q$ . It can be characterized by

$$[\varphi \otimes X, \psi \otimes Y] = \varphi \wedge \psi \otimes [X, Y] + \varphi \wedge \Theta(X) \psi \otimes Y - \Theta(Y) \varphi \wedge \psi \otimes X \\ + (-1)^p (d\varphi \wedge i(X) \psi \otimes Y + i(Y) \varphi \wedge d\psi \otimes X)$$

for all  $\varphi \in \Omega^p(M)$ ,  $\psi \in \Omega^q(M)$ ,  $X, Y \in \mathfrak{X}(M)$ , [9]. The coordinate expression of  $[P, Q]$  is

$$[P_\alpha^i d^\alpha \otimes \partial_i, Q_\beta^j d^\beta \otimes \partial_j] = (P_\alpha^i \dots \partial_j Q_\beta^j \delta_{p+1} \dots \delta_{p+q} \\ - (-1)^{pq} Q_\beta^j \delta_1 \dots \delta_q \partial_j P_\alpha^i \delta_{q+1} \dots \delta_{q+p} - P_\alpha^i \delta_1 \dots \delta_{p-1}^j \partial_j Q_\beta^j \delta_p \delta_{p+1} \dots \delta_{p+q} \\ (2) + (-1)^{pq} Q_\beta^j \delta_1 \dots \delta_{q-1}^j \partial_j P_\alpha^i \delta_q \delta_{q+1} \dots \delta_{q+p}) d^\alpha \otimes \partial_i$$

where  $d^\alpha = dx^{\alpha_1} \wedge \dots \wedge dx^{\alpha_p}$  and  $\partial_i = \partial / \partial x^i$  in local coordinates  $x = (x^1, \dots, x^n)$  on  $M$ . This follows from (1.9) of [9] by inserting the basic vector fields  $\partial_i$  into the global formula.

The wedge product of a differential  $q$ -form and a vector valued  $p$ -form is a bilinear map  $\Omega^q(M) \times \Omega^p(M, TM) \rightarrow \Omega^{p+q}(M, TM)$  characterized by  $\omega \wedge (\varphi \otimes X) = (\omega \wedge \varphi) \otimes X$  for all  $\omega \in \Omega^q(M)$ ,  $\varphi \in \Omega^p(M)$ ,  $X \in \mathfrak{X}(M)$ . Further, let  $C: \Omega^p(M, TM) \rightarrow \Omega^{p-1}(M)$  be the contraction operator defined by  $C(\omega \otimes X) = i(X)\omega$  for all  $\omega \in \Omega^p(M)$ ,  $X \in \mathfrak{X}(M)$ . In particular, for  $P \in \Omega^0(M, TM)$  we have  $C(P) = 0$ . Clearly  $C(i(P)Q)$  is a multiple of  $C(i(Q)P)$ ,  $P \in \Omega^p(M, TM)$ ,  $Q \in \Omega^q(M, TM)$ . Finally, write  $I = \text{Id}_{TM} \in \Omega^1(M, TM)$ .

**3. Theorem.** For  $\dim M > p+q+1$ , all natural concomitants of vector valued differential forms  $\Omega^p(M, TM) \times \Omega^q(M, TM) \rightarrow \Omega^{p+q}(M, TM)$  form a vector space linearly generated by the following 10 operators

$$[P, Q], dC(P) \wedge Q, dC(Q) \wedge P, \\ (3) \quad dC(P) \wedge C(Q) \wedge I, dC(Q) \wedge C(P) \wedge I, dC(i(P)Q) \wedge I, \\ i(P)dC(Q) \wedge I, i(Q)dC(P) \wedge I, di(P)C(Q) \wedge I, di(Q)C(P) \wedge I$$

The proof of this theorem will occupy the rest of the paper.

We remark that for  $p < 2$  or  $q < 2$  some expressions in (3) vanish identically. In particular, for  $p = q = 0$  we have only one non-trivial generator  $[P, Q]$ . One sees easily that the Frölicher-Nijenhuis bracket of two vector valued 0-forms, i.e. of two vector fields, coincides with the classical bracket. In this case we rededuce a result by S. van Strien, [14], and D. Krupka with V. Mikolášová, [7], that all natural concomitants of two vector fields are the scalar multiples of the classical bracket. An infinitesimal version of this result was deduced by M. de Wilde and P. Lecomte, [2].

4. From [4] or [9] we see that all expressions in the theorem are indeed natural. For the converse suppose that  $B = (B_M)$  is a natural concomitant as in 1. Since  $B_M$  is a local operator, it is a (finite order) differential operator by the bilinear Peetre theorem, [1], see also [13]. In [9] it was checked that  $B$  is a homogeneous bilinear differential operator of total order 1.

5. Let  $P^2M$  be the second order frame bundle of  $M$  (i.e. the space of all invertible 2-jets of  $R^n$  into  $M$  with source 0), which is a principal fibre bundle with structure group  $G_n^2$  of all invertible 2-jets of  $R^n$  into  $R^n$  with source and target 0. According to a general theory, see e.g. [6], the first order natural operators from  $\Omega^p(M, TM) \times \Omega^q(M, TM)$  into  $\Omega^{p+q}(M, TM)$  correspond to the  $G_n^2$ -equivariant maps from the standard fibre  $S$  of the first jet prolongation of the first bundle into the standard fibre  $R^n \otimes \wedge^{p+q} R^{n*}$  of the second bundle. Clearly,  $S$  is the product of four spaces  $R^n \otimes \wedge^p R^{n*}$ ,  $R^n \otimes \wedge^p R^{n*} \otimes R^{n*}$ ,  $R^n \otimes \wedge^q R^{n*}$ ,  $R^n \otimes \wedge^q R^{n*} \otimes R^{n*}$  and the action of  $G_n^2$  on  $S$  is given by the usual transformation law of tensor fields and their first order partial derivatives. In order to be completely clear we specify this in detail: the transformation law of a tensor field of type  $\binom{1}{p}$  is

$$(4) \quad \bar{P}_{j_1 \dots j_p}^i = P_{m_1 \dots m_p}^{\ell} \frac{\partial \bar{x}^i}{\partial x^{\ell}} \frac{\partial x^{m_1}}{\partial \bar{x}^{j_1}} \dots \frac{\partial x^{m_p}}{\partial \bar{x}^{j_p}}$$

and the transformation law of its first partial derivatives is

$$(5) \quad \bar{P}_{j_1 \dots j_p, k}^i = P_{m_1 \dots m_p, n}^{\ell} \frac{\partial \bar{x}^i}{\partial x^{\ell}} \frac{\partial x^{m_1}}{\partial \bar{x}^{j_1}} \dots \frac{\partial x^{m_p}}{\partial \bar{x}^{j_p}} \frac{\partial x^n}{\partial \bar{x}^k} +$$

$$\begin{aligned}
 (5) \quad & + p_{m_1 \dots m_p}^{\bar{x}^i} \left( \frac{\partial^2 \bar{x}^i}{\partial x^{\bar{x}^i} \partial x^n} \frac{\partial x^{m_1}}{\partial \bar{x}^{j_1}} \dots \frac{\partial x^{m_p}}{\partial \bar{x}^{j_p}} \frac{\partial x^n}{\partial \bar{x}^k} \right. \\
 & + \frac{\partial \bar{x}^i}{\partial x^{\bar{x}^i}} \frac{\partial^2 x^{m_1}}{\partial \bar{x}^{j_1} \partial \bar{x}^k} \frac{\partial x^{m_2}}{\partial \bar{x}^{j_2}} \dots \frac{\partial x^{m_p}}{\partial \bar{x}^{j_p}} \\
 & \left. + \dots + \frac{\partial \bar{x}^i}{\partial x^{\bar{x}^i}} \frac{\partial x^{m_1}}{\partial \bar{x}^{j_1}} \dots \frac{\partial^2 x^{m_p}}{\partial \bar{x}^{j_p} \partial \bar{x}^k} \right)
 \end{aligned}$$

Since  $B$  is bilinear and of total order 1, its associated map  $B_0: S \rightarrow R^n \otimes \wedge^{p+q} R^{n*}$  is a sum  $B_0 = B_1 + B_2$ , where  $B_1$  and  $B_2$  are bilinear maps

$$(6) \quad B_1: R^n \otimes \wedge^p R^{n*} \otimes R^{n*} \times R^n \otimes \wedge^q R^{n*} \rightarrow R^n \otimes \wedge^{p+q} R^{n*}$$

$$(7) \quad B_2: R^n \otimes \wedge^p R^{n*} \times R^n \otimes \wedge^q R^{n*} \otimes R^{n*} \rightarrow R^n \otimes \wedge^{p+q} R^{n*}$$

We remark that  $B_0$  corresponds to the canonical coordinate expression of the natural operator  $B$  on  $R^n$ .

6. Consider the canonical inclusion  $GL(n, R) \hookrightarrow G_n^2$  transforming every matrix into the 2-jet at 0 of the corresponding linear transformation of  $R^n$ . By (4) and (5), the linear maps associated with the bilinear maps  $B_1$  and  $B_2$ , which will be denoted by the same symbol, are  $GL(n, R)$ -equivariant, so that we can apply the classical theory of invariant tensors, [3]. Consider first the following diagram

$$\begin{array}{ccc}
 R^n \otimes \wedge^p R^{n*} \otimes R^{n*} \otimes R^n \otimes \wedge^q R^{n*} & \xrightarrow{B_1} & R^n \otimes \wedge^{p+q} R^{n*} \\
 (8) \quad \text{id} \otimes \text{Alt}_p \otimes \text{id} \otimes \text{Alt}_q \uparrow \downarrow & & \text{id} \otimes \text{Alt}_{p+q} \uparrow \downarrow \\
 R^n \otimes \bigotimes^p R^{n*} \otimes R^{n*} \otimes R^n \otimes \bigotimes^q R^{n*} & \longrightarrow & R^n \otimes \wedge^{p+q} R^{n*}
 \end{array}$$

where  $\text{Alt}$  denotes the alternator of the indicated degree. Since the vertical maps are also  $GL(n, R)$ -equivariant, it suffices to determine all  $GL(n, R)$ -equivariant maps in the bottom row and to restrict them and to take the alternator of the result. By the theory of invariant tensors, [3], all  $GL(n, R)$ -equivariant maps

$\bigotimes^2 R^n \otimes \bigotimes^{p+q+1} R^{n*} \rightarrow R^n \otimes \bigotimes^{p+q} R^{n*}$  are given by all kinds of permutations of the indices, all contractions and tensorizing with the identity. Since we apply this to alternating forms and use the

alternator on the result, permutations do not play a role.

7. In what follows we discuss the case  $p \geq 2$ ,  $q \geq 2$  only and we leave the other cases to the reader. (A direct discussion shows that in the remaining cases the list (9) below should be reduced by those terms that do not make sense, but our next procedure leads to (3) as well.) Constructing  $B_1$ , we may contract the vector field part of  $P$  into a non-derivation entry of  $P$  or into the derivation entry of  $P$  or into  $Q$ , and we may contract the vector field part of  $Q$  into  $Q$  or into a non-derivation entry of  $P$  or into the derivation entry of  $P$ , and then tensorize with  $I = \text{id}_{R^n}$ . This gives  $9-1=8$  po-

ssibilities. If we perform only one contraction, we get  $3+3=6$  further possibilities, so that we have a 14-parameter family, which corresponds to the lower case letters in the list (9) below. Constructing  $B_2$ , we obtain analogously another 14-parameter family denoted by upper case letters in the list (9) below. Hence by  $GL(n, R)$ -equivariancy we deduce the following expression for  $B_0: S \rightarrow R^n \otimes \wedge^{p+q} R^{n*}$  (we do not indicate alternation in the subscripts and we write  $\alpha, \beta$  for any kind of free form-index on the right hand side)

$$\begin{aligned}
 B_0^i = & a_{m\alpha, k}^p Q_{n\beta}^n \delta_{\frac{1}{2}}^i + b_{\alpha, m}^p Q_{n\beta}^n \delta_{\frac{1}{2}}^i + c_{\alpha, k}^p Q_{nm\beta}^n \delta_{\frac{1}{2}}^i + d_{mn\alpha, k}^p Q_{\beta}^n \delta_{\frac{1}{2}}^i \\
 & + e_{n\alpha, m}^p Q_{\beta}^n \delta_{\frac{1}{2}}^i + f_{n\alpha, k}^p Q_{m\beta}^n \delta_{\frac{1}{2}}^i + g_{m\alpha, n}^p Q_{\beta}^n \delta_{\frac{1}{2}}^i + h_{\alpha, n}^p Q_{m\beta}^n \delta_{\frac{1}{2}}^i \\
 (9) \quad & + i_{m\alpha, k}^p Q_{\beta}^i + j_{\alpha, m}^p Q_{\beta}^i + k_{\alpha, k}^p Q_{m\beta}^i + l_{\alpha, k}^p Q_{n\beta}^i + m_{n\alpha, k}^p Q_{\beta}^i + n_{\alpha, n}^p Q_{\beta}^i \\
 & + A_{m\alpha}^p Q_{n\beta, k}^n \delta_{\frac{1}{2}}^i + B_{m\alpha}^p Q_{\beta, n}^n \delta_{\frac{1}{2}}^i + C_{mn\alpha}^p Q_{\beta, k}^n \delta_{\frac{1}{2}}^i + D_{\alpha}^p Q_{nm\beta, k}^n \delta_{\frac{1}{2}}^i \\
 & + E_{\alpha}^p Q_{m\beta, n}^n \delta_{\frac{1}{2}}^i + F_{n\alpha}^p Q_{m\beta, k}^n \delta_{\frac{1}{2}}^i + G_{\alpha}^p Q_{n\beta, m}^n \delta_{\frac{1}{2}}^i + H_{n\alpha}^p Q_{\beta, m}^n \delta_{\frac{1}{2}}^i \\
 & + I_{\alpha}^p Q_{n\beta, k}^i + J_{\alpha}^p Q_{\beta, n}^i + K_{n\alpha}^p Q_{\beta, k}^i + L_{m\alpha}^p Q_{\beta, k}^i + M_{\alpha}^p Q_{m\beta, k}^i + N_{\alpha}^p Q_{\beta, m}^i
 \end{aligned}$$

8. The map  $B_0$  is also equivariant with respect to the kernel of the canonical projection  $G_n^2 \rightarrow GL(n, R)$  of 2-jets into 1-jets, which coincides with the abelian group  $R^n \otimes S^2 R^{n*}$ , provided  $S^2$  denotes the second symmetric tensor power. By (4) and (5), the action of an element  $(S_{jk}^i) \in R^n \otimes S^2 R^{n*}$  on  $R^n \otimes \wedge^p R^{n*}$  is just the identity and its action on  $R^n \otimes \wedge^p R^{n*} \otimes R^{n*}$  is

$$(10) \quad \bar{P}_{j_1 \dots j_p, k}^i = P_{j_1 \dots j_p, k}^i + P_{j_1 \dots j_p}^t S_{tk}^i - P_{tj_2 \dots j_p}^i S_{j_1 k}^t - \dots \\ - P_{j_1 \dots j_{p-1} t}^i S_{j_p k}^t$$

Hence the condition that (9) is equivariant under the action of  $R^n \otimes S^2 R^{n*}$  has the following form (the alternation in the subscripts is not indicated explicitly)

$$0 = a(P_{mj_2 \dots j_p}^t S_{tk}^m - P_{tj_2 \dots j_p}^m S_{mk}^t - P_{mtj_3 \dots j_p}^m S_{j_2 k}^t - \dots \\ - P_{mj_2 \dots j_{p-1} t}^m S_{j_p k}^t) Q_{n\beta}^n \delta_{\beta}^i + \dots + n(P_{j_1 \dots j_p}^t S_{tn}^i \\ (11) - P_{tj_2 \dots j_p}^i S_{j_1 n}^t - \dots - P_{j_1 \dots j_{p-1} t}^i S_{j_p n}^t) Q_{\beta}^n \\ + A P_{m\alpha}^m (Q_{nj_2 \dots j_p}^t S_{tk}^n - Q_{tj_2 \dots j_p}^n S_{nk}^t - Q_{ntj_3 \dots j_p}^n S_{j_2 k}^t - \dots \\ - Q_{nj_2 \dots j_{p-1} t}^n S_{j_p k}^t) \delta_{\beta}^i + \dots + N P_{m\alpha}^m (Q_{j_1 \dots j_p}^t S_{tm}^i \\ - Q_{tj_2 \dots j_p}^i S_{j_1 m}^t - \dots - Q_{j_1 \dots j_{p-1} t}^i S_{j_p m}^t)$$

We remark that every term containing an  $S$  with both free subscripts vanishes after applying the alternator.

9. The right hand side of (11) represents a trilinear map  $R^n \otimes \wedge^p R^{n*} \times R^n \otimes \wedge^q R^{n*} \times R^n \otimes S^2 R^{n*} \rightarrow R^n \otimes \wedge^{p+q} R^{n*}$  depending on the parameters  $a, \dots, N$ , which can be rewritten in the following form

$$\left[ B P_{m\alpha}^m Q_{\beta}^t S_{tn}^n + (-1)^q q B P_{m\alpha}^m Q_{t\beta}^n S_{nk}^t + b P_{\alpha}^t Q_{n\beta}^n S_{tm}^m - (-1)^{p+q} p b P_{t\alpha}^m Q_{n\beta}^n S_{mk}^t \right. \\ + ((-1)^q c - D - (-1)^q (q-1) G) P_{\alpha}^m Q_{nt\beta}^n S_{mk}^t + (C - (-1)^q d \\ - (-1)^{p+q} (p-1) g) P_{mt\alpha}^m Q_{\beta}^n S_{nk}^t + e P_{n\alpha}^m Q_{\beta}^n S_{mt}^t + (H - e) P_{t\alpha}^m Q_{\beta}^n S_{mn}^t \\ (12) + E P_{\alpha}^m Q_{m\beta}^t S_{tn}^n + (h - E) P_{\alpha}^m Q_{t\beta}^n S_{mn}^t + ((-1)^q q H - (-1)^q f - F) P_{n\alpha}^m Q_{t\beta}^n S_{mk}^t \\ + (F + (-1)^q f - (-1)^{p+q} p h) P_{n\alpha}^m Q_{m\beta}^t S_{tk}^n - (-1)^{p+q} (p-1) e P_{nt\alpha}^m Q_{\beta}^n S_{mk}^t \\ \left. - (-1)^q (q-1) E P_{\alpha}^m Q_{mt\beta}^n S_{nk}^t \right] \delta_{\beta}^i + j P_{\alpha}^t Q_{\beta}^i S_{tm}^m + (-1)^{p+q} p j P_{t\alpha}^m Q_{\beta}^i S_{mk}^t +$$

$$\begin{aligned}
 & + J P_{\alpha}^i Q_{\beta}^t S_{tm}^m + (-1)^q q J P_{\alpha}^i Q_{\beta}^m S_{mk}^t - ((-1)^q k - (-1)^q q_{N+M}) P_{\alpha}^m Q_{\beta}^i S_{mk}^t \\
 (12) & + (K + (-1)^{p+q} p n - (-1)^q m) P_{m\alpha}^i Q_{\beta}^t S_{tk}^m - \mathfrak{L}(-1)^q P_{\alpha}^t Q_{m\beta}^m S_{tk}^i \\
 & + L P_{m\alpha}^m Q_{\beta}^t S_{tk}^i - (-1)^q m P_{m\alpha}^t Q_{\beta}^m S_{tk}^i + M P_{\alpha}^m Q_{m\beta}^t S_{tk}^i + (n+N) P_{\alpha}^m Q_{\beta}^t S_{mt}^i
 \end{aligned}$$

The equivariancy of  $B_0$  with respect to  $R^n \otimes S^2 R^{n*}$  is equivalent to the fact that (12) is the zero map.

10. For  $\dim M > p+q+1$  the indices has at least  $p+q+2$  different values, which implies easily that all individual terms in (12) are linearly independent in  $\text{Hom}(R^n \otimes \wedge^{p+q} R^{n*} \otimes R^n \otimes \wedge^q R^{n*} \otimes R^n \otimes S^2 R^{n*}, R^n \otimes \wedge^{p+q} R^{n*})$ . Hence (12) is the zero map if and only if all the coefficients vanish. This leads to the following equations

$$\begin{aligned}
 (13) \quad & b = B = e = E = h = H = j = J = \mathfrak{L} = L = m = M = 0, \\
 & c = (q-1)G + (-1)^q D, \quad C = (-1)^q d + (-1)^{p+q}(p-1)g \\
 & F = (-1)^{q-1}f, \quad k = -qn, \quad K = (-1)^{p+q-1}pn, \quad N = -n
 \end{aligned}$$

while  $a, A, d, D, f, g, G, n, i, I$  are independent parameters.

11. Thus, for  $\dim M > p+q+1$  we have deduced 10-parameter family (13) and one verifies easily that (9) with (13) represents the coordinate expression of the linear combinations of the elements of (3). This completes the proof of our Theorem.

Obviously, for  $\dim M < p+q$  it holds  $\Omega^{p+q}(M, TM) = 0$ , so that we can construct the zero operator only. The study of the two remaining cases  $\dim M = p+q$  and  $\dim M = p+q+1$  consists in detailed discussion of the conditions for (12) to be the zero map, which is not included into the present paper.

#### REFERENCES

- [1] CAHEN M., DE WILDE M., GUTT S. "Local cohomology of the algebra of  $C^\infty$ -functions on a connected manifold", Lett. Math. Phys., 4(1980), 157-167.
- [2] DE WILDE M., LECOMTE P. "Algebraic characterizations of the algebra of functions and of the Lie algebra of vector fields on a manifold", Compositio Math., 40(1980), 79-85.

- [3] DIEUDONNÉ J. A., CARRELL J. B. "Invariant Theory, Old and New", Academic Press, New York - London 1971.
- [4] FRÖLICHER A., NIJENHUIS A. "Theory of vector valued differential forms, part I", Indag. Math., 18(1956), 338-359.
- [5] FRÖLICHER A., NIJENHUIS A. "Invariance of vector form operations under mappings", Comment. Math. Helv., 34(1960), 227-248.
- [6] KOLÁŘ I. "Some natural operators in differential geometry", to appear in Proc. Conf. Differential Geometry and its Applications, Brno 1986, D. Reidel.
- [7] KRUPKA D., MIKOLÁŠOVÁ V. "On the uniqueness of some differential invariants:  $d, [, ], \nabla$ ", Czechoslovak Math. J., (34) (109)(1984), 588-597.
- [8] MANGIAROTTI L., MODUGNO M. "Graded Lie algebras and connections on a fibred space", J. Math. Pures Appl., 63(1984), 111-120.
- [9] MICHOR P. W. "Remarks on the Frölicher-Nijenhuis bracket", to appear in Proc. Conf. Differential Geometry and its Applications, Brno 1986, D. Reidel.
- [10] NIJENHUIS A. "Jacobi type identities for bilinear differential concomitants for certain tensor fields I", Indag. Math., 17(1955), 390-403.
- [11] PALAIS R. S. "Natural operations on differential forms", Trans. Amer. Math. Soc., 92(1959), 125-141.
- [12] PALAIS R. S., TERNG C. L. "Natural bundles have finite order", Topology, 16(1977), 271-277.
- [13] SLOVÁK J. "On the finite order of some operators", to appear in Proc. Conf. Differential Geometry and its Applications, Brno 1986, D. Reidel.
- [14] VAN STRIEN S. "Unicity of the Lie product", Compositio Math., 45(1982), 199-205.

Authors' addresses:

MATHEMATICAL INSTITUTE OF THE ČSAV, BRANCH BRNO, MENDELOVO NÁM. 1,  
CS - 66282 BRNO.

INSTITUT FÜR MATHEMATIK, UNIVERSITÄT WIEN, STRUDLHOFGASSE 4,  
A - 1090 WIEN.