

Thomas Friedrich; Olga Pokorná

Twistor spinors and solutions of the equation (E) on Riemannian manifolds

In: Jarolím Bureš and Vladimír Souček (eds.): Proceedings of the Winter School "Geometry and Physics". Circolo Matematico di Palermo, Palermo, 1991. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 26. pp. [149]–154.

Persistent URL: <http://dml.cz/dmlcz/701488>

Terms of use:

© Circolo Matematico di Palermo, 1991

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

TWISTOR SPINORS AND SOLUTIONS OF THE EQUATION (E) ON RIEMANNIAN MANIFOLDS

by

Thomas Friedrich (Berlin) and Olga Pokorna (Prague)

1. Introduction

Let M^n be a Riemannian spin manifold and denote by R its scalar curvature. The conformally invariant twistor operator \mathfrak{D} acting on sections ψ of the spinor bundle S is defined by the covariant derivative and the projection onto the kernel of the Clifford multiplication. The kernel $\text{Ker}(\mathfrak{D})$ of this operator is the space of all spinor fields ψ satisfying the first order differential equation

$$\nabla_X \psi + \frac{1}{n} X \cdot D \psi = 0 \quad \langle 1 \rangle$$

where D denotes the Dirac operator (see [3],[4]).

Killing spinors, i.e. spinor fields satisfying for some complex number $\lambda \in \mathbb{C}$ the equation

$$\nabla_X \psi = \lambda X \cdot \psi \quad \langle 2 \rangle$$

are special solutions of the twistor equation. A. Lichnerowicz (see [5]) studied the so-called equation (E) for spinor fields:

$$\nabla_X (D\psi) + \frac{R}{4(n-1)} X \cdot \psi = 0. \quad \langle 3 \rangle$$

In particular, A. Lichnerowicz proved that if a connected Riemannian manifold admits a non-trivial solution of the equation (E) then its scalar curvature R is constant. Consequently, in case of a compact Riemannian manifold the space of all twistor spinors coincides with $\text{Ker}(E)$ and with the space

of all Killing spinors (see [2]). The aim of this note is to compare the kernel of the twistor operator with the space $\text{Ker}(E)$ of all solutions of equation (E).

2. A relation between $\text{Ker}(E)$ and $\text{Ker}(\mathcal{D})$.

We consider a Riemannian spin manifold (M^n, g) with a non-trivial solution of equation (E). Then its scalar curvature R is constant.

Proposition 1:

$$\text{Ker}(E) = \text{Ker}(D^2 - \frac{nR}{4(n-1)}) \cap D^{-1}(\text{Ker}(\mathcal{D}))$$

Proof: If $\psi \in \text{Ker}(E)$ we obtain from equation <3>

$$D^2\psi = \sum_{i=1}^n e_i \cdot \nabla_{e_i} (D\psi) = - \sum_{i=1}^n e_i \cdot \frac{R}{4(n-1)} e_i \cdot \psi = \frac{nR}{4(n-1)} \psi,$$

i.e. ψ belongs to the kernel of the operator $D^2 - \frac{nR}{4(n-1)}$.

Furthermore, we have

$$\begin{aligned} \nabla_X(D\psi) + \frac{1}{n} X \cdot D(D\psi) &= \bar{\nabla}_X(D\psi) + \frac{1}{n} X \cdot \frac{nR}{4(n-1)} \psi = \\ &= \bar{\nabla}_X(D\psi) + \frac{R}{4(n-1)} X \cdot \psi = 0 \end{aligned}$$

and, consequently, $D\psi$ is a twistor spinor. Conversely, if $\psi \in \text{Ker}(D^2 - \frac{nR}{4(n-1)}) \cap D^{-1}(\text{Ker}(\mathcal{D}))$ then we have

$D^2\psi = \frac{nR}{4(n-1)} \psi$ as well as $\bar{\nabla}_X(D\psi) + \frac{1}{n} X \cdot D(D\psi) = 0$ and ψ belongs to $\text{Ker}(E)$.

Theorem 1: Let (M^n, g) be a Riemannian spin manifold with constant scalar curvature $R \neq 0$. Then the map

$$\text{Ker}(E) \ni \psi \longrightarrow D\psi \in \text{Ker}(\mathcal{D})$$

is an isomorphism.

Proof: By proposition 1 the map is well-defined. If $D\psi = 0$

and $\psi \in \text{Ker}(E)$ the equation $\langle 3 \rangle$ implies

$$\frac{R}{4(n-1)} X \cdot \psi = 0.$$

Since $R \neq 0$ we conclude $\psi = 0$, i.e. the map $\text{Ker}(E) \rightarrow \text{Ker}(\mathcal{D})$ is injective. On the other hand, given a twistor spinor $\varphi \in \text{Ker}(\mathcal{D})$ we consider

$$\psi = \frac{4(n-1)}{n R} D\varphi.$$

$$\text{Then } D\psi = \frac{4(n-1)}{n R} D^2\varphi = \frac{4(n-1)}{n R} \frac{n R}{4(n-1)} \varphi = \varphi$$

$$\begin{aligned} \text{and } \nabla_X(D\psi) + \frac{R}{4(n-1)} X \cdot \psi &= \\ = \nabla_X\left(\frac{4(n-1)}{n R} D^2\varphi\right) + \frac{1}{n} X \cdot D\varphi &= \\ = \nabla_X(\varphi) + \frac{1}{n} X \cdot D\varphi &= 0. \end{aligned}$$

This means that ψ belongs to $\text{Ker}(E)$ and is the preimage of φ , i.e. the map $\text{Ker}(E) \rightarrow \text{Ker}(\mathcal{D})$ is surjective.

Corollary: Let (M^n, g) be a connected Riemannian manifold with scalar curvature $R \neq 0$. Then $\left[\frac{n}{2}\right] + 1$

$$\dim \text{Ker}(E) = \dim \text{Ker}(\mathcal{D}) \leq 2$$

Proof: see [3].

Proposition 2: If $\psi \in \text{Ker}(E)$ and $R \neq 0$ then

$$\nabla_X \psi = \frac{2(n-1)}{R(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot D\psi.$$

Proof: If ψ belongs to $\text{Ker}(E)$ then by proposition 1 $D\psi$ is a twistor spinor. The general Lichnerowicz formula

$$\nabla_X(D\psi) = \frac{n}{2(n-2)} \left(\frac{R}{2(n-1)} X \cdot \varphi - \text{Ric}(X) \cdot \varphi \right) \quad \langle 4 \rangle$$

valid for any twistor spinor φ (see [4]) now yields

$$\nabla_X(D^2\psi) = \frac{n}{2(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot D\psi.$$

Moreover, since $\psi \in \text{Ker}(E)$ it satisfies the equation

$$D^2\psi = \frac{n}{4(n-1)} R \psi$$

and we obtain

$$\frac{n}{4(n-1)} \nabla_X \psi = \frac{n}{2(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot D\psi.$$

Theorem 2:

a) Let (M^n, g) be an Einstein manifold with scalar curvature $R \neq 0$. Then $\text{Ker}(E)$ and $\text{Ker}(\mathcal{Q})$ coincide, i.e.

$$\text{Ker}(E) = \text{Ker}(\mathcal{Q}).$$

b) Let (M^n, g) be a connected Riemannian manifold such that $\text{Ker}(E) \cap \text{Ker}(\mathcal{Q}) \neq \{0\}$. Then (M^n, g) is an Einstein space.

Proof: Suppose first that M^n is an Einstein space, $\text{Ric}(X) = \frac{R}{n} X$. If $\psi \in \text{Ker}(E)$ we obtain from proposition 2

$$\nabla_X \psi = \frac{2(n-1)}{R(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot D\psi = -\frac{1}{n} X \cdot D\psi$$

and ψ is a twistor spinor. Conversely, if $\psi \in \text{Ker}(\mathcal{Q})$ we use the Lichnerowicz formula <4>

$$\nabla_X (D\psi) = \frac{n}{2(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot \psi$$

which reduces in an Einstein space to

$$\nabla_X (D\psi) + \frac{R}{4(n-1)} X \cdot \psi = 0.$$

This means that in an Einstein space every twistor spinor is a solution of equation (E). We consider now an arbitrary Riemannian manifold as well as a non-trivial solution $\psi \in \text{Ker}(E) \cap \text{Ker}(\mathcal{Q})$.

Using the formulas <3> and <4> we obtain the condition

$$-\frac{R}{4(n-1)} X \cdot \psi = \frac{n}{2(n-2)} \left(\frac{R}{2(n-1)} X - \text{Ric}(X) \right) \cdot \psi$$

and, finally,

$$\text{Ric}(X) \cdot \psi = \frac{R}{n} X \cdot \psi.$$

Since ψ is a twistor spinor, the zeroes of ψ are isolated points (see [3]) and therefore we conclude

$$\text{Ric}(X) = \frac{R}{n} X,$$

i.e. M^n is an Einstein space.

3. An example

Let M^2 be the simply connected Riemannian surface with constant scalar curvature R . Then M^2 admits two Killing spinors φ and ψ . According to the decomposition of the spinor bundle $S = S^+ \oplus S^-$ on M^2 we decompose the Killing spinors into $\psi = \psi^+ + \psi^-$, $\varphi = \varphi^+ + \varphi^-$. We consider the three-dimensional Riemannian manifold $M^3 = M^2 \times R^1$. The general solution of equation (E) on $M^3 = H^2 \times R^1$ (case $R < 0$) is given by

$$\begin{aligned} \psi(x, t) = & \{A_0 \cos(\tfrac{1}{2} t) + A_1 \sin(\tfrac{1}{2} t)\} \psi^+(x) + \\ & + \{A_0 \sin(\tfrac{1}{2} t) - A_1 \cos(\tfrac{1}{2} t)\} \psi^-(x) + \\ & + \{B_0 \cos(\tfrac{1}{2} t) + B_1 \sin(\tfrac{1}{2} t)\} \varphi^+(x) + \\ & + \{-B_0 \sin(\tfrac{1}{2} t) + B_1 \cos(\tfrac{1}{2} t)\} \varphi^-(x). \end{aligned}$$

A similar formula we can get on $M^3 = S^2 \times R^1$ (case $R > 0$):

$$\begin{aligned} \psi(x, t) = & \{A_0 \cosh(\tfrac{1}{2} t) + A_1 \sinh(\tfrac{1}{2} t)\} \psi^+(x) - \\ & - i \{A_0 \sinh(\tfrac{1}{2} t) + A_1 \cosh(\tfrac{1}{2} t)\} \psi^-(x) + \\ & + \{B_0 \cosh(\tfrac{1}{2} t) + B_1 \sinh(\tfrac{1}{2} t)\} \varphi^+(x) + \\ & + i \{B_0 \sinh(\tfrac{1}{2} t) + B_1 \cosh(\tfrac{1}{2} t)\} \varphi^-(x). \end{aligned}$$

References

- [1] Baum H. "Spin-Strukturen und Dirac-Operatoren über pseudo-Riemannschen Mannigfaltigkeiten", Teubner-Verlag Leipzig 1981.
- [2] Friedrich Th. "Der erste Eigenwert des Dirac-Operators einer kompakten Riemannschen Mannigfaltigkeit nichtnegativer Skalarkrümmung", Math. Nachr. 97(1980), 117-146.
- [3] Friedrich Th. "On the conformal relation between twistors and Killing spinors", to appear in Proc. Winter School Geometry and Physics, Srni, Suppl. Rend. Cive Mat. Palermo.

- [4] Lichnerowicz A. "Spin manifolds, Killing spinors and universality of the Hijazi inequality", Lett. Math. Phys. 13(1987), 331-344.
- [5] Lichnerowicz A. "Les spineurs twisteurs sur une variété spinorielle compacte", C.R. Acad. Sci. Paris Serie I 306(1988), 381-385.

THOMAS FRIEDRICH
SEKTION MATHEMATIK
HUMBOLDT-UNIVERSITÄT
1086 BERLIN, PSF 1297
G.D.R.

OLGA POKORNA
KATEDRA MATEMATIKY
MECHANIZAČNE FAKULTA
VŠZ
16000 PRAHA 6 - SUCHDOL
KAMYČKA UL.
CSSR