

Janusz Matkowski

On Nemytskii Lipschitzian operator

*Acta Universitatis Carolinae. Mathematica et Physica*, Vol. 28 (1987), No. 2, 79--82

Persistent URL: <http://dml.cz/dmlcz/701927>

**Terms of use:**

© Karolinum, Publishing House of Charles University, Prague, 1987

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## On Nemytskii Lipschitzian Operator

JANUSZ MATKOWSKI,\*

Bielsko-Biała, Poland

Received 31 March, 1987

In an earlier author's paper it has been proved that every Nemytskii operator  $N$  mapping the Banach space of Lipschitzian functions into itself and globally Lipschitzian with respect to the Lip-norm has to be of the form  $N(\varphi)(x) = A(x)\varphi(x) + B(x)$  where  $A$  and  $B$  are given Lipschitzian functions. In this paper we give a kind of local version of this result.

1. It has been proved in [3] that every Nemytskii operator  $N$  mapping  $\text{Lip}[a, b]$  into itself and globally Lipschitzian with respect to the  $\text{Lip}[a, b]$ -norm has to be of the form

$$N(\varphi)(x) = A(x)\varphi(x) + B(x), \quad x \in [a, b], \quad \text{Lip}[a, b],$$

where  $A, B \in \text{Lip}[a, b]$ . Recently this result has been extended to the Nemytskii operators mapping a normed space  $\text{Lip}(U, Y)$  into  $\text{Lip}(U, Z)$  where  $Y$  and  $Z$  are normed spaces and  $U$  is a convex (or starshaped) subset of a normed space  $X$  (cf. [4]).

Similar theorems have also been proved for the Banach spaces  $BV[a, b]$ ,  $C^r[a, b]$  and  $\text{Lip}^\alpha[a, b]$  (cf. [5], [6], [7]).

In the present paper we give a kind of local version of the above result. This "locality" is understood here in the sense of the supremum norm, i.e. a weaker one than any of the norms of Banach spaces mentioned above.

2. Let  $(X, |\cdot|)$ ,  $(Y, |\cdot|)$ ,  $(Z, |\cdot|)$  be normed spaces and let  $U \subset X$ . Denote by  $F(U, Y)$  the vector space of all functions  $\varphi: U \rightarrow Y$  and by  $\text{Lip}(U, Y)$  the vector space of all functions  $\varphi \in (U, Y)$  such that

$$\sup_{x \neq \bar{x}} \frac{|\varphi(x) - \varphi(\bar{x})|}{|x - \bar{x}|} < \infty,$$

where supremum is taken over all  $x, \bar{x} \in U$ . Assume that  $0 \in U$ . Clearly,  $\text{Lip}(U, Y)$  with the norm defined by the formula

$$(1) \quad \|\varphi\| := |\varphi(0)| + \sup_{x \neq \bar{x}} \frac{|\varphi(x) - \varphi(\bar{x})|}{|x - \bar{x}|}$$

\*) Department of Mathematics, Technical University, 43-300 Bielsko-Biała, Poland.

is a normed space. Let

$$\|\varphi\|_{\infty} := \sup_{x \in U} |\varphi(x)|, \quad \varphi \in \text{Lip}(U, Y)$$

and let  $(L(Y, Z), \|\cdot\|)$  be the normed space of all linear and continuous mappings  $A: Y \rightarrow Z$ .

Every function  $h: U \times Y \rightarrow Z$  generates the so called Nemytskii operator  $N = N_h: F(U, Y) \rightarrow F(U, Z)$  defined by the formula

$$(2) \quad N(\varphi)(x) := h(x, \varphi(x)), \quad x \in U, \quad \varphi \in F(U, Y).$$

In general it is, of course, a nonlinear operator.

We are going to prove the following

**Theorem.** *Let  $(X, |\cdot|)$ ,  $(Y, |\cdot|)$ ,  $(Z, |\cdot|)$  be normed spaces and suppose that  $U \subset X$  is star-shaped with respect to 0. If the Nemytskii operator  $N$  defined by (2) satisfies for a positive number  $r$  the following two conditions:*

$$1^{\circ}. N: \{\varphi \in \text{Lip}(U, Y): \|\varphi\|_{\infty} \leq r\} \rightarrow \text{Lip}(U, Z);$$

$$3^{\circ}. \text{ there is a } c \geq 0 \text{ such that}$$

$$(3) \quad \|N(\varphi_1) - N(\varphi_2)\| \leq c\|\varphi_1 - \varphi_2\|, \quad \varphi_i \in \text{Lip}(U, Y), \quad \|\varphi_i\|_{\infty} \leq r,$$

then there exist functions  $A: U \rightarrow L(Y, Z)$  and  $B \in \text{Lip}(U, Y)$  such that

$$(4) \quad h(x, y) = A(x)y + B(x), \quad x \in U, \quad y \in Y, \quad |y| \leq r.$$

If, moreover,  $(Y, |\cdot|)$  is a Banach space then  $A \in \text{Lip}(U, L(Y, Z))$ .

**Proof.** Since for every fixed  $y \in Y$  the constant function  $\varphi(x) = y$ ,  $x \in U$ , belongs to  $\text{Lip}(U, Y)$ , it follows from  $1^{\circ}$  that

$$h(\cdot, y) \in \text{Lip}(U, Y), \quad y \in Y, \quad |y| \leq r.$$

Therefore  $h$  is continuous with respect to the first variable for every fixed  $y$  from the ball  $B(0, r) := \{y \in Y: |y| \leq r\}$ .

Using definition (1) we may write assumption (3) in the following form

$$\begin{aligned} & |h(0, \varphi_1(0)) - h(0, \varphi_2(0))| + \\ & + \sup_{t \neq \bar{t}} \frac{|h(t, \varphi_1(t)) - h(t, \varphi_2(t)) - h(\bar{t}, \varphi_1(\bar{t})) + h(\bar{t}, \varphi_2(\bar{t}))|}{|t - \bar{t}|} \leq c\|\varphi_1 - \varphi_2\| \end{aligned}$$

where supremum is taken over all  $t, \bar{t} \in U$  and  $\|\varphi_i\|_{\infty} \leq r$ ,  $i = 1, 2$ . Hence it follows that

$$(5) \quad \frac{|h(t, \varphi_1(t)) - h(t, \varphi_2(t)) - h(\bar{t}, \varphi_1(\bar{t})) + h(\bar{t}, \varphi_2(\bar{t}))|}{|t - \bar{t}|} \leq c\|\varphi_1 - \varphi_2\|$$

for all  $\varphi_1, \varphi_2 \in \text{Lip}(U, Y)$  such that  $\|\varphi_i\|_{\infty} \leq r$ ,  $i = 1, 2$  and  $t, \bar{t} \in U$ ,  $t \neq \bar{t}$ .

Let us fix  $x \in U$ ,  $x \neq 0$ , and  $\bar{x}$  from the segment joining 0 with  $x$ . Take  $y_1, y_2, \bar{y}_1, \bar{y}_2 \in B(0, r)$  and define the functions

$$(6) \quad \varphi_i(t) := \begin{cases} \bar{y}_i & |t| < |\bar{x}| \\ \frac{y_i - \bar{y}_i}{|x| - |\bar{x}|} (|t| - |x|) + y_i, & |\bar{x}| \leq |t| \leq |x| \\ y_i & |t| > |x| \end{cases}$$

for  $t \in U$  and  $i = 1, 2$ . Evidently  $\varphi_i \in \text{Lip}(U, Y)$ ,  $\|\varphi_i\|_\infty \leq r$ ,  $i = 1, 2$ , and

$$\|\varphi_1 - \varphi_2\| = |y_1 - y_2| + \frac{|y_1 - y_2 - \bar{y}_1 + \bar{y}_2|}{|x| - |\bar{x}|}.$$

Hence, setting in (5)  $\varphi_1, \varphi_2$  defined by (6) and  $t := x, \bar{t} := \bar{x}$ , we obtain the inequality

$$\frac{|h(x, y_1) - h(x, y_2) - h(\bar{x}, \bar{y}_1) + h(\bar{x}, \bar{y}_2)|}{|x - \bar{x}|} \leq c \left( |y_1 - y_2| + \frac{|y_1 - y_2 - \bar{y}_1 + \bar{y}_2|}{|x| - |\bar{x}|} \right),$$

which can be rewritten in the following form

$$\begin{aligned} & |h(x, y_1) - h(x, y_2) - h(\bar{x}, \bar{y}_1) + h(\bar{x}, \bar{y}_2)| \leq \\ & \leq c \left( |y_1 - y_2| |x - \bar{x}| + \frac{|x - \bar{x}|}{|x| - |\bar{x}|} |y_1 - y_2 - \bar{y}_1 + \bar{y}_2| \right). \end{aligned}$$

Letting  $\bar{x}$  tend to  $x$ , using of the continuity of  $h(\cdot, y)$ , we hence get

$$(7) \quad |h(x, y_1) - h(x, y_2) - h(x, \bar{y}_1) + h(x, \bar{y}_2)| \leq c |y_1 - y_2 - \bar{y}_1 + \bar{y}_2|,$$

for  $x \neq 0, x \in U, y_1, y_2, \bar{y}_1, \bar{y}_2 \in B(0, r)$ .

By the continuity of  $h(\cdot, y)$  it follows that (7) holds for  $x = 0$ . Let us fix an  $x \in U$  and define the function  $A(x): B(0, r) \rightarrow Z$  by the formula

$$(8) \quad A(x)(y) := h(x, y) - h(x, 0).$$

Taking in (7)  $y_1 := y + w, y_2 := y, \bar{y}_1 := w, \bar{y}_2 := 0$  such that  $y, w \in B(0, r/2) \subset U$  we obtain

$$h(x, y + w) - h(x, y) - h(x, w) + h(x, 0) = 0,$$

which means that

$$A(x)(y + w) = A(x)(y) + A(x)(w), \quad y, w \in B(0, r/2),$$

i.e.  $A(x)$  is additive mapping in the ball  $B(0, r/2)$ . It is well known that  $A(x)$  has the unique extension to an additive map from  $Y$  to  $Z$  (cf. [1] and [2], Theorem 4.3). Denote this extension by  $A(x)$ . Setting  $\bar{y}_1 = \bar{y}_2 = 0$  in (7) we get

$$|A(x)(y_1) - A(x)(y_2)| \leq c |y_1 - y_2|, \quad y_1, y_2 \in B(0, r),$$

which implies the continuity of  $A(x)$ . Since every additive and continuous map is

linear we have proved that  $A(x) \in L(Y, Z)$ . Putting

$$B(x) := h(x, 0), \quad x \in U,$$

we have, according to (8),

$$h(x, y) = A(x)y + B(x), \quad x \in U, \quad y \in Y, \quad |y| \leq r,$$

where  $A \in F(U, L(Y, Z))$  and  $B \in \text{Lip}(U, Z)$ .

Suppose now that  $(Y, |\cdot|)$  is a Banach space. For every  $x, \bar{x} \in U, x \neq \bar{x}$ , we have

$$\frac{A(x) - A(\bar{x})}{|x - \bar{x}|} \in L(Y, Z).$$

From the just proved part of the theorem we have  $N(\varphi) - B = A(\cdot)y$ , for  $\varphi(x) = y$ . Consequently, for every  $y \in B(0, r)$ ,  $A(\cdot)y \in \text{Lip}(U, Z)$ , and, therefore

$$\sup_{\substack{x \neq \bar{x} \\ x, \bar{x} \in U}} \frac{|A(x)y - A(\bar{x})y|}{|x - \bar{x}|} = \sup_{\substack{x \neq \bar{x} \\ x, \bar{x} \in U}} \left| \frac{A(x) - A(\bar{x})}{|x - \bar{x}|} y \right| < \infty, \quad y \in B(0, r).$$

This shows that the family of linear maps

$$\left\{ \frac{A(x) - A(\bar{x})}{|x - \bar{x}|} \right\}_{x, \bar{x} \in U; x \neq \bar{x}}$$

is pointwise bounded. In view of Banach-Steinhaus Theorem the number

$$\sup_{\substack{x \neq \bar{x} \\ x, \bar{x} \in U}} \frac{\|A(x) - A(\bar{x})\|}{|x - \bar{x}|}$$

is finite. This completes the proof.

### References

- [1] DHOMBRES J., GER R., Conditional Cauchy Equations, *Glasnik Mat.* (1978) Vol. 13 (33), p. 39–62.
- [2] DHOMBRES, J., Some Aspects of Functional Equations, *Lecture Notes*, (1979), Chulalongkorn University, Bangkok.
- [3] MATKOWSKI J., Functional Equations and Nemytskii operator, *Funkcialaj Ekvacioj* 25 (1982), p. 127–132.
- [4] MATKOWSKI J., On Nemytskii operator, (will be published in *Mathematica Japonica*).
- [5] MATKOWSKI J., MIŠ J., On a Characterization of Lipschitzian operators of Substitution in the Space  $BV(a, b)$ , *Math. Nachr.* 117 (1984), p. 155–159.
- [6] MATKOWSKI J., Form of Lipschitz Operators of Substitution in Banach Spaces of Differentiable Functions, *Scient. Bull. Łódź Tech. Univ.*, 17 (1984), p. 5–10.
- [7] MATKOWSKA A., On Characterization of Lipschitzian Operators of Substitution in the class of Hölder's Functions, *Scient. Bull. Łódź Tech. Univ.* 17 (1984), p. 81–85.