

Lukáš Pospíšil; Zdeněk Dostál

Minimization of a convex quadratic function subject to separable conical constraints in granular dynamics

In: Jan Chleboun and Petr Přikryl and Karel Segeth and Jakub Šístek and Tomáš Vejchodský (eds.): Programs and Algorithms of Numerical Mathematics, Proceedings of Seminar. Dolní Maxov, June 8-13, 2014. Institute of Mathematics AS CR, Prague, 2015. pp. 175--180.

Persistent URL: <http://dml.cz/dmlcz/702681>

**Terms of use:**

© Institute of Mathematics AS CR, 2015

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library*  
<http://project.dml.cz>

## MINIMIZATION OF A CONVEX QUADRATIC FUNCTION SUBJECT TO SEPARABLE CONICAL CONSTRAINTS IN GRANULAR DYNAMICS

Lukáš Pospíšil, Zdeněk Dostál

FEECS VŠB-Technical University of Ostrava  
17. listopadu 15, CZ-70833 Ostrava, Czech Republic  
lukas.pospisil@vsb.cz, zdenek.dostal@vsb.cz

### Abstract

The numerical solution of granular dynamics problems with Coulomb friction leads to the problem of minimizing a convex quadratic function with semidefinite Hessian subject to a separable conical constraints. In this paper, we are interested in the numerical solution of this problem. We suggest a modification of an active-set optimal quadratic programming algorithm. The number of projection steps is decreased by using a projected Barzilai-Borwein method. In the numerical experiment, we compare our algorithm with Accelerated Projected Gradient method and Spectral Projected Gradient method on the solution of a particle dynamics problem with hundreds of spherical bodies and static obstacles.

### 1. Time-stepping scheme and formulation of optimization problem

In our simulation, we consider a system of  $nb \in \mathbb{N}$  particles in vector space  $\{(x, y, z) \in \mathbb{R}^3\}$ . The position of each particle in time  $t$  is defined by the vector of generalized position  $q_i^{(t)} \in \mathbb{R}^7$ , which consists of the position of the centre of gravity  $[r_x, r_y, r_z]^T$  and the unit quaternion of rotation  $[e_0, e_1, e_2, e_3]^T$ . The velocity of the body is defined by the vector of generalized velocities  $v_i^{(t)} \in \mathbb{R}^6$ , it includes the velocity corresponding to the position of the centre of the body and angular velocities represented in Euler angles.

We use the well-known time-stepping scheme, see Heyn [9] or Heyn et al. [10]

$$\begin{aligned} \mathbf{q}^{(t+h)} &= \mathbf{q}^{(t)} + hQ\mathbf{v}^{(t)}, \\ \mathbf{v}^{(t+h)} &= \mathbf{v}^{(t)} + hM^{-1}(\mathbf{F}_{ext} + \mathbf{F}_C), \end{aligned} \tag{1}$$

where  $h$  is a time step,  $Q$  denotes the matrix of linear mapping between the derivative of the position vector and the vector of velocities,  $M$  is a generalized mass matrix,  $\mathbf{F}_C$  is a vector of forces induced by contact constraints, and  $\mathbf{F}_{ext}$  is a vector of external forces. In our simulation, the vector of external forces represents the gravity force applied to each body.

Let us denote the number of contacts by  $m \in \mathbb{N} \cup \{0\}$ . The contact force applied to each body can be separated into the sum of the normal force and the tangential force, i.e.,

$$\mathbf{F}_C = \mathbf{F}_n + \mathbf{F}_T = \gamma_n \mathbf{n} + \gamma_u \mathbf{u} + \gamma_w \mathbf{w} ,$$

where  $\gamma_n > 0$  is the size of the normal component of the friction force, and  $\gamma_u, \gamma_w \in \mathbb{R}$  are the sizes of the tangential components of the friction force. Here,  $\{\mathbf{n}, \mathbf{u}, \mathbf{w}\}$  is an orthonormal basis of the tangential space at the contact point. The relation between the components of  $\boldsymbol{\gamma}_j := [\gamma_n, \gamma_u, \gamma_w]$  for  $j$ -th contact ( $j = 1, \dots, m$ ) can be described by the *Coulomb friction model*. The unknown vector of all components in all contacts can be denoted by  $\boldsymbol{\gamma} := [\gamma_1, \dots, \gamma_m] \in \mathbb{R}^{3m}$  and can be found by solving the problem of minimizing a convex quadratic function subject to separable conical constraints (see Heyn [9]). The proof of equivalency is based on the maximum dissipation principle and duality.

The optimization problem is given by

$$\text{find } \boldsymbol{\gamma} := \arg \min_{\boldsymbol{\gamma} \in \Omega} f(\boldsymbol{\gamma}), \quad f(\boldsymbol{\gamma}) := \frac{1}{2} \boldsymbol{\gamma}^T \mathbf{A} \boldsymbol{\gamma} - \mathbf{b}^T \boldsymbol{\gamma} , \quad (2)$$

where  $\mathbf{A} \in \mathbb{R}^{3m, 3m}$  is a symmetric positive semidefinite matrix,  $\mathbf{b} \in \mathbb{R}^{3m}$ , and  $\Omega \subset \mathbb{R}^{3m}$  is a non-empty convex feasible set defined by separable conical constraints

$$\Omega := \{ \boldsymbol{\gamma} \in \mathbb{R}^{3m} : h_j(x_{2j-2}, x_{2j-1}, x_{2j}) \leq 0, j = 1, \dots, m \} ,$$

where  $h_j : \mathbb{R}^3 \rightarrow \mathbb{R}$  are conical constraints functions

$$h_j(x, y, z) := \sqrt{y^2 + z^2} - \mu_j x, \quad j = 1, \dots, m ,$$

and  $\mu_j \geq 0$  are given friction coefficients that define the interior angles of cones. Let us notice, that if we consider the problem without friction, then  $\mu_j = 0$ , and the optimization problem (2) becomes a quadratic programming problem with bound constraints.

For the sake of simplicity we denote the triplet of components of  $\boldsymbol{\gamma} \in \mathbb{R}^{3m}$  constrained by  $j$ -th constraint function using the notation of index sets

$$\mathcal{I}_j := \{3j - 2, 3j - 1, 3j\}, \quad \bigcup_{j=1}^m \mathcal{I}_j = \{1, \dots, 3m\}, \quad j = 1, \dots, m .$$

## 2. Active-set method

For numerical solution of the problem (2), we are using the variant of Modified Proportioning with Gradient Projection (MPGP), see Dostál [5] and Dostál et al. [7, 4], or Pospíšil [12]. This active-set algorithm is based on the decomposition of the set of all constraint indices  $\mathcal{M} := \{1, \dots, 3m\}$  into two disjoint subsets based on the values of constraint functions

$$\begin{aligned} \mathcal{F}(\boldsymbol{\gamma}) &:= \{j \in \mathcal{M} : h_j(\boldsymbol{\gamma}_{\mathcal{I}_j}) < 0\} , \\ \mathcal{A}(\boldsymbol{\gamma}) &:= \{j \in \mathcal{M} : h_j(\boldsymbol{\gamma}_{\mathcal{I}_j}) = 0\} . \end{aligned}$$

The gradient of the objective function  $\mathbf{g} := \nabla f(\mathbf{x}) = A\mathbf{x} - b \in \mathbb{R}^n$  can be used to define the *free* and the *chopped* gradient with components

$$\begin{aligned} \varphi_{\mathcal{I}_j}(\mathbf{x}) &= \mathbf{g}_{\mathcal{I}_j} \text{ for } j \in \mathcal{F}(\mathbf{x}), & \varphi_{\mathcal{I}_j}(\mathbf{x}) &= 0 \text{ for } j \in \mathcal{A}(\mathbf{x}), \\ \beta_{\mathcal{I}_j}(\mathbf{x}) &= 0 \text{ for } j \in \mathcal{F}(\mathbf{x}), & \beta_{\mathcal{I}_j}(\mathbf{x}) &= \mathbf{g}_{\mathcal{I}_j} - \min\{n_j^T(\mathbf{x}_{\mathcal{I}_j})\mathbf{g}_{\mathcal{I}_j}, 0\}n_j(\mathbf{x}_{\mathcal{I}_j}) \\ & & & \text{for } j \in \mathcal{A}(\mathbf{x}), \end{aligned}$$

where  $n_j(x, y, z)$  is the unit outer normal of  $j$ -th constraint  $h_j(x, y, z)$ . We consider a problem with conical constraints, so outer normal is given by

$$n_j(x, y, z) := \begin{cases} [-1, 0, 0]^T & \text{if } x = y = z = 0, \\ [-\mu_j, y/\sqrt{y^2 + z^2}, z/\sqrt{y^2 + z^2}]^T & \text{elsewhere.} \end{cases}$$

**Algorithm 1: Modified Proportioning with Barzilai-Borwein Gradient Projection (MPGPS-BB).**

```

Choose  $\mathbf{x}^0 \in \Omega$ 
for  $k = 0, 1, 2, \dots$  (while a stopping criterion is not achieved)
  if  $\|\varphi(x_k)\| \geq \|\beta(x_k)\|$  (proportioning condition)
    Control the solvability
    if  $\min\{\alpha_f, \alpha_{cg}\} = \infty$ , then the problem has no solution.
    CG step or CG halfstep
    make one CG step to solve problem on free set
    if this step means leaving  $\Omega$ , do only a half-step and restart CG
  else
    Gradient projection step.
    make projected Barzilai-Borwein step
    restart CG on free set
  endif
   $k := k + 1$ 
endfor

```

Our algorithm is based on using the free and chopped gradient to minimize the objective function on the free set and afterwards on the active set. The switching between these processes is realized by the proportioning condition. The implementation details of each step are the same as in the original Modified Proportioning with Gradient Projections algorithm (MPGP) in Dostál [5], Dostál et al. [7, 4]. Nevertheless, MPGP was developed to solve the problems with a symmetric positive definite Hessian matrix. The recent generalization to the problems with symmetric positive semidefinite Hessian suggests only one difference from the original algorithm,

specifically a test of the problem solvability, see Algorithm 1. The coefficient  $\alpha_f$  is the maximal feasible step-size and  $\alpha_{cg}$  is a coefficient of the conjugate gradient computed from the free gradient. If both of these coefficient are equal to infinity, then the problem has no solution. The theory will be published in [6].

To solve a problem with separable conical constraints, we suggest to use the projected version of Barzilai-Borwein method [2] instead of the projected gradient method with constant step-length as in original MGP algorithm. Constant step-length always induces the descend of cost function, as it was shown by Dostál and Schöberl [8]. However, the numerical experiments show that using non-monotone algorithms, such as projected Barzilai-Borwein (PBB) given by

$$\mathbf{x}^{k+1} = P_{\Omega}(\mathbf{x}^k - \alpha_k^{BB} \nabla f(\mathbf{x}^k)), \quad \alpha_k^{BB} = \frac{\mathbf{s}_k^T \mathbf{s}_k}{\mathbf{s}_k^T A \mathbf{s}_k}, \quad \mathbf{s}_k = \mathbf{x}^k - \mathbf{x}^{k-1},$$

usually evokes the decrease of the projection steps number. This modification was inspired by the Spectral Projected Gradient method (SPG), which uses the similar type of steps, see Birgin et al. [3]. The idea of the combination of MGP and PBB was firstly presented by Pospíšil [12] and tested on the problem with separable quadratic constraints.

The main shortage of the presented MGP-SBB algorithm is the absence of the proof of convergence. The PBB method is non-monotone and hardly analyzable. Therefore, the SPG method is using an additional line-search method to control the descend of the objective function, i.e. the global convergence. In our algorithm, we tried to omit this line-search. Our idea is well-founded by the numerical experiment presented in the final section of this paper.

As a stopping criterion in our algorithm, we are using the norm of the *scaled projected gradient* defined by

$$\tilde{\mathbf{g}}_{\alpha}^P(\mathbf{x}) := \frac{1}{\alpha}(\mathbf{x} - P_{\Omega}(\mathbf{x} - \alpha \nabla f(\mathbf{x}))) .$$

The equivalency of this gradient and the fulfilment of Karush-Kuhn-Tucker optimality conditions for problems with feasible sets with strong curvature was discussed and proved by Bouchala et al. [4].

### 3. Numerical experiments

In this section, we present the numerical results showing the efficiency of our algorithm on the simulation of 339 spherical particles with friction. In our benchmark, the particles are scattered into simple box represented by six walls. The initial position of the partices and final position can be found in Fig. 1, where we depicted only the partices and the bottom side of the static box. The material of the bodies is represented by density  $\rho = 730 \text{ kg.m}^{-3}$  and friction parameter  $\mu = 0.3$ . The stepsize of the time-stepping scheme is  $h = 6.25 \cdot 10^{-4} \text{ s}$ .




Figure 1: State of testing benchmark at  $t = 0s$  (left) and  $t = 5s$  (right).

<b>t</b>	<b>contacts</b>	<b>n</b>	<b>active</b>	<b>MPGPS-BB</b>	<b>SPG</b>	<b>APGD</b>
1s	738	2214	562 (76%)	274 (7.6s)	2360 (37.8s)	754 (9.4s)
2s	702	2106	574 (82%)	137 (3.4s)	449 (6.0s)	346 (2.9s)
3s	730	2190	558 (76%)	137 (3.7s)	449 (6.0s)	346 (4.1s)
4s	814	2442	640 (79%)	338 (9.9s)	2931 (56.2s)	1345 (18.9s)
5s	818	2454	652 (80%)	425 (12.3s)	4176 (88.0s)	1742 (25.6s)

Table 1: The optimization problems at selected times of the simulation; number of contacts, dimension of the problem, the number of iterations and computing time of the algorithms.

We compare our algorithm with SPG and the Accelerated Projected Gradient Descend method (APGD [11]). In SPG, because the minimum of the quadratic function in a given direction is known, we use the Cauchy step-size instead of using an additional Grippo-Lampariello-Lucidi line-search. All algorithms were implemented in the Matlab environment. For contact detection, we are using our own implementation of the *Moving Bounding-Box algorithm* [13]. The number of iterations at selected times of the simulation can be found in Table 1. We demand the relative stopping tolerance  $\|\tilde{\mathbf{g}}_{\alpha}^P(\mathbf{x})\| < \epsilon\|b\|$ ,  $\epsilon = 10^{-6}$ .

#### 4. Conclusions

In our paper, we proposed the modification of our active-set algorithm for the solution of optimization problem in particle dynamics with friction. Our numerical experiment shows the efficiency of the modifications. Unfortunately, the basic disadvantage of using the projected Barzilai-Borwein method is the absence of a convergence proof as well as of an estimate of the speed of convergence.

#### 5. Acknowledgements

This work was supported by the European Regional Development Fund in the IT4Innovations Centre of Excellence project (CZ.1.05/1.1.00/02.0070) and project SGS SP2014/204.

## References

- [1] Anitescu, M.: Optimization-based simulation of nonsmooth rigid multibody dynamics. *Math. Program.* **105** (2006), 113–143.
- [2] Barzilai, J. and Borwein, J. M.: Two point step size gradient methods. *IMA J. Numer. Anal.* **8** (1988), 141–148.
- [3] Birgin, E. G., Martinez, J. M., and Raydan, M.: Nonmonotone spectral projected gradient methods on convex sets. *SIAM J. Optim.* **10** (2000), 1196–1211.
- [4] Bouchala, J., Dostál, Z., Kozubek, T., Pospíšil, L., and Vodstrčil, P.: On the solution of convex QPQC problems with elliptic and other separable constraints. *Appl. Math. Comput.* **247** (2014), 848–864.
- [5] Dostál, Z.: *Optimal quadratic programming algorithms, with applications to variational inequalities*, 1st edition. SOIA 23. Springer US, New York, 2009.
- [6] Dostál, Z. and Pospíšil, L.: Minimization of the quadratic function with semidefinite Hessian subject to the bound constraints. in preparation.
- [7] Dostál, Z. and Pospíšil, L.: Optimal iterative QP and QPQC algorithms. *Ann. Oper. Res.* (2013).
- [8] Dostál, Z., and Schöberl, J.: Minimizing quadratic functions subject to bound constraints with the rate of convergence and finite termination. *Comput. Optim. Appl.* **30** (2005), 23–44.
- [9] Heyn, T.: *On the modeling, simulation, and visualization of many-body dynamics problems with friction and contact*. Ph.D. Thesis, University of Wisconsin-Madison, 2013.
- [10] Heyn, T., Anitescu, M., Tasora, A., and Negrut, D.: Using Krylov subspace and spectral methods for solving complementarity problems in many-body contact dynamics simulation. *Internat. J. Numer. Methods Engrg.* **95** (2012), 541–561.
- [11] Nesterov, Y.: *Introductory lectures on convex optimization: a basic course*. Volume 87, Springer, 2003.
- [12] Pospíšil, L.: An optimal algorithm with Barzilai-Borwein steplength and super-relaxation for QPQC problem. In: J. Chleboun, K. Segeth, J. Šístek, T. Vejchodský (Eds.), *Proceedings of Programs and Algorithms of Numerical Mathematics 16*, pp. 155–161. IM ASCR, Prague, 2012.
- [13] Schinner, A.: Fast algorithms for the simulations of polygonal particles. *Springer-Verlag Granular Matter* **2** (1999), 35–43.