

Jing Cheng Tong

The best Diophantine approximation functions by continued fractions

Mathematica Bohemica, Vol. 121 (1996), No. 1, 89–94

Persistent URL: <http://dml.cz/dmlcz/125943>

Terms of use:

© Institute of Mathematics AS CR, 1996

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

THE BEST DIOPHANTINE APPROXIMATION FUNCTIONS
BY CONTINUED FRACTIONS

JINGCHENG TONG, Jacksonville

(Received September 14, 1994)

Summary. Let $\xi = [a_0; a_1, a_2, \dots, a_i, \dots]$ be an irrational number in simple continued fraction expansion, $p_i/q_i = [a_0; a_1, a_2, \dots, a_i]$, $M_i = q_i^2 |\xi - p_i/q_i|$. In this note we find a function $G(R, r)$ such that

$$\begin{aligned} M_{n+1} < R \text{ and } M_{n-1} < r &\text{ imply } M_n > G(R, r), \\ M_{n+1} > R \text{ and } M_{n-1} > r &\text{ imply } M_n < G(R, r). \end{aligned}$$

Together with a result the author obtained, this shows that to find two best approximation functions $\tilde{H}(R, r)$ and $\tilde{L}(R, r)$ is a well-posed problem. This problem has not been solved yet.

Keywords: continued fraction, diophantine approximation

AMS classification: 11J04, 11A55

1. INTRODUCTION

In 1891, Hurwitz [4] proved the following basic fact in the theory of Diophantine approximation:

Theorem 1. *Let ξ be an irrational number. Then there are infinitely many reduced fractions p/q such that $|\xi - p/q| < 1/(\sqrt{5}q^2)$.*

In order to find these fractions p/q , Borel [2] found the theorem below.

Theorem 2. *Let ξ be an irrational number with simple continued fraction expansion $\xi = [a_0; a_1, a_2, \dots, a_i, \dots]$. If $p_i/q_i = [a_0; a_1, a_2, \dots, a_i]$, then at least one of the three consecutive p_i/q_i ($i = n-1, n, n+1$) satisfies $|\xi - p_i/q_i| < 1/(\sqrt{5}q_i^2)$.*

For nearly a century, Borel's theorem has been extensively investigated. There are numerous results about it [1, 3, 6, 7, 10, 11]. The following result of the present author [10] is one of the highlights since it contains asymmetric approximations [5, 9] as a special case.

Theorem 3. *Let ξ be an irrational number with simple continued fraction expansion $\xi = [a_0; a_1, a_2, \dots, a_i, \dots]$. If $M_i = q_i^2 |\xi - p_i/q_i|$ and r is a real number exceeding a_{n+1} , then*

$$\begin{aligned} M_n < r &\text{ implies } \max(M_{n-1}, M_{n+1}) > 4r/(r^2 - a_{n+1}^2), \\ M_n > r &\text{ implies } \min(M_{n-1}, M_{n+1}) < 4r/(r^2 - a_{n+1}^2). \end{aligned}$$

Recently the author generalized Theorem 3 to the inhomogeneous case [12].

Theorem 4. *Let R, r be two real numbers exceeding $1+1/a_{n+1}$, let M_i be defined as in Theorem 3. Then*

$$\begin{aligned} M_{n+1} < R \text{ and } M_{n-1} < r &\text{ imply } M_n > F(R, r), \\ M_{n+1} > R \text{ and } M_{n-1} > r &\text{ imply } M_n < F(R, r), \end{aligned}$$

where $F(R, r) = \left(R^{-1} + r^{-1} + \sqrt{a_{n+1}^2 + 4/(Rr)} \right) / \left(1 - (R^{-1} - r^{-1})^2/a_{n+1}^2 \right)$.

Theorem 3 is a special case of Theorem 4 for $R = r$.

In this paper we give another estimation function $G(R, r)$ to replace $F(R, r)$ in Theorem 4. Using $G(R, r)$, the proof is simpler, it also gives Theorem 3 for $R = r$, the approximation is better than $F(R, r)$ for many values of R, r . These advantages are not essentially important, the significance of introducing $G(R, r)$ is the following consideration.

Suppose \mathcal{H} is the set of all functions $H(R, r)$ such that

$$M_{n+1} < R \text{ and } M_{n-1} < r \text{ imply } M_n > H(R, r),$$

and \mathcal{L} is the set of all functions $L(R, r)$ such that

$$M_{n+1} > R \text{ and } M_{n-1} > r \text{ imply } M_n < L(R, r).$$

Then $\tilde{H}(R, r) = \sup \mathcal{H}$ and $\tilde{L}(R, r) = \inf \mathcal{L}$ are the best approximation functions. Theorem 4 guarantees that both \mathcal{H} and \mathcal{L} are not empty sets. Nevertheless, it could be true that $\tilde{H}(R, r) = F(R, r)$ and $\tilde{L}(R, r) = F(R, r)$. The function $G(R, r)$ which we are going to introduce satisfies $G(R, r) \neq F(R, r)$. This ensures that to find the best approximation functions $\tilde{H}(R, r)$ and $\tilde{L}(R, r)$ is a well-posed problem. These two functions have not been found yet. It is still a challenging Problem.

2. PRELIMINARIES

Let $\xi = [a_0; a_1, a_2, \dots, a_i, \dots]$ be an irrational number in simple continued fraction expansion, $p_i/q_i = [a_0; a_1, a_2, \dots, a_i]$ and $M_i = q_i^2|\xi - p_i/q_i|$. Then it is known [8] that

$$(1) \quad M_i = [a_{i+1}; a_{i+2}, \dots] + [0; a_i, a_{i-1}, \dots, a_1].$$

Let $P = [a_{n+2}; a_{n+3}, \dots]$, $Q = [a_n; a_{n-1}, \dots, a_1]$. Then it is easily checked that the following relations are true.

$$(2) \quad M_{n+1} = P + (a_{n+1} + Q^{-1})^{-1},$$

$$(3) \quad M_n = a_{n+1} + P^{-1} + Q^{-1},$$

$$(4) \quad M_{n-1} = Q + (a_{n+1} + P^{-1})^{-1}.$$

Now we turn to some inequalities.

If $M_{n+1} < R$, then $P < R - (a_{n+1} + Q^{-1})^{-1}$.

If $M_{n-1} < r$, then $Q < r - (a_{n+1} + P^{-1})^{-1}$.

If both $M_{n+1} < R$ and $M_{n-1} < r$ are true, then by (3) we know that the following two inequalities are correct.

$$(5) \quad M_n > a_{n+1} + P^{-1} + (r - (a_{n+1} + P^{-1})^{-1})^{-1},$$

$$(6) \quad M_n > a_{n+1} + Q^{-1} + (r - (a_{n+1} + Q^{-1})^{-1})^{-1}.$$

We will use these inequalities in the next section.

3. THE MAIN THEOREM

We need a useful lemma.

Lemma 1. *The function $f(P) = a + P^{-1} + (r - (a + P^{-1})^{-1})^{-1}$ is decreasing in P for $P^{-1} > 2r^{-1} - a$, where $a \geq 1$, $r > 1$ are constants.*

Proof. Since $f'(P) = -P^{-2}(1 - (ra + rP^{-1} - 1)^{-2})$, if $P^{-1} > 2r^{-1} - a$, we have $ra + rP^{-1} - 1 > 1$, hence $f'(P) < 0$. \square

Theorem 5. *Let M_i be defined as in Theorem 3. Then*

$$M_{n+1} < R \text{ and } M_{n-1} < r \text{ imply } M_n > G(R, r),$$

$$M_{n+1} > R \text{ and } M_{n-1} > r \text{ imply } M_n < G(R, r),$$

where

$$G(R, r) = a_{n+1} + 4a_{n+1} / \left((3r - R)a_{n+1} - 4 + \sqrt{((R+r)a_{n+1})^2 + 16} \right) \\ + \left(\sqrt{((R+r)a_{n+1})^2 + 16} + (R+r)a_{n+1} + 4 \right) / (2(R+r)).$$

Proof. If $M_{n+1} < R$ and $M_{n-1} < r$, then $M_{n+1} + M_{n-1} < R + r$. From (2), (4) we have one of the following inequalities:

$$(7) \quad P + (a_{n+1} + P^{-1})^{-1} < \frac{1}{2}(R+r),$$

$$(8) \quad Q + (a_{n+1} + Q^{-1})^{-1} < \frac{1}{2}(R+r).$$

Suppose (7) holds. Then

$$a_{n+1}P^2 + (2 - \frac{1}{2}(R+r)a_{n+1})P - \frac{1}{2}(R+r) = 0.$$

Hence

$$(9) \quad P < \frac{(R+r)a_{n+1} - 4 + \sqrt{((R+r)a_{n+1})^2 + 16}}{4a_{n+1}}.$$

It is obvious that $1 > 2/(a_{n+1} + P^{-1} + 1)$, hence

$$a_{n+1} + P^{-1} > 2/(1 + (a_{n+1} + P^{-1})^{-1}).$$

Since $1 + (a_{n+1} + P^{-1})^{-1} \leq Q + (a_{n+1} + P^{-1})^{-1} = M_{n-1} < r$, we have $P^{-1} > 2r^{-1} - a_{n+1}$. By (5), (9) and Lemma 1 we have

$$M_n > G(R, r).$$

Suppose (8) holds. Noticing (6), we can have $M_n > G(R, r)$ by the similar discussion.

Reversing all the directions of the inequalities above, we can prove that

$$M_{n+1} > R \text{ and } M_{n-1} > r \text{ imply } M_n < G(R, r).$$

□

4. COMPARISON OF THE APPROXIMATION FUNCTIONS $F(R, r)$ AND $G(R, r)$

It is easily checked that

$$F(r, r) = G(r, r) = (2 + \sqrt{(ra_{n+1})^2 + 4})/r.$$

The function $F(R, r)$ is a symmetric function while $G(R, r)$ is not.

If $a_{n+1} = 1$, then

$$F(4, 2) = 2.1063946,$$

$$G(4, 2) = 2.2018503.$$

In this case $G(R, r)$ gives a better estimation than $F(R, r)$ in the set \mathcal{H} , while $F(R, r)$ gives a better estimation than $G(R, r)$ in the set \mathcal{L} .

If $a_{n+1} = 3$, then

$$F(2, 6) = 3.7682386,$$

$$G(2, 6) = 3.4463062.$$

In this case the situation reverses.

The function $H(R, r) = \max(F(R, r), G(R, r))$ is a better approximation than both $F(R, r)$ and $G(R, r)$ in the set \mathcal{H} .

The function $L(R, r) = \min(F(R, r), G(R, r))$ is a better approximation than both $F(R, r)$ and $G(R, r)$ in the set \mathcal{L} .

The best approximation functions satisfy $\tilde{H}(R, r) \geq \tilde{L}(R, r)$ and $\tilde{H}(R, r) \neq \tilde{L}(R, r)$. Now it is an open problem how to find them.

References

- [1] *F. Bagemihl and J. R. McLaughlin*: Generalization of some classical theorems concerning triples of consecutive convergents to simple continued fractions. *J. Reine Angew. Math.* **221** (1966), 146–149.
- [2] *E. Borel*: Contribution a l'analyse arithmetique du continu. *J. Math. Pures Appl.* **9** (1903), 329–375.
- [3] *M. Fujiwara*: Bemerkung zur Theorie der Approximation der irrationalen Zahlen durch rationale Zahlen. *Tôhoku Math. J.* **14** (1918), 109–115.
- [4] *A. Hurwitz*: Über die angenäherte Darstellung der irrationalzahlen durch rationale Brüche. *Math. Ann.* **39** (1891), 279–284.
- [5] *W. J. LeVeque*: On asymmetric approximations. *Michigan Math. J.* **2** (1953), 1–6.
- [6] *M. Müller*: Über die Approximation reeller Zahlen durch die Näherungsbrüche ihres regelmässigen Kettenbruches. *Arch. Math. (Basel)* **6** (1955), 253–258.
- [7] *M. B. Nathanson*: Approximation by continued fractions. *Proc. Amer. Math. Soc.* **45** (1974), 323–324.

- [8] *O. Perron*: Die Lehre von den Kettenbrüchen I, II. 3rd ed. Teubner, Leipzig, 1954.
- [9] *B. Segre*: Lattice points in the infinite domains and asymmetric Diophantine approximation. *Duke Math. J.* *12* (1945), 337–365.
- [10] *J. Tong*: The conjugate property for Diophantine approximation of continued fraction. *Proc. Amer. Math. Soc.* *105* (1989), 535–539.
- [11] *J. Tong*: Diophantine approximation of a single irrational number. *J. Number Theory* *34* (1990), 53–57.
- [12] *J. Tong*: Diophantine approximation by continued fractions. *J. Austral. Math. Soc. Ser. A* *51* (1991), 324–330.

Author's address: Jingcheng Tong, Department of Mathematics and Statistics, University of North Florida, Jacksonville, FL 32224, U.S.A.