

Danica Jakubíková-Studenovská

Representation of connected monounary algebras by means of irreducibles

Czechoslovak Mathematical Journal, Vol. 55 (2005), No. 4, 1067–1078

Persistent URL: <http://dml.cz/dmlcz/128045>

Terms of use:

© Institute of Mathematics AS CR, 2005

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

REPRESENTATION OF CONNECTED MONOUNARY ALGEBRAS
BY MEANS OF IRREDUCIBLES

DANIČA JAKUBÍKOVÁ-STUDENOVSKÁ, Košice

(Received April 18, 2003)

Abstract. The aim of the present paper is to describe all connected monounary algebras for which there exists a representation by means of connected monounary algebras which are retract irreducible in the class \mathcal{U}_c (or in \mathcal{U}).

Keywords: monounary algebra, connectedness, retract, retract irreducibility, representation

MSC 2000: 08A60

0. INTRODUCTION

The relations between retracts and direct product decompositions have been investigated by Duffus and Rival [1] (for ordered sets), by Imrich and Klavžar [2] and by Klavžar [11] (for graphs) and by Kuczmanov, Reich, Schmidt and Stachura [12] (for metric spaces).

The author [5]–[10] dealt with these relations for monounary algebras. Let us denote by \mathcal{U} (and \mathcal{U}_c , respectively) the class of all (all connected) monounary algebras.

In [5] and [4] all connected monounary algebras which are retract irreducible in the class \mathcal{U}_c were described. Next, in [6] all connected monounary algebras retract irreducible in the class \mathcal{U} were found. In [10] it was proved that there exist connected monounary algebras which are not representable by means of connected monounary algebras which are retract irreducible in \mathcal{U}_c . An analogous result was shown for the representability in \mathcal{U} .

This work was supported by the Slovak Grant VEGA 1/0423/03.

The aim of the present paper is to describe all connected monounary algebras for which there exists a representation by means of connected monounary algebras which are retract irreducible in \mathcal{U}_c (or in \mathcal{U}).

1. PRELIMINARIES

Let $\underline{A} = (A, F)$ be an algebra. A subalgebra $\underline{B} = (B, F)$ of \underline{A} is called a retract of \underline{A} if there is an endomorphism φ of \underline{A} such that $\varphi(b) = b$ for each $b \in B$. For an algebra \underline{A} let $R(\underline{A})$ be the class of all algebras which are isomorphic to a retract of \underline{A} .

Let \mathcal{K} be a class of algebras of the same type. If $\underline{A} \in R\left(\prod_{i \in I} \underline{A}_i\right)$, $\underline{A}_i \in \mathcal{K}$ for each $i \in I$, then the system $\{\underline{A}_i\}_{i \in I}$ is called a representation of \underline{A} in the class \mathcal{K} ; we say that \underline{A} is representable by means of $\{\underline{A}_i\}_{i \in I}$.

The algebra \underline{A} is said to be retract irreducible in a class of algebras \mathcal{K} if the following condition is satisfied:

$$(1) \quad \text{If } \underline{A} \in R\left(\prod_{i \in I} \underline{A}_i\right), \underline{A}_i \in \mathcal{K} \text{ for each } i \in I, \\ \text{then there is } j \in I \text{ such that } \underline{A} \in R(\underline{A}_j).$$

We will say that \underline{A} is representable by means of irreducibles in \mathcal{K} if there is a representation $\{\underline{A}_i\}_{i \in I}$ of \underline{A} such that each \underline{A}_i is retract irreducible in \mathcal{K} .

Let us remark that if $\{\underline{A}_i\}_{i \in I}$ is a representation of \underline{A} in \mathcal{K} and if for each $i \in I$ the system $\{\underline{A}_{ij_i}\}_{j_i \in J_i}$ is a representation of \underline{A}_i in \mathcal{K} , then the system

$$\{\underline{A}_{ij_i}\}_{i \in I, j_i \in J_i}$$

is a representation of \underline{A} , too.

In what follows we will apply the previous notion to the case of monounary algebras.

For the sake of completeness we recall some basic notions concerning monounary algebras.

Let $\underline{A} = (A, f)$ be a monounary algebra. As usual, a nonempty subset M of A is said to be a retract of \underline{A} if there is a mapping h of A onto M such that h is an endomorphism of \underline{A} and $h(x) = x$ for each $x \in M$. The mapping h is then called a retraction endomorphism corresponding to the retract M . Let us remark that then $\underline{M} = (M, f)$ is a subalgebra of \underline{A} . (In fact, for each subalgebra \underline{A}_1 of $\underline{A} = (A, f)$, the corresponding operation in \underline{A}_1 is denoted by the same symbol f .)

Let \mathbb{Z} be the set of all integers, \mathbb{N} the set of all positive integers and $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. A connected monounary algebra \underline{A} is called unbounded if

$$(\forall x \in A)(\forall n \in \mathbb{N})(\exists m \in \mathbb{N})(f^{-(n+m)}(f^m(x)) \neq \emptyset).$$

The notion of the degree $s_f(x)$ of an element $x \in A$ was introduced in [13] (cf. also [3]) as follows. Let us denote by $A^{(\infty)}$ the set of all elements $x \in A$ such that there exists a sequence $\{x_n\}_{n \in \mathbb{N} \cup \{0\}}$ of elements belonging to A with the property $x_0 = x$ and $f(x_n) = x_{n-1}$ for each $n \in \mathbb{N}$. Further, we put $A^{(0)} = \{x \in A : f^{-1}(x) = \emptyset\}$. Now we define a set $A^{(\lambda)} \subseteq A$ for each ordinal λ by induction. Assume that we have defined $A^{(\alpha)}$ for each ordinal $\alpha < \lambda$. Then we put

$$A^{(\lambda)} = \left\{ x \in A - \bigcup_{\alpha < \lambda} A^{(\alpha)} : f^{-1}(x) \subseteq \bigcup_{\alpha < \lambda} A^{(\alpha)} \right\}.$$

The sets $A^{(\lambda)}$ are pairwise disjoint. For each $x \in A$, either $x \in A^{(\infty)}$ or there is an ordinal λ with $x \in A^{(\lambda)}$. In the former case we put $s_f(x) = \infty$, in the latter we set $s_f(x) = \lambda$. We put $\lambda < \infty$ for each ordinal λ .

The following assertions are consequences of the definition of $s_f(x)$ (cf. also [14]) and we will sometimes use them without further reference.

(T1) If $s_f(x) \neq \infty$, then $s_f(f(x)) > s_f(x)$.

(T2) If h is a homomorphism of (A, f) into (B, g) , then $s_g(h(x)) \geq s_f(x)$ for each $x \in A$.

(T3) Let $\{(A_i, f_i) : i \in I\}$ be a system of monounary algebras, $(A, f) = \prod_{i \in I} (A_i, f_i)$. If $z \in \prod_{i \in I} A_i$, then $s_f(z) \leq s_{f_i}(z(i))$ for each $i \in I$.

Further, by [5], Thm. 1.3, in which a characterization of retracts was given, we obtain

1.1. Lemma. *Let $\underline{A} = (A, f)$ be a monounary algebra and let $\underline{M} = (M, f)$ be a subalgebra of \underline{A} . Then M is a retract of \underline{A} if and only if the following conditions are satisfied:*

- (a) *if $y \in f^{-1}(M)$, then there is $z \in M$ such that $f(y) = f(z)$ and $s_f(y) \leq s_f(z)$;*
- (b) *for any connected component K of \underline{A} with $K \cap M = \emptyset$ there is a homomorphism of \underline{K} into \underline{M} .*

1.2. Notation. For $k \in \mathbb{N}$ we denote by $\mathbb{Z}_k = \{0_k, 1_k, \dots, (k-1)_k\}$ the set of all integers modulo k . Let us consider the following five types of monounary algebras:

- a) $\underline{\mathbb{Z}} = (\mathbb{Z}, f)$, where $f(n) = n + 1$ for each $n \in \mathbb{Z}$;
- b) $\underline{\mathbb{N}} = (\mathbb{N}, f)$, where $f(n) = n + 1$ for each $n \in \mathbb{N}$;
- c) $\underline{\mathbb{N}}_\infty = (\mathbb{N}, f)$, where $f(n) = n - 1$ for each $n \in \mathbb{N} - \{1\}$, $f(1) = 1$;
- d) $\underline{\mathbb{Z}}_k = (\mathbb{Z}_k, f)$, where $f(n_k) = (n + 1)_k$ for each $n_k \in \mathbb{Z}_k$;
- e) $\underline{\mathbb{N}}_k$ is a subalgebra of $\underline{\mathbb{N}}_\infty$, where $\mathbb{N}_k = \{1, 2, \dots, k\}$.

Let \mathcal{U} and \mathcal{U}_c be as above. Connected monounary algebras which are retract irreducible in the above classes were characterized in [4], [5] and [6] as follows:

1.3. Theorem (cf. [5], R1 and [4], R). *Let \underline{A} be a connected monounary algebra. Then \underline{A} is retract irreducible in \mathcal{U}_c if and only if \underline{A} is isomorphic to one of the algebras $\underline{\mathbb{N}}$, $\underline{\mathbb{N}}_\infty$, $\underline{\mathbb{N}}_k$ or $\underline{\mathbb{Z}}_{p^k}$ for some $k \in \mathbb{N}$, p prime.*

1.4. Theorem (cf. [6], Thm.). *Let \underline{A} be a connected monounary algebra. Then \underline{A} is retract irreducible in \mathcal{U} if and only if \underline{A} is isomorphic to one of the algebras $\underline{\mathbb{N}}$, $\underline{\mathbb{N}}_\infty$ or $\underline{\mathbb{N}}_k$ for some $k \in \mathbb{N}$.*

In the remaining part of this section we assume that $\underline{A} = (A, f)$ is a connected monounary algebra. We start by proving some auxiliary results.

1.5. Lemma. *Let \underline{A} be representable by means of irreducibles in \mathcal{U}_c or in \mathcal{U} . Then $\{s_f(x) : x \in A\} \subseteq \mathbb{N}_0 \cup \{\infty\}$.*

Proof. The assumption yields that $\underline{A} \in R\left(\prod_{i \in I} \underline{A}_i\right)$, where \underline{A}_i are irreducibles (in \mathcal{U}_c , in \mathcal{U}). There is a retract T of $\prod_{i \in I} \underline{A}_i$ such that $\underline{A} \cong \underline{T}$. By way of contradiction, let $a \in A$ be such that $s_f(a) = \lambda \in \text{Ord-}\mathbb{N}_0$. Let $\iota: \underline{A} \rightarrow \underline{T}$ be an isomorphism, $\iota(a) = b$. Then, for each $i \in I$, $s_f(b(i)) \geq s_f(b) = s(a) = \lambda$ in view of T3). By 1.3 or 1.4, $\{s_f(x) : x \in A_i\} \subseteq \mathbb{N}_0 \cup \{\infty\}$, which implies that $s_f(b(i)) = \infty$ for each $i \in I$, hence $s_f(b) = \infty \neq \lambda = s_f(a)$, which is a contradiction. \square

Now suppose that

$$(*) \quad \{s_f(x) : x \in A\} \subseteq \mathbb{N}_0 \cup \{\infty\}$$

is valid.

1.6. Lemma. *Let A_0 be a retract of \underline{A} , $A_0 \neq A$. Then \underline{A}_0 is a subalgebra of \underline{A} and there exists a representation $\{\underline{A}_i\}_{i \in I \cup \{0\}}$ of \underline{A} such that if $i \in I$, then $\underline{A}_i = (A_i, f_i)$, $A_i \subseteq A$ and*

- (i) $A_i \cap A_0$ is a one-element cycle of A_i ,
- (ii) if $x \in A_i$ is noncyclic, then $f_i(x) = f(x)$,
- (iii) if $x \in A_i$ is cyclic, then $|f_i^{-1}(x)| = 2$.

First, we introduce some notation and then we prove Lemmas 1.6.1–1.6.3; as a consequence we then obtain that 1.6 is valid.

We denote $I = f^{-1}(A_0) - A_0$. If $I = \emptyset$, then $\{\underline{A_0}\}$ is a required representation. Suppose that $I \neq \emptyset$. For $i \in I$ let us put

$$A_i = \bigcup_{l \in \mathbb{N} \cup \{0\}} f^{-l}(i) \cup \{f(i)\}$$

and let the corresponding unary operation f_i on A_i be defined by the formula

$$f_i(x) = \begin{cases} f(x) & \text{if } x \in A_i - \{f(i)\}, \\ f(i) & \text{if } x = f(i). \end{cases}$$

Obviously, (i)–(iii) are valid. We denote

$$\underline{B} = (B, f) = \prod_{i \in I \cup \{0\}} \underline{A_i}.$$

Since A_0 is a retract of \underline{A} , there exists a retraction homomorphism φ of A onto A_0 . Let us define a mapping $\tau: \underline{A} \rightarrow \underline{B}$ as follows:

a) If $a \in A_0$, then $\tau(a) \in B$ is such that

$$(\tau(a))(j) = \begin{cases} a & \text{if } j = 0, \\ f(j) & \text{if } j \in I. \end{cases}$$

b) If $a \in A - A_0$, then there exists a uniquely determined $i \in I$ such that $a \in A_i - A_0$; let us denote by $\tau(a) \in B$ such an element that

$$(\tau(a))(j) = \begin{cases} \varphi(a) & \text{if } j = 0, \\ a & \text{if } j = i, \\ f(j) & \text{if } j \in I - \{i\}. \end{cases}$$

Let $T = \{\tau(a) : a \in A\}$.

1.6.1. Lemma. *The mapping $\tau: A \rightarrow B$ is injective.*

Proof. Assume that $a, b \in A$, $\tau(a) = \tau(b)$. If $a, b \in A_0$, then

$$a = (\tau(a))(0) = (\tau(b))(0) = b.$$

Let $a \notin A_0$, thus $a \in A_i - A_0$ for some $a \in I$. Then

$$(\tau(b))(i) = (\tau(a))(i) = a \in A_i - A_0$$

and the definition of $\tau(b)$ implies that $b = a$. □

1.6.2. Lemma. *The mapping τ is a homomorphism of \underline{A} into \underline{B} .*

Proof. First let $a \in A$ be such that $\{a, f(a)\} \subseteq A_i - A_0$, $i \in I$. Then

$$(\tau(f(a)))(j) = \begin{cases} \varphi(f(a)) & \text{if } j = 0, \\ f(a) & \text{if } j = i, \\ f(j) & \text{otherwise.} \end{cases}$$

Therefore $\tau(f(a)) = f(\tau(a))$.

Now let $a \in A_i - A_0$, $f(a) \in A_0$, $i \in I$. Then $a = i$ and for $j \in I - \{i\}$ we obtain

$$\begin{aligned} f_0(\tau(a)(0)) &= f_0(\varphi(a)) = f(\varphi(a)) = \varphi(f(a)) = f(a) = (\tau(f(a)))(0); \\ f_i(\tau(a)(i)) &= f_i(a) = f(a) = f(i) = (\tau(f(i)))(i) = (\tau(f(a)))(a); \\ f_j(\tau(a)(j)) &= f_j(f(j)) = f(j) = (\tau(f(a)))(j); \end{aligned}$$

i.e., $f(\tau(a)) = \tau(f(a))$.

Finally, if $\{a, f(a)\} \subseteq A_0$, then obviously $\tau(f(a)) = f(\tau(a))$. □

1.6.3. Lemma. *T is a retract of \underline{B} .*

Proof. a) First let us prove the validity of a) of 1.1. Assume that $y \in f^{-1}(T)$, $f(y) = \tau(a)$, $a \in A$. First let $a \in A_0$. We obtain

$$f(y(0)) = (f(y))(0) = (\tau(a))(0) = a.$$

Put $z = \tau(y(0))$. Then $z \in T$ and $f(z) = f(\tau(y(0))) = \tau(f(y(0))) = \tau(a) = f(y)$. Further, by T3) and T2),

$$s_f(y) \leq s_f(y(0)) \leq s_f(\tau(y(0))) = s_f(z).$$

Now suppose that $a \in A_i - A_0$, $i \in I$. Then

$$f(y(i)) = f_i(y(i)) = (f(y))(i) = a.$$

Put $z = \tau(y(i))$. Then $z \in T$ and $f(z) = f(\tau(y(i))) = \tau(f(y(i))) = \tau(a) = f(y)$. Similarly as above,

$$s_f(y) \leq s_{f_i}(y(i)) \leq s_f(\tau(y(i))) = s_f(z).$$

To prove b) of 1.1 let K be a connected component of B with $K \cap T = \emptyset$. There is a projection p_i of K onto A_i for any $i \in I$. Then $p_i \circ \tau: \underline{K} \rightarrow \underline{T}$ is a homomorphism. □

Proof of 1.6. According to 1.6.1–1.6.3, $\underline{A} \in R\left(\prod_{i \in I \cup \{0\}} \underline{A}_i\right)$ and we have constructed the required representation. □

2. \underline{A} WITH A ONE-ELEMENT CYCLE

As above, in 2.1–2.6 we suppose that $\underline{A} = (A, f)$ is a connected monounary algebra such that

$$(*) \quad \{s_f(x) : x \in A\} \subseteq \mathbb{N}_0 \cup \{\infty\}$$

is valid. Further, we will speak shortly about a representation by means of irreducibles and we will mean a representation of \underline{A} by means of connected monounary algebras which are retract irreducible in the class \mathcal{U}_c .

Let us consider the following conditions:

- (c1) \underline{A} possesses a one-element cycle $\{c\}$ and $|f^{-1}(c)| = 2$;
- (c2) (c1) is valid and $s_f(x) = \infty$ for each $x \in A$;
- (c3) (c1) is valid and $s_f(x) \neq \infty$ for each $x \in A - \{c\}$.

2.1. Lemma. *If (c2) holds in \underline{A} , then \underline{A} is representable by means of irreducibles.*

Proof. Assume that (c2) is valid and that \underline{A} is not irreducible. By 1.3, there are $a, b \in A$ with $a \neq b$ such that $f(a) = f(b) \neq c$. Then the condition (C4) from [4] is satisfied and in view of the construction 4.1 and the corresponding lemmas in [4] we obtain that \underline{A} is representable by means of irreducibles (these irreducibles are isomorphic to $\underline{\mathbb{N}}_\infty$). □

2.2. Notation. Let \underline{A} satisfy (c3). There exists a unique element $a \in f^{-1}(c) - \{c\}$. Put

$$d(\underline{A}) = s_f(a) + 1;$$

we will call this positive integer the depth of \underline{A} .

2.3. Lemma. *Let \underline{A} satisfy (c3). Then there exists a representation $\{\underline{A}_j\}_{j \in J}$ of \underline{A} such that, for $j \in J$, \underline{A}_j satisfies (c3) and either \underline{A}_j is irreducible or $d(\underline{A}_j) < d(\underline{A})$.*

Proof. In view of the notation introduced in 2.2, there exist distinct elements $a = a_1, a_2, \dots, a_{d(\underline{A})} \in A$ such that

$$f(a_k) = \begin{cases} a_{k-1} & \text{if } k > 1, \\ c & \text{if } k = 1, \end{cases}$$

$$f^{-1}(a_{d(\underline{A})}) = \emptyset.$$

It is easy to verify that $A_0 = \{c, a_1, a_2, \dots, a_{d(\underline{A})}\}$ is a retract of \underline{A} . If $A_0 = A$, then the assertion holds. Let $A_0 \neq A$. Thus there is a representation of \underline{A} satisfying the

assertion of 1.6. Put $J = I \cup \{0\}$. From 1.3 it follows that \underline{A}_0 is irreducible. Let $i \in I$. By 1.6 (ii), if $x \in A_i$ and x fails to be cyclic, then $s_{f_i}(x) \leq s_f(x)$. Further, let $\{c_i\}$ be a cycle of A_i and $\{a_i\} = f_i^{-1}(c_i) - \{c_i\}$. Then $c_i \in A_0$ by virtue of 1.6 (i). If $c_i = c$, then $c \neq a_i \in f^{-1}(c) = \{c, a\}$, hence $\{a_i, c_i\} = \{a, c\} \subseteq A_i \cap A_0$ and $|A_i \cap A_0| \geq 2$, which is a contradiction to (i) of 1.6. Thus $c_i = a_k$ for some $k \in \{1, \dots, d(\underline{A})\}$. Hence

$$d(\underline{A}_i) = s_{f_i}(a_i) + 1 \leq s_f(a_i) + 1 \leq s_f(c_i) = s_f(a_k) < s_f(a) + 1 = d(\underline{A}).$$

□

2.4. Lemma. *Let \underline{A} satisfy (c3). Then \underline{A} is representable by means of irreducibles.*

Proof. If \underline{A} is irreducible, then \underline{A} is representable. Assume that \underline{A} is not irreducible. In view of 2.3, \underline{A} has a representation in which the factors are either irreducible or have smaller depth than \underline{A} . The factors \underline{A}_i which are not irreducible have representations in which the factors have depths smaller than \underline{A}_i has. After finitely many steps we obtain a representation in which all factors are irreducible. □

2.5. Lemma. *Let \underline{A} be a monounary algebra with a cycle $\{c\}$, $|A| \geq 2$. Then there exists a representation $\{\underline{A}_i\}_{i \in I}$ of \underline{A} such that \underline{A}_i satisfies (c1) for each $i \in I$.*

Proof. The assertion follows from 1.6, if we take $A_0 = \{c\}$. □

2.6. Lemma. *Let (c1) hold in \underline{A} . There exists a representation $\{\underline{A}_i\}_{i \in I}$ of \underline{A} such that if $i \in I$, then \underline{A}_i satisfies either (c2) or (c3).*

Proof. Let $A_0 = \{x \in A : s_f(x) = \infty\}$. If $A_0 = \{c\}$, then \underline{A} satisfies (c3) and $\{\underline{A}\}$ is a one-element representation of \underline{A} . Suppose that $A_0 \neq \{c\}$. It is easy to verify that A_0 is a retract of \underline{A} and we can apply 1.6, i.e., there is a representation $\{\underline{A}_i\}_{i \in I \cup \{0\}}$ satisfying the assertion of 1.6. Then 1.6 (i) and (ii) imply that $s_{f_i}(x) \neq \infty$ for each noncyclic element x of A_i , $i \in I$. Therefore \underline{A}_0 satisfies (c2) and \underline{A}_i , for $i \in I$, satisfies (c3). □

2.7. Proposition. *Suppose that \underline{A} is a connected monounary algebra with a one-element cycle. Then \underline{A} is representable by means of irreducibles in \mathcal{U}_c ; moreover, each factor of the representation contains a one-element cycle.*

Proof. According to 2.5, there exists a representation $\{\underline{A}_i\}_{i \in I}$ of \underline{A} such that \underline{A}_i satisfies (c1) for each $i \in I$. Further, 2.6 yields that for each \underline{A}_i for $i \in I$ there exists a representation $\{\underline{A}_{ij_i}\}_{j_i \in J_i}$ such that \underline{A}_{ij_i} satisfies either (c2) or (c3) for

each $j_i \in J_i$. All factors are then representable by means of irreducibles in view of 2.1 or 2.4, respectively, therefore \underline{A} is representable by means of irreducibles. All factors which were used contain a one-element cycle. \square

3. REPRESENTATION IN \mathcal{U}_c

In 3.1–3.5 we will suppose that \underline{A} is a connected monounary algebra and that (*) is valid.

3.1. Lemma. *If A is a cycle, then \underline{A} is representable by means of irreducibles.*

Proof. Suppose that \underline{A} is an n -element cycle, where the canonical form of n is $p_1^{\alpha_1} \dots p_k^{\alpha_k}$ (p_1, \dots, p_k being distinct primes, $\alpha_1, \dots, \alpha_k \in \mathbb{N}$). For $i \in \{1, \dots, k\}$ let \underline{A}_i be a cycle with $p_i^{\alpha_i}$ elements. Denote $\underline{B} = \prod_{i=1}^k \underline{A}_i$. Then \underline{B} is a cycle with $p_1^{\alpha_1} p_i^{\alpha_i} \dots p_k^{\alpha_k} = n$ elements, i.e., $\underline{B} \cong \underline{A}$. The algebras \underline{A}_i are irreducible for $i \in 1, \dots, k$ in view of 1.3, therefore \underline{A} is representable by means of irreducibles. \square

3.2. Lemma. *The algebra $\underline{\mathbb{Z}}$ is representable by means of cycles.*

Proof. For $i \in \mathbb{N}$ let \underline{A}_i be a 2^i -element cycle. Put $\underline{B} = \prod_{i \in \mathbb{N}} \underline{A}_i$. Then \underline{B} consists of \aleph_0 connected components, each component being isomorphic to $\underline{\mathbb{Z}}$. Obviously, $\underline{\mathbb{Z}} \in R(\underline{B})$, thus $\underline{\mathbb{Z}}$ is representable by the cycles \underline{A}_i , $i \in \mathbb{N}$. \square

3.3. Lemma. *Assume that there is $x \in A$ with $s_f(x) = \infty$. Then \underline{A} is representable by means of irreducibles.*

Proof. The assumption implies that there is a subalgebra \underline{A}_0 of \underline{A} such that either

- 1) $\underline{A}_0 \cong \underline{\mathbb{Z}}$, or
- 2) \underline{A}_0 is a cycle.

It is easy to see that \underline{A}_0 is a retract of \underline{A} . We get by 1.6 that there is a representation $\{\underline{A}_i\}_{i \in I \cup \{0\}}$ such that if $i \in I$, then \underline{A}_i contains a one-element cycle. According to 2.7, each \underline{A}_i , $i \in I$, is representable by means of irreducibles. Further, \underline{A}_0 is representable by means of cycles in view of 3.2, thus it is representable by means of irreducibles in view of 3.1. Therefore \underline{A} is representable by means of irreducibles. \square

3.4. Lemma. Assume that $s_f(x) \neq \infty$ for each $x \in A$ and that \underline{A} is bounded. Then \underline{A} is representable by means of irreducibles, which are not cycles.

Proof. From [7], 1.6 it follows that there is a subalgebra \underline{A}_0 of \underline{A} such that $\mathbb{N} \cong \underline{A}_0$ and that \underline{A}_0 is a retract of \underline{A} . If $A_0 = A$, then \underline{A} is irreducible. Let $A \neq A_0$. Then according to 1.6, \underline{A} is representable by \underline{A}_0 and by nontrivial algebras containing one-element cycles. Then \underline{A} is representable by means of irreducibles in view of 2.7; one of the factors is $\underline{A}_0 \cong \mathbb{N}$ and the other factors contain a one-element cycle and are nontrivial, thus none of the factors is a cycle. \square

3.5. Lemma. Assume that $s_f(x) \neq \infty$ for each $x \in A$ and that \underline{A} is unbounded. Then there is no representation by means of irreducibles for the algebra \underline{A} .

Proof. By way of contradiction, suppose that there is a representation $\{\underline{A}_i\}_{i \in I}$ of \underline{A} by irreducibles. Denote $\underline{B} = \prod_{i \in I} \underline{A}_i$ and assume that T is a retract of \underline{B} with $\underline{T} \cong \underline{A}$. Then $s_f(x) \neq \infty$ for each $x \in T$. If for each $i \in I$ there is $a_i \in A_i$ with $s_f(a_i) = \infty$, then the element $b \in B$ such that $b(i) = a_i$ for each $i \in I$ satisfies the relation $s_f(b) = \infty$; by T2) this element cannot be mapped by any endomorphism of \underline{B} into T , which yields a contradiction. Thus there is $i \in I$ such that $s_f(x) \neq \infty$ for each $x \in A_i$. Since \underline{A}_i is irreducible, $\underline{A}_i \cong \mathbb{N}$.

Further, we have supposed that \underline{A} is unbounded, thus \underline{T} is unbounded and

$$(2) \quad (\forall n \in \mathbb{N})(\exists m \in \mathbb{N})(f^{-(n+m)}(f^m(x)) \neq \emptyset) \text{ is valid for each } x \in T.$$

Then

$$(3) \quad (\forall n \in \mathbb{N})(\exists m \in \mathbb{N})(f^{-(n+m)}(f^m(x(i))) \neq \emptyset),$$

which is a contradiction to the relation $\underline{A}_i \cong \mathbb{N}$. \square

3.6. Theorem. Let \underline{A} be a connected monounary algebra. Then \underline{A} is representable by means of connected monounary algebras which are retract irreducible in the class \mathcal{U}_c if and only if

- 1) $\{s_f(x) : x \in A\} \subseteq \mathbb{N}_0 \cup \{\infty\}$,
- 2) if $\{s_f(x) : x \in A\} \subseteq \mathbb{N}_0$, then A is bounded.

Proof. Suppose that \underline{A} is representable by means of irreducibles in \mathcal{U}_c . By 1.5, 1) is valid. Let $\{s_f(x) : x \in A\} \subseteq \mathbb{N}_0$. If \underline{A} is unbounded, then 3.5 yields a contradiction; therefore the condition 2) is satisfied, too.

Conversely, let 1) and 2) hold. If there is $x \in A$ with $s_f(x) = \infty$, then \underline{A} is representable by means of irreducibles according to 3.3. If $\{s_f(x) : x \in A\} \subseteq \mathbb{N}_0$, then \underline{A} is bounded by 2), and then 3.4 implies that \underline{A} is representable by means of irreducibles in \mathcal{U}_c . \square

4. REPRESENTATION IN \mathcal{U}

The aim of this section is to characterize the connected monounary algebras which are representable by means of connected monounary algebras which are retract irreducible in the class \mathcal{U} .

Let \underline{A} be a connected monounary algebra. It follows from 1.3 and 1.4 that if \underline{A} is representable by means of connected monounary algebras which are retract irreducible in \mathcal{U} , then \underline{A} is representable by means of irreducibles in \mathcal{U}_c .

4.1. Lemma. *Suppose there is a subalgebra \underline{C} of \underline{A} such that either $\underline{C} \cong \underline{\mathbb{Z}}$ or $\underline{C} \cong \underline{\mathbb{Z}}_n$ for some $n \in A$, $n > 1$. Then \underline{A} is not representable by means of connected monounary algebras which are retract irreducible in \mathcal{U} .*

Proof. Assume that $\{A_i\}_{i \in I}$ is a representation by means of connected monounary algebras which are retract irreducible in \mathcal{U} , corresponding to the algebra \underline{A} . By assumption, there is $a \in A$ with $s_f(a) = \infty$. Put $\underline{B} = \prod_{i \in I} A_i$ and let $\iota: A \rightarrow T$ be an isomorphism of \underline{A} onto a retract T of \underline{B} . Then $s_f(\iota(a)) = \infty$, hence T3) yields that $s_f((\iota(a))(i)) = \infty$ for each $i \in I$. Thus if $i \in I$, then there is $a_i \in A_i$ with $s_f(a_i) = \infty$. Since \underline{A}_i is irreducible in \mathcal{U} , 1.4 implies that a_i forms a one-element cycle. This yields that \underline{B} is a connected monounary algebra possessing a one-element cycle. Therefore neither \underline{B} nor the retract T of \underline{B} contains a subalgebra isomorphic to $\underline{\mathbb{Z}}$ ($\underline{\mathbb{Z}}_n$ for $n > 1$, respectively), which contradicts the relation $\underline{T} \cong \underline{A}$. \square

4.2. Theorem. *Let \underline{A} be a connected monounary algebra. Then \underline{A} is representable by means of connected monounary algebras which are retract irreducible in \mathcal{U} if and only if the following conditions are satisfied:*

- (1) $\{s_f(x): x \in A\} \subseteq \mathbb{N}_0 \cup \{\infty\}$,
- (2) if $\{s_f(x): x \in A\} \subseteq \mathbb{N}_0$, then \underline{A} is bounded,
- (3) if $x \in A$, $s_f(x) = \infty$, then $f(x) = x$.

Proof. As was already remarked, if \underline{A} is representable by means of connected monounary algebras which are retract irreducible in \mathcal{U} , then \underline{A} is representable by means of irreducibles in \mathcal{U}_c , thus then (1) and (2) hold. If in this case (3) fails to hold, then 4.1 yields a contradiction.

Conversely, suppose that (1)–(3) are valid. Therefore one of the following possibilities occurs:

- (a) \underline{A} contains a one-element cycle;
- (b) \underline{A} contains no cycle, $\{s_f(x): x \in A\} \subseteq \mathbb{N}_0$, \underline{A} is bounded.

Let (a) hold. By 2.7, \underline{A} is representable by means of irreducibles in \mathcal{U}_c , each factor of the representation containing a one-element cycle, i.e., it is irreducible in \mathcal{U} as well. Thus \underline{A} is representable by means of irreducibles in \mathcal{U} .

Now let (b) be valid. According to 3.4 there exists a representation of \underline{A} by means of irreducibles in \mathcal{U}_c , where none of the factors in the representation is a cycle. Thus this is a representation of \underline{A} by means of irreducibles in \mathcal{U} , which concludes the proof. \square

References

- [1] *D. Duffus, I. Rival*: A structure theory for ordered sets. *Discrete Math.* 35 (1981), 53–118.
- [2] *W. Imrich, S. Klavžar*: Retracts of strong products of graphs. *Discrete Math.* 109 (1992), 147–154.
- [3] *D. Jakubíková-Studenovská*: Systems of unary algebras with common endomorphisms I. *Czechoslovak Math. J.* 29 (1979), 406–420.
- [4] *D. Jakubíková-Studenovská*: Retract irreducibility of connected monounary algebras I. *Czechoslovak Math. J.* 46(121) (1996), 291–308.
- [5] *D. Jakubíková-Studenovská*: Retract irreducibility of connected monounary algebras II. *Czechoslovak Math. J.* 47(122) (1997), 113–126.
- [6] *D. Jakubíková-Studenovská*: Two types of retract irreducibility of connected monounary algebras. *Math. Bohemica* 121 (1996), 143–150.
- [7] *D. Jakubíková-Studenovská*: Antiatomic retract varieties of monounary algebras. *Czechoslovak Math. J.* 48(123) (1998), 793–808.
- [8] *D. Jakubíková-Studenovská*: DR-irreducibility of connected monounary algebras with a cycle. *Czechoslovak Math. J.* 50(125) (2000), 681–698.
- [9] *D. Jakubíková-Studenovská*: DR-irreducibility of connected monounary algebras. *Czechoslovak Math. J.* 50(125) (2000), 705–720.
- [10] *D. Jakubíková-Studenovská*: On a representation of monounary algebras. *Czechoslovak Math. J.* To appear.
- [11] *S. Klavžar*: Two remarks on retracts of graph products. *Discrete Math.* 109 (1992), 155–160.
- [12] *T. Kuczmanov, S. Reich, M. Schmidt, and A. Stachura*: The product retraction property for the c_0 -product of countable many metric spaces. *Math. Japonica* 39 (1994), 73–79.
- [13] *M. Novotný*: Über Abbildungen von Mengen. *Pacif. J. Math.* 13 (1963), 1359–1369.
- [14] *M. Novotný*: Construction of all homomorphisms of unary algebras. *Proceedings of the Seminar on New Directions in Mathematical Education, Brno, 1973.* Dept. Math. Ped. Fac. Univ. J. E. Purkyně, Brno, 1973. (In Czech.)

Author's address: Institute of Mathematics, Faculty of Science, P. J. Šafárik University, Jesenná 5, 041 54 Košice, Slovakia, e-mail: studenovska@science.upjs.sk.