

Dănuț Marcu

A note on the domination number of a graph and its complement

Mathematica Bohemica, Vol. 126 (2001), No. 1, 63–65

Persistent URL: <http://dml.cz/dmlcz/133925>

Terms of use:

© Institute of Mathematics AS CR, 2001

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

A NOTE ON THE DOMINATION NUMBER OF A GRAPH
AND ITS COMPLEMENT

DĂNUȚ MARCU, Bucharest

(Received February 19, 1999)

Abstract. If G is a simple graph of size n without isolated vertices and \overline{G} is its complement, we show that the domination numbers of G and \overline{G} satisfy

$$\gamma(G) + \gamma(\overline{G}) \leq \begin{cases} n - \delta + 2 & \text{if } \gamma(G) > 3, \\ \delta + 3 & \text{if } \gamma(\overline{G}) > 3, \end{cases}$$

where δ is the minimum degree of vertices in G .

Keywords: graphs, domination number, graph's complement

MSC 2000: 05C40

INTRODUCTION

Graphs, considered here, are *finite* and *simple* (without loops or multiple edges), and [1,2] are followed for terminology and notation.

Let $G = (V, E)$ be an *undirected graph* with the set of *vertices* V and the set of edges E . The *complement* \overline{G} of G is the graph with vertex set V , two vertices being adjacent in \overline{G} if and only if they are not adjacent in G .

For any vertex v of G , the *neighbour set* of v is the set of all vertices adjacent to v ; this set is denoted by $N(v)$. A vertex is said to be *isolated* if its neighbour is empty. Suppose that W is a nonempty subset of V . The subgraph of G , whose vertex set is W and whose edge set is the set of those edges of G that have both ends in W , is called the subgraph of G *induced* by W and is denoted by $G[W]$. A set of vertices in a graph is said to be *dominating* if every vertex not in the set is adjacent to one

or more vertices in the set. A *minimal dominating set* is a dominating set such that no proper subset of it is also a dominating set.

The *domination number* $\gamma(G)$ of G is the size of the smallest minimal dominating set.

THE MAIN RESULTS

In the sequel, we will denote $n = |V|$ and $\delta = \min_{v \in V} |N(v)|$.

Theorem 1. *If $G = (V, E)$ is a graph without isolated vertices and $\gamma(G) > 3$, then $\gamma(G) + \gamma(\overline{G}) \leq n - \delta + 2$.*

Proof. Let $v \in V$ be such that $\delta = |N(v)|$ (obviously, since G has no isolated vertices, we have $\delta \geq 1$) and $W = V - (N(v) \cup \{v\})$. If W is empty, then $\gamma(G) = 1$, contradicting the hypothesis. Thus $|W| \geq 1$ and, by the choice of v , it follows that $|N(w)| \geq \delta$ for each $w \in W$.

Consequently, if all vertices of W are isolated in $G[W]$, then $(w, u) \in E$ for every $w \in W$ and $u \in N(v)$, that is, $\{v, u\}$ is a dominating set in G for each $u \in N(v)$. Thus, $\gamma(G) = 2$, contradicting the hypothesis. Let now $Z \subset W$ ($Z \neq W$) be the set of isolated vertices in $G[W]$ (Z can be empty or nonempty), and $Z^* = W - Z$. Let also $D \subseteq Z^*$ be a minimal dominating set in $G[Z^*]$.

If Z is empty, then $D \cup \{v\}$ is a dominating set of G , and we have $\gamma(G) \leq |D \cup \{v\}| = 1 + |D|$. Hence, $|D| \geq \gamma(G) - 1$.

If Z is nonempty, then, since $\delta \leq |N(z)|$ for each $z \in Z$, we have $(z, u) \in E$ for every $z \in Z$ and $u \in N(v)$. Consequently, for each $u \in N(v)$, $D \cup \{v\} \cup \{u\}$ is a dominating set of G and, therefore, we have $\gamma(G) \leq |D \cup \{v\} \cup \{u\}| = 2 + |D|$. Hence, $|D| \geq \gamma(G) - 2$. Thus we always have

$$(1) \quad |D| \geq \gamma(G) - 2.$$

By (1), since $\gamma(G) > 3$, we can choose $B \subseteq D$ such that $|B| = \gamma(G) - 3$.

Let $C \subseteq Z^*$ be the set of vertices in $G[Z^*]$ dominated by B , and $C^* = Z^* - C$. Suppose Z to be empty. If there exists $c \in C$ such that $(c, c^*) \in E$ for each $c^* \in C^*$, then $B \cup \{v\} \cup \{c\}$ is a dominating set in G , that is, $\gamma(G) \leq |B \cup \{v\} \cup \{c\}| = 2 + |B| = \gamma(G) - 1$; a contradiction. Thus for every $c \in C$ there exists $c^* \in C^*$ such that $(c, c^*) \notin E$. If there exists $u \in N(v)$ such that $(u, c^*) \in E$ for each $c^* \in C^*$, then $B \cup \{v\} \cup \{u\}$ is a dominating set in G , that is, $\gamma(G) \leq |B \cup \{v\} \cup \{u\}| = 2 + |B| = \gamma(G) - 1$; a contradiction. Thus for every $u \in N(v)$ there exists $c^* \in C^*$ such that $(u, c^*) \notin E$. On the other hand, by the choice of v , for each $c^* \in C^*$ we have $(v, c^*) \notin E$. Consequently, $C^* = C^* \cup Z$ is a dominating set in \overline{G} .

Suppose Z to be nonempty. By the choice of v , we have $(v, z) \notin E$ for each $z \in Z$. Also, $(z, c) \notin E$ for every $z \in Z$ and $c \in C$. Suppose that there exists $u \in N(v)$ such that $(u, c^*) \in E$ for each $c^* \in C^*$. Since $\delta \leq |N(z)|$ for each $z \in Z$, we have $(t, z) \in E$ for every $z \in Z$ and $t \in N(v)$. Hence $B \cup \{u\} \cup \{v\}$ is a dominating set in G , that is, $\gamma(G) \leq |B \cup \{u\} \cup \{v\}| = 2 + |B| = \gamma(G) - 1$; a contradiction. Thus for every $u \in N(v)$ there exists $c^* \in C^*$ such that $(u, c^*) \notin E$. Consequently, $C^* \cup Z$ is a dominating set in \overline{G} .

So we have

$$(2) \quad \gamma(\overline{G}) \leq |C^* \cup Z| = |C^*| + |Z| = |Z^*| - |C| + |Z| = |W| - |C| = n - \delta - 1 - |C|.$$

However, because $G[Z^*]$ does not contain isolated vertices, it follows that $|B| \leq |C|$ and, by (2), since $|B| = \gamma(G) - 3$, we obtain $\gamma(G) + \gamma(\overline{G}) \leq n - \delta + 2$. \square

Theorem 2. *If $G = (V, E)$ is a graph without isolated vertices and $\gamma(\overline{G}) > 3$, then $\gamma(G) + \gamma(\overline{G}) \leq \delta + 3$.*

Proof. Let $v \in V$ be such that $\delta = |N(v)|$ (obviously, since G has no isolated vertices, we have $\delta \geq 1$). Obviously, $N(v) \cup \{v\}$ is a dominating set in \overline{G} , that is, $\gamma(\overline{G}) \leq |N(v) \cup \{v\}| = 1 + \delta$. Thus $\delta \geq \gamma(\overline{G}) - 1$ and, since $\gamma(\overline{G}) > 3$, we can choose $B \subseteq N(v)$ such that $|B| = \gamma(\overline{G}) - 3$. Let $B^* = N(v) - B$ and $W = V - (N(v) \cup \{v\})$. If W is empty, then the minimum degree of vertices in G is less than δ , contradicting the choice of v . Hence $|W| \geq 1$. Let $w \in W$. We have $|B \cup \{v\} \cup \{w\}| = 2 + |B| = \gamma(\overline{G}) - 1$, that is, $B \cup \{v\} \cup \{w\}$ is not a dominating set in \overline{G} . Consequently, there exists $x \in V$ such that $(x, v) \in E$, $(x, w) \in E$ and $(x, b) \in E$ for each $b \in B$. Obviously, since G does not contain loops, $x \in B^*$. So for every $w \in W$ there exists $b_w^* \in B^*$ such that $(b_w^*, w) \in E$, $(b_w^*, v) \in E$ and $(b_w^*, b) \in E$ for each $b \in B$. Hence B^* is a dominating set in G , that is, $\gamma(G) \leq |B^*| = |N(v)| - |B| = \delta - \gamma(\overline{G}) + 3$. Therefore, $\gamma(G) + \gamma(\overline{G}) \leq \delta + 3$. \square

Corollary. *If G is a graph without isolated vertices such that $\gamma(G) > 3$ and $\gamma(\overline{G}) > 3$, then $\gamma(G) + \gamma(\overline{G}) \leq \lfloor (n + 5)/2 \rfloor$ (we use $\lfloor x \rfloor$ to denote the integer less than or equal to x).*

Proof. It follows from the above theorems. \square

References

- [1] C. Berge: Graphes et Hypergraphes. Dunod, Paris, 1970.
- [2] J. A. Bondy, U. S. R. Murty: Graph Theory with Applications. Macmillan Press, 1976.

Author's address: Dănuț Marcu, Str. Pasului 3, Sect. 2, 70241-Bucharest, Romania.