

Danica Jakubíková-Studenovská

On a cancellation law for monounary algebras

*Mathematica Bohemica*, Vol. 128 (2003), No. 1, 77–90

Persistent URL: <http://dml.cz/dmlcz/133930>

## Terms of use:

© Institute of Mathematics AS CR, 2003

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

## ON A CANCELLATION LAW FOR MONOUNARY ALGEBRAS

DANICA JAKUBÍKOVÁ-STUDENOVSKÁ, Košice

(Received August 24, 2001)

*Abstract.* In this paper we investigate the validity of a cancellation law for some classes of monounary algebras.

*Keywords:* monounary algebra, direct product, connected component, cancellation law

*MSC 2000:* 08A60

## 1. INTRODUCTION

For monounary algebras we apply the standard notation (cf., e.g., [1]).

In this paper we deal with the implication

$$(1) \quad AB \cong AC \Rightarrow B \cong C,$$

where  $A$ ,  $B$  and  $C$  are monounary algebras.

If  $\mathcal{K}$  is a class of monounary algebras such that for each  $A, B, C \in \mathcal{K}$  the implication (1) is valid, then we say that the cancellation law (1) holds in  $\mathcal{K}$ .

For a given monounary algebra  $D$  we denote by  $\mathcal{U}(D)$  the class of all monounary algebras  $A$  such that

- (i) the number of connected components of  $A$  is finite;
- (ii) if  $E$  is a connected component of  $A$ , then  $E$  can be expressed as the direct product of a finite number of subalgebras  $A_1, A_2, \dots, A_n$  of  $D$  such that no  $A_i$  ( $i = 1, 2, \dots, n$ ) is a cycle.

We denote by  $\mathbb{Z} = (\mathbb{Z}, f)$  the monounary algebra such that  $f(x) = x + 1$  for each  $x \in \mathbb{Z}$ .

Let  $n \in \mathbb{N}$ . Then  $D_n$  denotes a connected monounary algebra such that  $D_n = \{a_0, a_1, \dots, a_{n-1}\} \cup \mathbb{N}$ , where  $\{a_0, a_1, \dots, a_{n-1}\}$  is an  $n$ -element cycle and for  $1 \neq k \in \mathbb{N}$  we have  $f(k) = k - 1$ ,  $f(1) = a_0$ .

We prove the following results:

- ( $\alpha$ ) The class  $\mathcal{U}(\mathbb{Z})$  does not satisfy the cancellation law (1).
- ( $\beta$ ) For each  $n \in \mathbb{N}$ , the cancellation law holds in the class  $\mathcal{U}(D_n)$ .

When proving ( $\beta$ ), we apply different methods for the case  $n = 1$  and for the case  $n > 1$ .

The validity of a cancellation law for finite unary algebras was investigated in [7]. In [6], a cancellation law for monounary algebras which are sums of cycles was dealt with.

The cancellation law (1) for finite algebras was studied in [3], [4]; cf. also the monograph [5], Section 5.7. In [2], the implication (1) for partially ordered sets was investigated.

## 2. PRELIMINARIES

In this section we recall some definitions and prove some auxiliary results concerning the class  $\mathcal{U}(D_1)$ .

By a monounary algebra we understand a pair  $(A, f)$ , where  $A$  is a non-empty set and  $f$  is a mapping of  $A$  into  $A$ . If no misunderstanding can occur, then we write  $A$  instead of  $(A, f)$ .

A monounary algebra  $(A, f)$  is said to be connected if for each  $x, y \in A$  there are  $m, n \in \mathbb{N} \cup \{0\}$  such that  $f^m(x) = f^n(y)$ . A maximal connected subalgebra of a monounary algebra  $(A, f)$  is called a connected component of  $(A, f)$ .

Let  $A$  be a monounary algebra. An element  $a \in A$  is cyclic if  $f^n(a) = a$  for some  $n \in \mathbb{N}$ . Let  $B$  be a connected subalgebra of  $A$ . If each element of  $B$  is cyclic, then  $B$  is called a cycle of  $A$ .

Let  $n \in \mathbb{N}$ . For  $i \in \mathbb{Z}$  we denote  $i_n = \{j \in \mathbb{Z} : j \equiv i \pmod{n}\}$ . Next, let  $\mathbb{Z}_n = \{0_n, 1_n, \dots, (n-1)_n\}$  be the set of all integers modulo  $n$ . We define a monounary algebra  $D_n = (D_n, f)$  putting

$$D_n = \mathbb{Z}_n \cup \mathbb{N},$$

$$f(a) = \begin{cases} a + 1_n & \text{if } a \in \mathbb{Z}_n, \\ a - 1 & \text{if } a \in \mathbb{N}, a \neq 1, \\ 0_n & \text{if } a = 1. \end{cases}$$

For  $n = 1$  we write 0 instead of the symbol  $0_n$ , i.e.,  $D_1 = \{0\} \cup \mathbb{N}$ .

Let  $n \in \mathbb{N}$  and let  $X_1, \dots, X_n$  be subalgebras of  $D_1$  having more than one element. Further, let  $\xi$  be an isomorphism of  $X_1 X_2 \dots X_n$  onto a monounary algebra  $A$ . We will omit brackets and write just  $\xi(x_1, \dots, x_n)$  instead of  $\xi((x_1, \dots, x_n))$ . We denote

$$\begin{aligned} X_1^{(0)} &= \{\xi(x_1, 0, \dots, 0) : x_1 \in X_1\}, \\ X_2^{(0)} &= \{\xi(0, x_2, 0, \dots, 0) : x_2 \in X_2\}, \dots, \\ X_n^{(0)} &= \{\xi(0, 0, \dots, x_n) : x_n \in X_n\}. \end{aligned}$$

**2.1. Lemma.** *A is a connected monounary algebra with a one-element cycle  $\{\xi(0, 0, \dots, 0)\}$ . Further,*

$$|f^{-1}(\xi(0, 0, \dots, 0))| = 2^n.$$

*Proof.* Let  $x = \xi(x_1, \dots, x_n) \in A$ ,  $k = \max\{x_1, \dots, x_n\} + 1$ . Then

$$f^k(x) = \xi(f^k(x_1), \dots, f^k(x_n)) = \xi(0, \dots, 0) = f(\xi(0, \dots, 0)),$$

which implies that the element  $\xi(0, \dots, 0)$  forms a one-element cycle of  $A$  and that  $A$  is connected. Next,

$$\begin{aligned} f^{-1}(\xi(0, \dots, 0)) &= \{\xi(y_1, \dots, y_n) : y_i \in \{0, 1\} \text{ for each } i \in \{1, \dots, n\}\}, \\ |f^{-1}(\xi(0, \dots, 0))| &= 2^n. \end{aligned}$$

□

**2.2. Lemma.** *Let  $x \in A$  be such that  $f^{-1}(x) \neq \emptyset$ . Then  $x \in \bigcup_{i=1}^n X_i^{(0)}$  if and only if  $|f^{-1}(x)| \in \{2^n, 2^{n-1}\}$ .*

*Proof.* Suppose that  $x \in \bigcup_{i=1}^n X_i^{(0)}$ . There are  $i \in I$  and  $x_i \in X_i$  with  $x = \xi(0, 0, \dots, x_i, \dots, 0)$ . We have supposed that  $f^{-1}(x) \neq \emptyset$ , thus  $f^{-1}(x_i) \neq \emptyset$ ; if  $x_i = 0$ , then  $f^{-1}(x_i) = \{0, 1\}$  and if  $x_i \neq 0$ , then  $f^{-1}(x_i) = x_i + 1$ . Let  $y \in f^{-1}(x)$ . If  $j \neq i$ , then the  $j$ -th projection of  $\xi^{-1}(y)$  belongs to the set  $\{0, 1\}$ . Hence

- (a) if  $x_i = 0$ , then  $|f^{-1}(x)| = 2^n$  by 2.1,
- (b) if  $x_i \neq 0$ , then

$$\begin{aligned} f^{-1}(x) &= \{\xi(y_1, y_2, \dots, x_i + 1, \dots, y_n) : y_j \in \{0, 1\} \text{ for } j \neq i\}, \\ |f^{-1}(x)| &= 2^{n-1}. \end{aligned}$$

Therefore

$$|f^{-1}(x)| \in \{2^n, 2^{n-1}\}.$$

Conversely, assume that  $x \in A - \bigcup_{i=1}^n X_i^{(0)}$ . Then the number of projections of  $\xi^{-1}(x)$  which are equal to 0 is less than  $n - 1$ ; without loss of generality,  $x = \xi(x_1, \dots, x_k, 0, \dots, 0)$ ,  $\{x_1, \dots, x_k\} \cap \{0\} = \emptyset$ ,  $k > 1$ . We obtain

$$f^{-1}(x) = \{\xi(x_1 + 1, \dots, x_k + 1, y_{k+1}, \dots, y_n) : y_{k+1}, \dots, y_n \in \{0, 1\}\},$$

which implies that  $|f^{-1}(x)| \leq 2^{n-2}$ .  $\square$

Now let  $n, m \in \mathbb{N}$  and let  $X_1, \dots, X_n, Y_1, \dots, Y_m$  be subalgebras of  $D_1$  having more than one element. Further, let  $A$  be a monounary algebra such that  $\xi$  is an isomorphism of  $X_1 X_2 \dots X_n$  onto  $A$  and let  $\eta$  be an isomorphism of  $Y_1 Y_2 \dots Y_m$  onto  $A$ . We suppose that  $X_1^{(0)}, \dots, X_n^{(0)}, Y_1^{(0)}, \dots, Y_m^{(0)}$  have an analogous meaning as above.

### 2.3. Lemma.

(1)  $n = m$ ;

(2)  $\left\{x \in \bigcup_{i=1}^n X_i^{(0)} : f^{-1}(x) \neq \emptyset\right\} = \left\{y \in \bigcup_{i=1}^n Y_i^{(0)} : f^{-1}(y) \neq \emptyset\right\}$ ;

(3) *there is a permutation  $\varphi$  of the set  $\{1, 2, \dots, n\}$  such that  $X_k \cong X_k^{(0)} \cong Y_{\varphi(k)}^{(0)} \cong Y_{\varphi(k)}$  for each  $k \in \{1, \dots, n\}$ .*

*Proof.* In view of 2.1 we obtain that  $\{\xi(0, \dots, 0)\} = \{\eta(0, \dots, 0)\}$  is a cycle of  $A$  and

$$2^n = |f^{-1}(\xi(0, \dots, 0))| = |f^{-1}(\eta(0, \dots, 0))| = 2^m,$$

therefore  $n = m$ .

The assertion (2) follows from 2.2.

The set  $\bigcup_{i=1}^n X_i^{(0)}$  is a subalgebra of  $A$ . Further, for  $i \in \{1, \dots, n\}$ ,  $X_i^{(0)} \cong X_i$  is a subalgebra of  $D_1$  and  $X_i^{(0)} \cap X_j^{(0)}$  is a one-element cycle of  $A$  whenever  $i, j \in \{1, \dots, n\}$ ,  $i \neq j$ . (Analogously for  $Y_1, \dots, Y_n$ .) Notice that if  $x \in X_j^{(0)}$ ,  $j \in \{1, \dots, n\}$  and  $f^{-1}(x) = \emptyset$ , then the element  $f(x) = z$  has the property that  $f^{-1}(z) \neq \emptyset$ ,  $z \in \left(\bigcup_{i=1}^n X_i^{(0)}\right) \cap \left(\bigcup_{i=1}^n Y_i^{(0)}\right)$ .

Let  $k \in \{1, \dots, n\}$ .

(a) Suppose that  $X_k \cong D_1$ . Then  $X_k^{(0)} \cong D_1$  and for each  $x \in X_k^{(0)}$  we have  $f^{-1}(x) \neq \emptyset$ . According to (2),

$$(4) \quad X_k^{(0)} \subseteq \left\{y \in \bigcup_{i=1}^n Y_i^{(0)} : f^{-1}(y) \neq \emptyset\right\}.$$

Take  $t \in X_k^{(0)}$  such that  $t$  does not belong to a cycle but  $f(t)$  does;  $t$  is uniquely determined. Further, (4) implies that there is  $j \in \{1, \dots, n\}$  with  $t \in Y_j$ . By the above consideration, the set  $Y_i^{(0)} \cap Y_j^{(0)}$  is a one-element cycle of  $A$  for each  $i \neq j$ , therefore we obtain in view of (4) that  $Y_j^{(0)} = X_k^{(0)}$ ; let us denote  $j = \varphi(k)$ .

(b) Now let  $X_k \neq D_1$ . There is exactly one  $x \in X_k^{(0)}$  such that  $f^{-1}(x) = \emptyset$ . Let  $z = f(x)$ . According to (2) the subalgebra  $\{f^l(z) : l \in \mathbb{N} \cup \{0\}\}$  of  $X_k^{(0)}$  is a subalgebra of  $\left\{y \in \bigcup_{i=1}^n Y_i^{(0)} : f^{-1}(y) \neq \emptyset\right\}$  and analogously as above, there is exactly one  $j \in \{1, \dots, n\}$  such that

$$(5) \quad \{f^l(z) : l \in \mathbb{N} \cup \{0\}\} = \{y \in Y_j^{(0)} : f^{-1}(y) \neq \emptyset\}.$$

We have  $X_k^{(0)} = \{x\} \cup \{f^l(z) : l \in \mathbb{N} \cup \{0\}\}$ . Similarly,  $Y_j^{(0)}$  consists of the elements of the right set in (5) and of one element  $q$  with the property  $f^{-1}(q) = \emptyset$ . Hence  $X_k^{(0)} \cong Y_j^{(0)}$ . We denote  $j = \varphi(k)$ .

The mapping  $\varphi$  is a permutation and  $X_k^{(0)} \cong Y_{\varphi(k)}^{(0)}$  for each  $k \in \{1, \dots, n\}$ , i.e., (3) is valid.  $\square$

**2.4. Proposition.** *If  $A$  is isomorphic to a direct product of subalgebras of  $D_1$  such that these subalgebras are not cycles, then the decomposition of  $A$  into such a direct product is unique up to isomorphism.*

*Proof.* This is a corollary of 2.3.  $\square$

### 3. CANCELLATION LAW IN $\mathcal{U}(D_1)$

For investigating the properties of  $\mathcal{U}(D_1)$ , in this section we deal with the system  $\mathbb{Z}[y_1, \dots, y_n]$  of polynomials with unknowns  $y_1, \dots, y_n$  over the integrity domain of integers  $\mathbb{Z}$ ; it is known that  $\mathbb{Z}[y_1, \dots, y_n]$  is an integrity domain as well.

A similar consideration has been used in [2] for investigating the cancellation law for partially ordered sets, where generalized polynomials over  $\mathbb{Z}$  have been taken into account.

Let  $A, B, C$  be monounary algebras belonging to the class  $\mathcal{U}(D_1)$ . Next, let  $\{Y_1, \dots, Y_n\}$  be a system of monounary algebras such that

- (a) if  $i, j \in \{1, \dots, n\}, i \neq j$ , then  $Y_i \not\cong Y_j$ ,
- (b) if  $E \in \{A, B, C\}$ ,  $F$  is a connected component of  $E$  and  $F = X_1 X_2 \dots X_k$ , where  $X_1, \dots, X_k$  are subalgebras of  $D_1$  which are not cycles, then for each  $j \in \{1, \dots, k\}$  there is  $l \in \{1, \dots, n\}$  such that  $X_i \cong Y_l$ ,
- (c) if  $l \in \{1, \dots, n\}$ , then  $|Y_l| > 1$ .

Let us remark that it follows from the definition of  $\mathcal{U}(D_1)$  that a system  $\{Y_1, \dots, Y_n\}$  with the required properties exists.

**3.1. Notation.** If  $X$  is a monounary algebra,  $k \in \mathbb{N}$ , then we denote by  $X^0$  a one-element monounary algebra and

$$X^k = XX \dots X \text{ (} k\text{-times)}.$$

If  $k \in \mathbb{N}$  and  $X_1, \dots, X_k$  are mutually disjoint monounary algebras, then  $\sum_{i=1}^k X_i$  is a disjoint union of the given algebras. Further, if  $X_1 \cong X_2 \cong \dots \cong X_k$ , then we write also  $kX_1$  instead of  $\sum_{i=1}^k X_i$ ; thus  $kX_2$  is an algebra consisting of  $k$  copies of  $X_1$ .

We denote  $0X = \emptyset$  for each monounary algebra  $X$ .

**3.2. Lemma.** *Let  $t_1, \dots, t_n, s_1, \dots, s_n \in \mathbb{N} \cup \{0\}$ . Then  $Y_1^{t_1} Y_2^{t_2} \dots Y_n^{t_n} \cong Y_1^{s_1} Y_2^{s_2} \dots Y_n^{s_n}$  if and only if  $t_1 = s_1, \dots, t_n = s_n$ .*

**Proof.** Since the condition (a) is satisfied, we obtain the assertion by virtue of 2.4. □

**3.3. Corollary.** *Let  $E \in \{A, B, C\}$ , let  $F$  be a connected component of  $E$ . Then  $F$  can be expressed in the form  $F \cong Y_1^{t_1} \dots Y_n^{t_n}$ , where  $t_1, \dots, t_n \in \mathbb{N} \cup \{0\}$ ; further,  $t_1, \dots, t_n$  are uniquely determined.*

**3.4. Notation.** Let  $f(y_1, \dots, y_n) \in \mathbb{Z}[y_1, \dots, y_n]$  be a polynomial with non-negative coefficients. Then we can write it in the form

$$f(y_1, \dots, y_n) = \sum_{i=1}^m p_i y_1^{t_{i1}} y_2^{t_{i2}} \dots y_n^{t_{in}}$$

such that

- (i)  $p_i \geq 0$  for each  $i \in \{1, \dots, m\}$ ,
- (ii) if  $j, k$  are distinct elements of the set  $\{1, \dots, m\}$ , then  $y_1^{t_{j1}} y_2^{t_{j2}} \dots y_n^{t_{jn}} \neq y_1^{t_{k1}} y_2^{t_{k2}} \dots y_n^{t_{kn}}$ ; we will say that  $f(y_1, \dots, y_n)$  is written in a normal form. By  $f(Y_1, \dots, Y_n)$  we denote the monounary algebra

$$\sum_{i=1}^m p_i Y_1^{t_{i1}} Y_2^{t_{i2}} \dots Y_n^{t_{in}}.$$

**3.5. Lemma.** Let  $f(y_1, \dots, y_n), g(y_1, \dots, y_n) \in \mathbb{Z}[y_1, \dots, y_n]$  be polynomials with non-negative coefficients which are written in a normal form. Then  $f(Y_1, \dots, Y_n) = g(Y_1, \dots, Y_n)$  if and only if  $f(y_1, \dots, y_n) = g(y_1, \dots, y_n)$ .

*Proof.* If  $f(y_1, \dots, y_n) = g(y_1, \dots, y_n)$ , then 3.4 implies that  $f(Y_1, \dots, Y_n) = g(Y_1, \dots, Y_n)$ . The converse implication follows from 3.2 in view of the fact that the polynomials are written in a normal form.  $\square$

**3.6. Corollary.** There are uniquely determined polynomials  $f_A(y_1, \dots, y_n), f_B(y_1, \dots, y_n), f_C(y_1, \dots, y_n)$  with non-negative coefficients such that

$$\begin{aligned} A &\cong f_A(Y_1, \dots, Y_n), \\ B &\cong f_B(Y_1, \dots, Y_n), \\ C &\cong f_C(Y_1, \dots, Y_n). \end{aligned}$$

*Proof.* This is a consequence of the definition of the system  $\{Y_1, \dots, Y_n\}$  and of 3.5.  $\square$

**3.7. Corollary.** Let  $(f_A \cdot f_B)(y_1, \dots, y_n) = f_A(y_1, \dots, y_n) \cdot f_B(y_1, \dots, y_n)$ . Then

$$AB \cong (f_A \cdot f_B)(Y_1, \dots, Y_n).$$

*Proof.* By 3.6,  $AB \cong f_A(Y_1, \dots, Y_n)f_B(Y_1, \dots, Y_n)$ , thus we get the assertion in view of 3.4.  $\square$

**3.8. Theorem.** Let  $A, B, C \in \mathcal{U}(D_1)$ ,  $AB \cong AC$ . Then  $B \cong C$ .

*Proof.* According to 3.7 we obtain  $AB \cong (f_A \cdot f_B)(Y_1, \dots, Y_n)$ , and similarly,  $AC \cong (f_A \cdot f_C)(Y_1, \dots, Y_n)$ . Thus

$$(f_A \cdot f_B)(Y_1, \dots, Y_n) \cong (f_A \cdot f_C)(Y_1, \dots, Y_n).$$

According to 3.5,

$$\begin{aligned} (f_A \cdot f_B)(y_1, \dots, y_n) &= (f_A \cdot f_C)(y_1, \dots, y_n), \\ f_A(y_1, \dots, y_n) \cdot f_B(y_1, \dots, y_n) &= f_A(y_1, \dots, y_n) \cdot f_C(y_1, \dots, y_n). \end{aligned}$$

The polynomial  $f_A(y_1, \dots, y_n)$  is a non-zero polynomial, thus we can apply the cancellation law in the integrity domain  $\mathbb{Z}[y_1, \dots, y_n]$ , which implies

$$f_B(y_1, \dots, y_n) = f_C(y_1, \dots, y_n).$$

Again by 3.5,  $f_B(Y_1, \dots, Y_n) = f_C(Y_1, \dots, Y_n)$ , thus  $B \cong C$ .  $\square$



Now we will give two examples showing that if some of the conditions (i), (ii) in the definition of the class  $\mathcal{U}(D)$  fails to hold, then the cancellation law need not be valid in general.

3.9. **Example.** Let  $E$  be an arbitrary subalgebra of  $D_1$ ,  $|E| > 1$ . Put  $A = \aleph_0 E$ ,  $B = E$ ,  $C = 2E$ . Then

$$\begin{aligned} AB &= (\aleph_0 E)E = \aleph_0(EE), \\ AC &= (\aleph_0 E)(2E) \cong \aleph_0(EE). \end{aligned}$$

Hence  $AB \cong AC$ , but  $B \not\cong C$ . Notice that here each connected component  $F$  of  $A$ ,  $B$ ,  $C$  is a subalgebra of  $D_1$  and  $|F| > 1$ , i.e.,  $A$ ,  $B$  and  $C$  fulfil the condition (ii).

3.10. **Example.** Let  $E$  be as in 3.9. Take  $A = E^{\aleph_0}$ ,  $B = E$ ,  $C = E^2$ . Then

$$AB \cong E^{\aleph_0} \cong AC, \quad B \not\cong C.$$

Here  $A$ ,  $B$ ,  $C$  have finitely many connected components, each connected component is a direct product of subalgebras of  $D_1$ , but there are infinitely many factors in the product in  $A$ .

#### 4. THE CLASS $\mathcal{U}(\mathbb{Z})$

Let  $\mathbb{N} = (\mathbb{N}, f)$  be a monounary algebra such that  $f(x) = x + 1$  for each  $x \in \mathbb{N}$ . We will show that the cancellation law (1) in  $\mathcal{U}(\mathbb{N})$  is not valid in general.

**4.1. Lemma.**  $\mathbb{N}\mathbb{N} \cong \aleph_0\mathbb{N}$ .

**Proof.** Let  $E$  be a connected component of  $\mathbb{N}\mathbb{N}$ ,  $u = (u_1, u_2) \in E$ . Without loss of generality, suppose that  $u_1 \leq u_2$ . Let  $a = (1, u_2 - u_1 + 1)$ . Then

$$f^{u_1-1}(a) = (1 + (u_1 - 1), u_2 - u_1 + 1 + (u_1 - 1)) = (u_1, u_2) = u,$$

thus  $a \in E$ . Further,  $f^{-1}(a) = \emptyset$ . If  $i, j \in \mathbb{N} \cup \{0\}$ ,  $f^i(a) = f^j(a)$ , then

$$(1 + i, u_2 - u_2 - u_1 + 1 + i) = (1 + j, u_2 - u_1 + 1 + j),$$

which implies that  $i = j$ . We will show that

$$E = \{f^i(a) : i \in \mathbb{N} \cup \{0\}\}.$$

Let  $(w_1, w_2) = w \in E$ . Then there are  $m, n \in \mathbb{N} \cup \{0\}$  such that  $f^m(a) = f^n(w)$ . Denote  $a_2 = w_2 - w_1 + 1$ . We obtain

$$\begin{aligned}(1 + m, a_2 + m) &= (w_1 + n, w_2 + n), \\ 1 + m &= w_1 + n, a_2 + m = w_2 + n, \\ w_1 &= 1 + m - n, w_2 = a_2 + m - n.\end{aligned}$$

Since  $w_1 \geq 1$ , we have  $m - n \geq 0$  and  $f^{m-n}(a) = w$ . Therefore  $E \subseteq \{f^i(a) : i \in \mathbb{N} \cup \{0\}\}$ . The converse inclusion is obvious. Hence each connected component of  $\mathbb{NN}$  is isomorphic to  $\mathbb{N}$ .

Further, if  $i, j \in \mathbb{N}$ ,  $i \neq j$ , then  $(1, i)$  and  $(1, j)$  do not belong to the same connected component. We have  $|\mathbb{NN}| = \aleph_0$ , thus  $\mathbb{NN}$  consists of  $\aleph_0$  connected components which are all isomorphic to  $\mathbb{N}$ .  $\square$

**4.2. Lemma.** *A monounary algebra  $E$  belongs to  $\mathcal{U}(\mathbb{N})$  if and only if  $E \cong k\mathbb{N}$ ,  $k \in \mathbb{N}$ .*

*Proof.* Let  $E \in \mathcal{U}(\mathbb{N})$ . If  $F$  is a connected component of  $E$ , then  $F$  is a direct product of finitely many subalgebras of  $\mathbb{N}$ . Since each subalgebra of  $\mathbb{N}$  is isomorphic to  $\mathbb{N}$  and a product of at least two algebras isomorphic to  $\mathbb{N}$  is non-connected by 4.1, we obtain that each connected component  $F$  of  $E$  is isomorphic to  $\mathbb{N}$ . Next,  $E$  consists of finitely many connected components, which implies that  $E \cong k\mathbb{N}$ ,  $k \in \mathbb{N}$ .

The relation  $\{k\mathbb{N} : k \in \mathbb{N}\} \subseteq \mathcal{U}(\mathbb{N})$  is obvious.  $\square$

**4.3. Lemma.** *Let  $A, B \in \mathcal{U}(\mathbb{N})$ . Then  $AB \cong \aleph_0\mathbb{N}$ .*

*Proof.* By 4.2 there are  $k, m \in \mathbb{N}$  with  $A \cong k\mathbb{N}$ ,  $B \cong m\mathbb{N}$ . According to 4.1 we obtain

$$AB \cong (k\mathbb{N})(m\mathbb{N}) = (km)(\mathbb{NN}) \cong \aleph_0\mathbb{N}.$$

$\square$

From 4.3 we infer that the cancellation law (1) does not hold in  $\mathcal{U}(\mathbb{N})$  in general. Further, as a corollary we obtain

**4.4. Theorem.**

- (a) *For each  $A \in \mathcal{U}(\mathbb{N})$  there are  $B, C \in \mathcal{U}(\mathbb{N})$  with  $B \not\cong C$ ,  $AB \cong AC$ .*
- (b) *For each  $B, C \in \mathcal{U}(\mathbb{N})$  there is  $A \in \mathcal{U}(\mathbb{N})$  such that  $AB \cong AC$ .*

**4.5. Corollary.** *The cancellation law (1) in  $\mathcal{U}(\mathbb{Z})$  does not hold in general.*

*Proof.* This is a consequence of the fact that the class  $\mathcal{U}(\mathbb{N})$  is a subclass of  $\mathcal{U}(\mathbb{Z})$  and that the cancellation law (1) does not hold in  $\mathcal{U}(\mathbb{N})$  in general.  $\square$

## 5. CANCELLATION LAW IN $\mathcal{U}(D_n)$

Let  $n \in \mathbb{N}, n > 1$ . According to the notation of Section 2 we have  $D_n = \mathbb{Z}_n \cup \mathbb{N}$ , where  $i_n \in \mathbb{Z}_n$  is the set of all integers  $k$  with  $k \equiv i \pmod{n}$ .

**5.1. Lemma.** *If  $X, Y$  are subalgebras of  $D_n$ , then  $XY$  is non-connected.*

*Proof.* Let  $a = (0_n, 1_n), b \in (1_n, 0_n) \in XY$ . By way of contradiction, suppose that  $XY$  is connected. Then there are  $k, m \in \mathbb{N} \cup \{0\}$  such that  $f^k(a) = f^m(b)$ . We obtain

$$((0+k)_n, (1+k)_n) = ((1+m)_n, (0+m)_n),$$

i.e.,  $k \equiv m+1, 1+k \equiv m \pmod{n}$ . This implies that  $n/2$  and since  $n > 1$ , we have  $n = 2$ . Take  $c = (0_2, 0_2)$ . There exist  $p, q \in \mathbb{N} \cup \{0\}$  such that  $f^p(a) = f^q(c)$ , thus

$$(p_2, (1+p)_2) = (q_2, q_2),$$

i.e.,  $p \equiv q, 1+p \equiv q \pmod{2}$ , which is a contradiction. □

**5.2. Lemma.** *A monounary algebra  $E$  belongs to  $\mathcal{U}(D_n)$  if and only if  $E$  consists of finitely many connected components and each connected component  $F$  of  $E$  is a subalgebra of  $D_n, |F| > n$ .*

*Proof.* Let  $E \in \mathcal{U}(D_n)$ . Then it has finitely many connected components. By the definition of  $\mathcal{U}(D_n)$ , no connected component  $F$  of  $E$  is a cycle, thus  $|F| > n$ . The remaining part of the proof is analogous to 4.2 provided we apply 5.1. □

In 5.3.1–5.5.3 let  $X, Y$  be subalgebras of  $D_n$  such that  $|X| > n, |Y| > n$ . There are  $k, m \in \mathbb{N} \cup \{\aleph_0\}$  with

$$X = \mathbb{Z}_n \cup \{i \in \mathbb{N}: i \leq k\},$$

$$Y = \mathbb{Z}_n \cup \{i \in \mathbb{N}: i \leq m\}.$$

Let  $E = XY$ . The following two lemmas are easy to verify by a routine calculation.

**5.3.1. Lemma.**

- (a) *Let  $v = (v_1, v_2) \in E$ . Then  $v$  belongs to a cycle of  $E$  if and only if  $v_1, v_2 \in \mathbb{Z}_n$ .*
- (b) *Each connected component of  $E$  contains a cycle with  $n$  elements.*

**5.3.2. Lemma.**

- (a)  $f^{-1}((0_n, 0_n)) = \{((n-1)_n, (n-1)_n), (1, (n-1)_n), ((n-1)_n, 1),;$   
 $((n-1)_n, 1), (1, 1)\}$

- (b) if  $0_n \neq j_n \in \mathbb{Z}_n$ , then  $f^{-1}((j_n, 0_n)) = \{((j-1)_n, (n-1)_n), ((j-1)_n, 1)\}$ ,  
 $f^{-1}((0_n, j_n)) = \{((n-1)_n, (j-1)_n), (1, (j-1)_n)\}$ ;  
(c) if  $0_n \neq j_n \in \mathbb{Z}_n$ ,  $0_n \neq l_n \in \mathbb{Z}_n$ , then  $f^{-1}((j_n, l_n)) = \{((j-1)_n, (l-1)_n)\}$ .

**5.3.3. Corollary.** *Let  $v$  be a cyclic element of  $E$ . Then  $v = (0_n, 0_n)$  if and only if  $|f^{-1}(v)| = 4$ .*

**5.4. Lemma.** *Let  $F$  be the connected component of  $E$  containing the element  $(0_n, 0_n)$ .*

- (a)  $|\{v \in F : v \text{ is cyclic, } |f^{-1}(v)| > 1\}| = 1$ ;  
(b) if  $F_1$  is a connected component of  $E$  such that  $F_1 \neq F$ , then  $|\{v \in F_1 : v \text{ is cyclic, } |f^{-1}(v)| > 1\}| > 1$ .

*Proof.* Let  $v$  be a cyclic element of  $F$ . Then  $v = (i_n, i_n)$ ,  $i_n \in \mathbb{Z}_n$ . If  $i_n = 0_n$ , then 5.3.3 implies that  $|f^{-1}(v)| = 4$ . If  $i_n \neq 0_n$ , then 5.3.2(c) yields that  $|f^{-1}(v)| = 1$ . Hence (a) is valid.

Now let  $F_1$  be a connected component of  $E$  such that  $F_1 \neq F$ . Then there is  $j \in \{1, 2, \dots, n-1\}$  such that  $(0_n, j_n) \in F_1$ . Denote  $v = (0_n, j_n)$ ,  $w = f^{n-j}(v)$ . Thus  $w$  is a cyclic element of  $F_1$ ,

$$w = ((n-j)_n, (j+n-j)_n) = ((-j)_n, 0_n).$$

According to 5.3.2(b),  $|f^{-1}(v)| = 2 = |f^{-1}(w)|$ , which implies that (b) holds.  $\square$

Denote  $u = (0_n, 0_n)$ ,  $u^{(1)} = (1, 1)$ ,  $u^{(2)} = (1, (n-1)_n)$ ,  $u^{(3)} = ((n-1)_n, 1)$ .

**5.5.1. Lemma.**

- (a) If  $k = m = \aleph_0$ , then  $f^{-1}(u^{(\alpha)}) \neq \emptyset$  for each  $\alpha \in \{1, 2, 3\}$ ,  $i \in \mathbb{N}$ .  
(b) If  $k < m = \aleph_0$ , then  $f^{-1}(u^{(3)}) \neq \emptyset$  for each  $i \in \mathbb{N}$  and  $f^{-k}(u^{(1)}) = \emptyset = f^{-(k-1)}(u^{(1)})$ ,  $f^{-k}(u^{(2)}) = \emptyset \neq f^{-(k-1)}(u^{(2)})$ .  
(c) If  $k \leq m < \aleph_0$ , then  $f^{-m}(u^{(3)}) = \emptyset \neq f^{-(m-1)}(u^{(3)})$ ,  $f^{-k}(u^{(1)}) = \emptyset \neq f^{-(k-1)}(u^{(1)})$ ,  $f^{-k}(u^{(2)}) = \emptyset \neq f^{-(k-1)}(u^{(2)})$ .

*Proof.*

- (a) Let  $k = m = \aleph_0$ ,  $i \in \mathbb{N}$ . Then

$$\begin{aligned} f^i((i+1, i+1)) &= (i+1-i, i+1-i) = (1, 1) = u^{(1)}, \\ f^i((i+1, (n-1-i)_n)) &= (i+1-i, (n-1-i+i)_n) = (1, (n-1)_n) = u^{(2)}, \\ f^i(((n-1-i)_n, i+1)) &= ((n-1-i+i)_n, i+1-i) = u^{(3)}. \end{aligned}$$

- (b) Let  $k < m = \aleph_0$ ,  $i \in \mathbb{N}$ . Similarly as above,  $f^i((n-1-i)_n, i+1) = u^{(3)}$ .  
Further,

$$f^{k-1}((k, (-k)_n)) = (k - (k-1), (-k + k - 1)_n) = (1, (n-1)_n) = u^{(2)},$$

$$f^{k-1}((k, k)) = (k - (k-1), k - (k-1)) = (1, 1) = u^{(1)}.$$

Suppose that  $f^{-k}(u^{(1)}) \neq \emptyset$  (the case for  $u^{(2)}$  is analogous). Then there is  $(t_1, t_2) \in E$  with

$$u^{(1)} = (1, 1) = f^k((t_1, t_2)) = (f^k(t_1), f^k(t_2)).$$

This implies that in  $X$  the set  $f^{-k}(1)$  is non-empty, which is a contradiction. Therefore (b) is valid.

- (c) The proof of this assertion is similar to that of (b). □

In view of 5.3.2(a), the set  $f^{-1}(u) = f^{-1}((0_n, 0_n))$  consists of a cyclic element  $((n-1)_n, (n-1)_n)$  and of three non-cyclic elements; let  $w^{(1)}$ ,  $w^{(2)}$ ,  $w^{(3)}$  be these elements.

### 5.5.2. Lemma.

- (a) If  $k = m = \aleph_0$ , then  $f^{-i}(w^{(\alpha)}) \neq \emptyset$  for each  $\alpha \in \{1, 2, 3\}$ ,  $i \in \mathbb{N}$ .  
(b) If  $k < m = \aleph_0$ , then there is  $\alpha \in \{1, 2, 3\}$  such that  $f^{-i}(w^{(\alpha)}) \neq \emptyset$  for each  $i \in \mathbb{N}$  and if  $\alpha \neq \beta \in \{1, 2, 3\}$ , then  $f^{-k}(w^{(\beta)}) = \emptyset \neq f^{-(k-1)}(w^{(\beta)})$ .  
(c) If  $k \leq m < \aleph_0$ , then there is  $\alpha \in \{1, 2, 3\}$  such that  $f^{-m}(w^{(\alpha)}) = \emptyset \neq f^{-(m-1)}(w^{(\alpha)})$  and if  $\alpha \neq \beta \in \{1, 2, 3\}$ , then  $f^{-k}(w^{(\beta)}) = \emptyset \neq f^{-(k-1)}(w^{(\beta)})$ .

### 5.5.3. Corollary.

- (a) Let  $f^{-i}(w^{(\alpha)}) \neq \emptyset$  for each  $\alpha \in \{1, 2, 3\}$ ,  $i \in \mathbb{N}$ . Then  $k = m = \aleph_0$ .  
(b) Let the assumption of (a) be not valid and suppose that there is  $\alpha \in \{1, 2, 3\}$  such that  $f^{-i}(w^{(\alpha)}) \neq \emptyset$  for each  $i \in \mathbb{N}$ . Then there is  $j \in \mathbb{N}$  such that if  $\alpha \neq \beta \in \{1, 2, 3\}$ , then  $f^{-j}(w^{(\beta)}) = \emptyset \neq f^{-(j-1)}(w^{(\beta)})$ . Further, this yields that  $\{k, m\} = \{j, \aleph_0\}$ .  
(c) Let neither the assumption of (a) nor the assumption of (b) be valid. There are  $j, l \in \mathbb{N}$  and  $\alpha \in \{1, 2, 3\}$  such that  $f^{-j}(w^{(\alpha)}) = \emptyset \neq f^{-(j-1)}(w^{(\alpha)})$  and if  $\alpha \neq \beta \in \{1, 2, 3\}$ , then  $f^{-l}(w^{(\beta)}) = \emptyset \neq f^{-(l-1)}(w^{(\beta)})$ . Then  $\{k, m\} = \{j, l\}$ .

**5.6. Lemma.** Suppose that  $X, Y, X', Y'$  are subalgebras of  $D_n$  which are not cycles. If  $XY \cong X'Y'$ , then either  $X \cong X'$ ,  $Y \cong Y'$  or  $X \cong Y'$ ,  $Y \cong X'$ .

*Proof.* Let  $k, m, E, u, F$  be as above and assume that  $k', m', E', u', F'$  have an analogous meaning in the product  $X'Y'$ . There is an isomorphism  $\xi: XY \rightarrow X'Y'$ . By 5.3.1,  $\xi$  maps cyclic elements into cyclic elements and by 5.3.3,  $\xi(u) = u'$ . Next, 5.4 implies that  $\xi(F) = F'$ . It follows from 5.5.3 that

$$\{k, m\} = \{k', m'\}.$$

If  $k = k', m = m'$ , then  $X \cong X', Y \cong Y'$ . If  $k = m', m = k'$ , then  $X \cong Y', Y \cong X'$ .  $\square$

**5.7. Theorem.** *Let  $A, B, C \in \mathcal{U}(D_n)$ ,  $n \in \mathbb{N}$ ,  $n > 1$ . Then  $AB \cong AC$  implies  $B \cong C$ .*

*Proof.* It follows from 5.2 that  $A, B, C$  are sums of finitely many subalgebras of  $D_n$ . Let  $\{Y_1, \dots, Y_n\}$  be a system of monounary algebras such that

- (1)  $Y_i \not\cong Y_j$  for  $i, j \in \{1, \dots, n\}$ ,  $i \neq j$ ,
- (2) if  $F$  is a connected component of  $A, B$  or  $C$ , then  $F \cong Y_i$  for some  $i \in \{1, \dots, n\}$ .

Then there are non-negative integers  $\alpha_i, \beta_i, \gamma_i$  ( $i \in \{1, \dots, n\}$ ) such that

$$A \cong \sum_{i=1}^n \alpha_i Y_i, \quad B \cong \sum_{i=1}^n \beta_i Y_i, \quad C \cong \sum_{i=1}^n \gamma_i Y_i.$$

Suppose that  $AB \cong AC$ , i.e.,

$$\sum_{i,j} (\alpha_i \beta_j) (Y_i Y_j) \cong \sum_{i,j} (\alpha_i \gamma_j) (Y_i Y_j).$$

Since (1) is valid, we obtain by virtue of 5.6 that  $\beta_j = \gamma_j$  for each  $j \in \{1, \dots, n\}$ . Therefore  $B \cong C$ .  $\square$

### *References*

- [1] *B. Jónsson*: Topics in Universal Algebra. Springer, Berlin, 1972.
- [2] *J. Jakubík, L. Lihová*: On the cancellation law for disconnected partially ordered sets. Math. Bohem. Submitted.
- [3] *L. Lovász*: Operations with structures. Acta Math. Acad. Sci. Hungar. 18 (1967), 321–328.
- [4] *L. Lovász*: On the cancellation law among finite relational structures. Period. Math. Hungar. 1 (1971), 145–156.
- [5] *R. McKenzie, G. McNulty, W. Taylor*: Algebras, Lattices, Varieties. Vol. I, Wadsworth, Belmont, 1987.
- [6] *J. Novotný*: On the characterization of a certain class of monounary algebras. Math. Slovaca 40 (1990), 123–126.

- [7] *M. Ploščica, M. Zelina*: Cancellation among finite unary algebras. *Discrete Mathematics* 159 (1996), 191–198.

*Author's address*: *D. Jakubíková-Studenovská*, Department of Geometry and Algebra, P. J. Šafárik University, Jesenná 5, 041 54 Košice, Slovak Republic, e-mail: **studenovska@duro.science.upjs.sk**.