

Diane M. Donovan; James G. Lefevre; Thomas A. McCourt; Nicholas J. Cavenagh

Distinct equilateral triangle dissections of convex regions

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 53 (2012), No. 2, 189--210

Persistent URL: <http://dml.cz/dmlcz/142884>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

## Distinct equilateral triangle dissections of convex regions

DIANE M. DONOVAN, JAMES G. LEFEVRE,  
THOMAS A. MCCOURT, NICHOLAS J. CAVENAGH

*Abstract.* We define a proper triangulation to be a dissection of an integer sided equilateral triangle into smaller, integer sided equilateral triangles such that no point is the vertex of more than three of the smaller triangles. In this paper we establish necessary and sufficient conditions for a proper triangulation of a convex region to exist. Moreover we establish precisely when at least two such equilateral triangle dissections exist.

We also provide necessary and sufficient conditions for some convex regions with up to four sides to have either one, or at least two, proper triangulations when an internal triangle is specified.

*Keywords:* equilateral triangle dissection, latin trade

*Classification:* 05B45

### 1. Introduction

The dissection of an integer sided equilateral triangle into smaller, integer sided equilateral triangles is a classic problem considered by Tutte [12]. He showed various properties of such a dissection, including the fact that some of the smaller triangles must have equal sides.

If we apply an extra restriction to such a dissection, namely that no point is the vertex of more than three of the smaller triangles, then the dissection gives rise to a *latin trade* within the addition table for the integers modulo  $n$  ([3]). We call such a dissection a proper triangulation. (It was Drápal, in [3], who first observed the connection between latin trades and proper triangulations, and as a consequence of this in some papers (see [2] and [11]) proper triangulations are also known as Drápal Triangulations.) More details about this connection and latin trades may be found in [1], [3], [4], [6] and [7].

This application of triangle dissections to latin trades is our key motivation. In particular, the results in this paper are applied to classify flaws in cryptographic applications of latin squares [2]. However, the results have some geometric interest in their own right.

It is conjectured that there exists a constant  $c$  such that for each integer  $n$ , there exists a non-trivial proper triangulation of an integer sided equilateral triangle containing at most  $c \log p$  triangles, where  $p$  is the least prime that divides  $n$  ([3]). The results in this paper may provide insights into this question. A further

possible application is the enumeration of proper triangulations (see [5]). Other, laterally related result on triangulations include [8], [9] and [10].

In Section 2 we introduce necessary terminology. In Section 3 we establish precisely when a convex regions has at least one or at least two proper triangulations (Theorem 3.6). In Section 4 we consider the same question when an internal triangle is specified; our results are restricted to convex region with at most four sides which are not rectangles.

## 2. Proper triangulations

For ease of notation we consider the equivalent problem of dissections of right-angled isosceles triangles into smaller such objects, where each triangle has hypotenuse of gradient  $-1$ . To see this equivalence, consider such a dissection with the large triangle lying in the first quadrant with its right angle at the origin. Then the linear transformation  $T(x) = xA$  given by

$$A = \begin{bmatrix} 1 & 0 \\ 0.5 & \sqrt{3}/2 \end{bmatrix}$$

shows the equivalence to an equilateral triangle dissection.

Let  $k, i, x_i, y_i \in \mathbb{Z}$  and  $0 \leq i \leq k - 1$ . Let  $R = (x_0, y_0), (x_1, y_1), \dots, (x_i, y_i), \dots, (x_{k-1}, y_{k-1})$  be a sequence of points which satisfies the following condition: for all  $0 \leq i \leq k - 1$ ,

$$x_i = x_{i+1 \pmod k} \text{ OR } y_i = y_{i+1 \pmod k} \text{ OR } x_i + y_i = x_{i+1 \pmod k} + y_{i+1 \pmod k}.$$

Then we say that  $R$  is a *region* in the plane  $\mathbb{R}^2$ . The *reduced form*  $R'$  of  $R$  is formed by successively deleting any points  $(x_i, y_i)$  from  $R$  whenever  $(x_{i-1}, y_{i-1}), (x_i, y_i)$  and  $(x_{i+1}, y_{i+1})$  are collinear.

If the straight line segments between  $(u_i, v_i) \in R'$  and  $(u_{i+1 \pmod l}, v_{i+1 \pmod l}) \in R'$ ,  $0 \leq i \leq l - 1 = |R'| - 1$ , form the boundary of a convex polygon (where  $R'$  is the reduced form of  $R$ ), then  $R$  is called a *convex region*. Furthermore, the region  $R$  is denoted by  $R = (x_0, y_0) \rightarrow (x_1, y_1) \rightarrow \dots \rightarrow (x_i, y_i) \rightarrow \dots \rightarrow (x_{k-1}, y_{k-1})$ , and if  $1 < |R|$ , we refer to the elements of the reduced form of  $R$  as the *corners* of  $R$ .

If the reduced form of  $R$  has precisely three corners, then  $R$  is said to be a *triangle*. Let  $0 \leq x$ , denote the region

$$FT_x^{(z_1, z_2)} = (z_1, z_2) \rightarrow (z_1 + x, z_2) \rightarrow (z_1, z_2 + x) \text{ as a forward triangle and}$$


$$BT_x^{(z_1, z_2)} = (z_1, z_2) \rightarrow (z_1 - x, z_2) \rightarrow (z_1, z_2 - x) \text{ as a backward triangle.}$$

Let  $R$  be the union of regions  $R_1, R_2, \dots, R_t$ ; that is,  $R = \bigcup_{1 \leq i \leq t} R_i$ . If for each  $1 \leq i < j \leq t$ , the regions  $R_i$  and  $R_j$  intersect in at most their respective boundaries, then  $\{R_i \mid 1 \leq i \leq t\}$  is called a *tessellation* of  $R$  and each  $R_i$  is a *subregion* of  $R$ .

If each of the subregions  $R_i$  is a triangle,  $R$  is said to have a *triangulation*, namely  $\{R_i \mid 1 \leq i \leq t\}$ , furthermore each subregion,  $R_i$ , is referred to as a *subtriangle* of  $R$ . If, in addition, each element  $(a,b) \in R$  is the corner of at most three distinct subtriangles,  $\{R_i \mid 1 \leq i \leq t\}$  is called a *proper triangulation* of the region  $R$ . It is this property which makes the problem of finding proper triangulations of a specified region non-trivial.

**Example 2.1.** In Figure 1 we provide an example of a region,  $R$ , that has a triangulation but no possible proper triangulation and a region  $S$  that has a proper triangulation.

FIGURE 1. A triangulation and a proper triangulation



Consider the following group of matrices, isomorphic to the Dihedral group  $D_6$ :

$$G = \left\langle \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} -1 & 1 \\ -1 & 1 \end{bmatrix} \right\rangle.$$

Let  $\lambda \in G$ ,  $(p, q) \in \mathbb{R}^2$  and  $S \subset \mathbb{R}^2$ . In this paper the set  $\{(m, n)\lambda + (p, q) \mid (m, n) \in S\}$  is denoted by  $S\lambda + (p, q)$ .

If there exists some  $(i, j) \in \mathbb{Z}^2$  and some  $\lambda \in G$  such that  $R_2 = R_1\lambda + (i, j)$ , then  $R_1$  and  $R_2$  are said to be *equivalent*. Observe that the property of possessing a proper triangulation is invariant under this equivalence, even though the gradients of lines may change. We frequently make use of this observation.

Recall that, for a proper triangulation, the condition that each vertex of a triangle is the vertex of at most three triangles must be satisfied. Suppose that  $\{R_i \mid 1 \leq i \leq p\}$  is a tessellation of a region  $R$  and that each subregion  $R_i$  has a proper triangulation  $Q_i$ , where  $1 \leq i \leq p$ . Then the set of triangles

$$\bigcup_{1 \leq i \leq p} Q_i$$


does not necessarily form a proper triangulation of  $R$ .

**Example 2.2.** In Figure 2 we provide an example of a tessellation of a region  $R$  for which each region has a proper triangulation and the union of the subtriangles does not yield a proper triangulation of the region  $R$ .


To avoid this problem, whenever two distinct regions  $R_i$  and  $R_j$  in the tessellation of  $R$  both have a triangulation containing more than one triangle, then we ensure that their boundaries do not share a line segment of non zero length.

FIGURE 2. Failing to construct a proper triangulation


Original tessellation of  $R$



Drápal Triangulations of each subregion



Triangulation of  $R$



**Example 2.3.** In Figure 3 we provide an example of a tessellation of a region  $R$  for which each region has a proper triangulation and the union of the subtriangles yields a proper triangulation of the region  $R$ .

### 3. Proper triangulations of convex regions

For  $\alpha, \beta \in \mathbb{Z}$ , let:

$$Z_0 = (0, 0),$$

$$Z_1 = (0, 0) \rightarrow (1, 0) \rightarrow (1, \alpha) \rightarrow (0, \alpha) \text{ where } 0 < \alpha,$$

$$Z_2 = (1, 0) \rightarrow (\alpha, 0) \rightarrow (\alpha, 1) \rightarrow (0, 1) \text{ where } 0 < \alpha,$$

$$Z_3 = (\alpha, 0) \rightarrow (\alpha, \beta) \rightarrow (0, \beta) \rightarrow (0, \beta - 1) \rightarrow (\alpha - 1, \beta - 1) \rightarrow (\alpha - 1, 0) \text{ where } 1 < \alpha, \beta,$$

$$Z_4 = (2, 0) \rightarrow (2, 2) \rightarrow (0, 2) \rightarrow (0, 1) \rightarrow (1, 0),$$

$$Z_5 = (3, 0) \rightarrow (3, 1) \rightarrow (1, 3) \rightarrow (0, 3) \rightarrow (0, 1) \rightarrow (1, 0) \text{ and}$$


$$Z_6 = (2, 0) \rightarrow (2, 1) \rightarrow (1, 2) \rightarrow (0, 2) \rightarrow (0, 1) \rightarrow (1, 0).$$

Let  $\mathcal{Z}$  be the set of all regions equivalent to any  $Z_i$ , where  $0 \leq i \leq 6$  (see the Appendix for an illustration of these regions).


By inspection, the regions equivalent to  $Z_i$  where  $0 \leq i \leq 5$  have a unique proper triangulation, while  $Z_6$  has no proper triangulation. The aim of this

FIGURE 3. Constructing a proper triangulation


Original tessellation of  $R$



Drápal Triangulations of each subregion



Drápal Triangulation of  $R$



section is to show that any convex region not belonging to  $\mathcal{Z}$  has at least two proper triangulations (Theorem 3.6).

We begin by investigating when it is possible for a region with three or four sides to have at least two distinct proper triangulations.

**Lemma 3.1.** *Let  $R$  be a region with three or four corners (sides). Thus  $R$  is equivalent to*

$$\begin{aligned}
 R_1 &= (\delta, 0) \rightarrow (0, \delta) \rightarrow (0, 0), \text{ or} \\
 R_2 &= (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (0, \alpha) \rightarrow (\gamma, 0),
 \end{aligned}$$

where  $0 < \delta$ , and either  $\gamma = 0$  and  $0 < \alpha \leq \beta$ , or  $\gamma = \alpha$  and  $0 < \alpha < \beta$ . Then  $R$  has a proper triangulation. Further, if  $1 < \delta$ , there exists a second distinct proper triangulation of  $R_1$  and, unless  $R_2$  is equivalent to  $Z_1$  or  $Z_2$ , there exists a second distinct proper triangulation of  $R_2$ .

PROOF: Since  $R_1$  is a triangle, a proper triangulation trivially exists. If  $\delta = 1$ , then, by inspection,  $R_1 = FT_1^{(0,0)}$  has precisely one proper triangulation. However

if  $1 < \delta$ , then

$$\{FT_{\delta-1}^{(1,0)}\} \cup \{FT_1^{(0,0)}\} \cup \bigcup_{1 \leq i \leq \delta-1} \{FT_1^{(0,i)}, BT_1^{(1,i)}\}$$

is a second distinct proper triangulation of  $R_1$ .

There are two cases to consider for  $R_2$ : Case A,  $\gamma = 0$  and Case B,  $\gamma = \alpha$ .

**Case A:**  $\gamma = 0$ .

Consider the tessellation  $\{FT_{\alpha}^{(0,0)}, BT_{\alpha}^{(\alpha,\alpha)}\}$   $R_3 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (\alpha, \alpha) \rightarrow (\alpha, 0)$  of  $R_2$ . If  $\alpha = \beta$ , then  $R_3$  is empty and we are done. Otherwise,  $R_3$  is a rectangular region with area strictly less than  $\alpha\beta$ . Thus, by recursion,  $R_1$  has a proper triangulation.

If  $\alpha = 1$ , then  $R$  is equivalent to  $Z_1$  and by inspection it has precisely one proper triangulation. When  $1 < \alpha$ , the second distinct proper triangulation is obtained by applying the argument given for  $R_1$  to the triangle  $FT_{\alpha}^{(0,0)}$ .

**Case B:**  $\gamma = \alpha$ .

Let  $1 < \alpha$ . Consider the tessellation  $\{BT_{\alpha}^{(\alpha,\alpha)}, S = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (\alpha, \alpha) \rightarrow (\alpha, 0)\}$  of  $R_2$ . If  $1 < \beta - \alpha$ , the argument presented in Case A implies  $S$  has two distinct proper triangulations. If  $\beta - \alpha = 1$ , then the above gives one proper triangulation. Consider the proper triangulation  $\{BT_{\alpha}^{(\beta,\alpha)}\} \cup \bigcup_{1 \leq i \leq \alpha} \{FT_1^{(i,\alpha-i)}, BT_1^{(i,\alpha-i+1)}\}$  of  $R_2$ . This yields a second distinct proper triangulation of  $R_2$ .

If  $\alpha = 1$ , then  $R_2$  is equivalent to  $Z_2$  and by inspection it has precisely one proper triangulation. □

An L-region will be defined to be a region equivalent to

$$(\delta, 0) \rightarrow (\delta, \beta) \rightarrow (0, \beta) \rightarrow (0, \alpha) \rightarrow (\gamma, \alpha) \rightarrow (\gamma, 0),$$

where  $0 < \alpha < \beta$  and  $0 < \gamma < \delta$ .

In order to obtain a similar result to Lemma 3.1 for convex regions with five sides we first prove the following result detailing when an L-region has at least two distinct proper triangulations.

**Lemma 3.2.** *Let  $0 < \alpha < \beta$  and  $0 < \gamma < \delta$ . The L-region  $L_1 = (\delta, 0) \rightarrow (\delta, \beta) \rightarrow (0, \beta) \rightarrow (0, \alpha) \rightarrow (\gamma, \alpha) \rightarrow (\gamma, 0)$  has a proper triangulation, and a second distinct proper triangulation when  $L_1$  is not equivalent to  $Z_3$ .*

PROOF: Several cases are considered which, together with the associated conditions, are summarized in the following table.

|                                      |                                       |                                       |                                   |
|--------------------------------------|---------------------------------------|---------------------------------------|-----------------------------------|
| <b>Case A</b>                        | <b>Case B</b>                         | <b>Case C</b>                         | <b>Case D</b>                     |
| $\alpha + \gamma \geq \delta, \beta$ | $\delta \leq \alpha + \gamma < \beta$ | $\beta \leq \alpha + \gamma < \delta$ | $\alpha + \gamma < \beta, \delta$ |

**Case A:**  $\alpha + \gamma \geq \delta, \beta$ .

Consider the tessellation  $R = \{BT_{\beta+\delta-\alpha-\gamma}^{(\delta,\beta)}, R_1 = (\delta, 0) \rightarrow (\delta, \alpha + \gamma - \delta) \rightarrow (\gamma, \alpha) \rightarrow (\gamma, 0), R_2 = (\gamma, \alpha) \rightarrow (\alpha + \gamma - \beta, \beta) \rightarrow (0, \beta) \rightarrow (0, \alpha)\}$  of  $L_1$ . By Lemma 3.1,  $L_1$  has a proper triangulation.

Provided  $R_1$  or  $R_2$  are not both equivalent to  $Z_2$  then Lemma 3.1 implies there exists a second distinct proper triangulation of  $L_1$ . When both  $R_1$  and  $R_2$  are equivalent to  $Z_2$ , then  $\beta - \alpha = \delta - \gamma = 1$ ,  $L_1$  is equivalent to  $Z_3$  and by inspection it has precisely one proper triangulation.

**Case B:**  $\delta \leq \alpha + \gamma < \beta$ .

Consider the tessellation  $\{BT_{\delta}^{(\delta,\alpha+\gamma)}, FT_{\gamma}^{(0,\alpha)}, R_1 = (\delta, 0) \rightarrow (\delta, \alpha + \gamma - \delta) \rightarrow (\gamma, \alpha) \rightarrow (\gamma, 0), R_2 = (\delta, \alpha + \gamma) \rightarrow (\delta, \beta) \rightarrow (0, \beta) \rightarrow (0, \alpha + \gamma)\}$  of  $L_1$ . As  $\alpha + \gamma < \beta$ ,  $R_2$  is not equivalent to  $Z_0$ . By Lemma 3.1,  $L_1$  has a proper triangulation. If at least one of  $R_1$  and  $R_2$  is equivalent to neither  $Z_1$  nor  $Z_2$ , then  $L_1$  has a second distinct proper triangulation.

If  $1 < \gamma$ , then, by Lemma 3.1,  $FT_{\gamma}^{(0,\alpha)}$  (and hence  $L_1$ ) has a second distinct proper triangulation.

If  $R_1$  and  $R_2$  are each equivalent to either  $Z_1$  or  $Z_2$  and  $\gamma = 1$ , then  $\delta - \gamma = \gamma = \beta - \alpha - \gamma = 1$ , hence,  $\delta = 2$ . Consider the tessellation  $\{BT_2^{(2,\alpha+2)}, FT_2^{(0,\alpha)}, R_1 = (2, 0) \rightarrow (2, \alpha) \rightarrow (1, \alpha) \rightarrow (1, 0)\}$  of  $L_1$ . By Lemma 3.1 there exists a second distinct proper triangulation of  $L_1$ .

**Case C:**  $\beta \leq \alpha + \gamma < \delta$ .

Via the transformation  $L_1 \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$  this region is equivalent to the region in Case B.

**Case D:**  $\alpha + \gamma < \beta, \delta$ .

Consider the tessellation  $\{BT_{\alpha+\gamma}^{(\alpha+\gamma,\alpha+\gamma)}, FT_{\gamma}^{(0,\alpha)}, FT_{\alpha}^{(\gamma,0)}, R_1 = (\delta, 0) \rightarrow (\delta, \beta) \rightarrow (0, \beta) \rightarrow (0, \alpha + \gamma) \rightarrow (\alpha + \gamma, \alpha + \gamma) \rightarrow (\alpha + \gamma, 0)\}$  of  $L_1$ . Observe that  $R_1$  is an  $L$ -region, so is equivalent to one of the regions given in Case A, B, C or (recursively) D. For Cases B and C, there exists at least two distinct proper triangulations so we are done. Otherwise we have the following subcases:

**Subcase D.1:** Suppose that  $R_1$  is equivalent to a region given in Case A. If either  $\alpha \neq 1, \gamma \neq 1, \beta - \alpha - \gamma \neq 1$  or  $\delta - \alpha - \gamma \neq 1$ , then, by Lemma 3.1, there exist two distinct proper triangulations of  $L_1$ .

Otherwise,  $\alpha = \gamma = \beta - \alpha - \gamma = \delta - \alpha - \gamma = 1$ . Then  $\alpha = \gamma = 1$  and  $\beta = \delta = 3$ . Consider the triangulation  $\{BT_3^{(3,3)}, BT_1^{(2,1)}, FT_2^{(0,1)}, FT_1^{(1,0)}, FT_1^{(2,0)}\}$  of  $L_1$ . This is a second distinct (to the above) proper triangulation of  $L_1$ .

**Subcase D.2:**

Otherwise  $R_1$  is equivalent to a region given in Case D. Note that  $R_1$  has area strictly less than  $L_1$ , so by recursion there exists a proper triangulation of  $R_1$ . Moreover, the tessellation of  $R_2$  contains the triangle  $FT_{\alpha+\gamma}^{(0,\alpha+\gamma)}$ . But  $\alpha + \gamma \geq 2$ , so by Lemma 3.1 there exists a second distinct proper triangulation.  $\square$

We will now make use of Lemmas 3.1 and 3.2 to determine when there exists precisely one and when there exists at least two distinct proper triangulations of convex regions with five sides.



**Lemma 3.3.** *Let  $R$  be a convex region with five corners (sides). Then  $R$  has a proper triangulation. Moreover, whenever  $R$  is not equivalent to  $Z_4$  then  $R$  has a second distinct proper triangulation.*

PROOF: Under the appropriate transformation, we may assume without loss of generality that  $R = (\beta, 0) \rightarrow (\beta, \gamma) \rightarrow (0, \gamma) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$ , where  $0 < \alpha < \beta \leq \gamma$ .

Consider the tessellation  $\{BT_\alpha^{(\alpha, \alpha)}, R_1 = (\beta, 0) \rightarrow (\beta, \gamma) \rightarrow (0, \gamma) \rightarrow (0, \alpha) \rightarrow (\alpha, \alpha) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemma 3.2 the region  $R$  has a proper triangulation.

Provided  $R_1$  is not equivalent to  $Z_3$  ( $\gamma - \alpha \neq 1$  or  $\beta - \alpha \neq 1$ ), Lemma 3.1 and 3.2 imply a second triangulation of  $R_1$ , and so  $R$  has a second distinct proper triangulation.

Suppose that  $R_1$  is equivalent to  $Z_3$ . Then  $\gamma - \alpha = 1$  and  $\gamma = \beta$ . If in addition  $\alpha = 1$ , then  $\beta = \gamma = 2$ , and hence  $R = Z_4$ ; otherwise  $2 < \beta = \gamma$  and  $\{BT_\gamma^{(\beta, \gamma)}\} \cup_{1 \leq i \leq \beta-1} \{FT_1^{(i-1, \gamma-i)}, BT_1^{(i, \gamma-i)}\} \cup FT_1^{(\beta-1, 0)}$  is a second distinct proper triangulation of  $R$ .  $\square$


We will now prove another technical lemma which we will use to establish when a convex region with six sides has a proper triangulation and when it has at least two distinct proper triangulations.

Let  $R$  be a region equivalent to

$$(\beta, 0) \rightarrow (\beta, \delta + \gamma - \beta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$$

with  $0 \leq \alpha < \beta, \delta; 0 < \gamma < \beta, \delta + 1$ ; and  $0 \leq \delta + \gamma - \beta$  (this region is illustrated in Figure 4).

FIGURE 4. Illustration of the region  $R$ .



It will be shown that  $R$  possesses at least two distinct proper triangulations, except when  $R$  is equivalent to any of the following (see the Appendix for illustrations):

$$\begin{aligned}
 X_1 &= (\beta, 0) \rightarrow (\beta, \beta - 1) \rightarrow (\beta - 1, \beta) \rightarrow (\beta - 1, 1) \rightarrow (0, \beta) \rightarrow (0, \beta - 1) \rightarrow (\beta - 1, 0); \\
 X_2 &= (3, 0) \rightarrow (1, 2) \rightarrow (1, 1) \rightarrow (0, 2) \rightarrow (0, 1) \rightarrow (1, 0); \\
 X_3 &= (3, 0) \rightarrow (3, 1) \rightarrow (1, 3) \rightarrow (1, 2) \rightarrow (0, 3) \rightarrow (0, 1) \rightarrow (1, 0); \\
 X_4 &= (3, 0) \rightarrow (3, 2) \rightarrow (2, 3) \rightarrow (2, 1) \rightarrow (0, 3) \rightarrow (0, 1) \rightarrow (1, 0); \\
 X_5 &= (\beta, 0) \rightarrow (\beta, \beta - 2) \rightarrow (\beta - 1, \beta - 1) \rightarrow (\beta - 1, 0) \rightarrow (0, \beta - 1) \rightarrow (0, \beta - 2) \rightarrow (\beta - 2, 0).
 \end{aligned}$$

Let  $\mathcal{X}$  be the set of all regions equivalent to any  $X_i$ , where  $1 \leq i \leq 5$ .

By inspection the regions equivalent to  $X_i$ , where  $2 \leq i \leq 5$ , have a unique proper triangulation. Furthermore, by inspection, the region  $X_1$  has no proper triangulation.

**Lemma 3.4.** *Let  $0 \leq \alpha < \beta, \delta; 0 < \gamma < \beta, \delta + 1$ ; and  $0 \leq \delta + \gamma - \beta$ . The region  $R = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$  has a proper triangulation if and only if  $R \neq X_1$  and a second distinct proper triangulation if and only if  $R$  is not equivalent to any  $X_i$ , where  $1 \leq i \leq 5$ .*

PROOF: Consider the transformation  $R\lambda + (\delta + \gamma, 0)$ , where  $\lambda = \begin{bmatrix} -1 & 0 \\ -1 & 1 \end{bmatrix} \in G$ . This transformation replaces  $\beta$  with  $\delta + \gamma - \alpha$  and  $\alpha$  with  $\delta + \gamma - \beta$ . Hence, without loss of generality, we may assume that  $\delta + \gamma - \beta \leq \alpha$ .

Several cases are considered which, together with the associated conditions, are summarized in the following table.

|   |   |                                       |                   |
|---|---|---------------------------------------|-------------------|
| <b>Case A</b>                             | <b>Case B</b>                             | <b>Case C</b>                         | <b>Case D</b>     |
| $\gamma < \delta; \gamma + 2 \leq \beta;$ | $\gamma < \delta; \gamma + 2 \leq \beta;$ | $\gamma < \delta; \beta = \gamma + 1$ | $\delta = \gamma$ |
| $\alpha < \gamma$                         | $\alpha \geq \gamma$                      |                                       |                   |

For Case B several additional subcases are considered which are summarized in the following table.

|           |  |                                    |  |
|-----------|--|------------------------------------|--|
| <b>B1</b> | $\alpha - \gamma \geq \gamma + \delta - \beta$ |                                    |  |
| <b>B2</b> | $\alpha - \gamma < \gamma + \delta - \beta$    |                                    | <b>B2.1</b> $\beta - \gamma > \delta - \alpha$ |
|           |  | $\beta - \gamma = \delta - \alpha$ | <b>B2.2.1</b> $\alpha \geq 2$                  |
|           |  |                                    | <b>B2.2.2</b> $\alpha \leq 1$                  |

**Case A:**  $\gamma < \delta, \gamma + 2 \leq \beta$  and  $\alpha < \gamma$ .

The conditions for this case together with the assumption  $\delta + \gamma - \beta \leq \alpha$  imply  $\delta < \beta$ .

Consider the tessellation  $\{FT_{\delta-\alpha}^{(\alpha,0)}, R_1 = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - \gamma) \rightarrow (\delta, 0), R_2 = (\alpha, 0) \rightarrow (\alpha, \delta - \alpha) \rightarrow (0, \delta) \rightarrow (0, \alpha)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a proper triangulation of  $R$ .

Unless either  $R_1$  is equivalent to one of  $Z_2$  or  $Z_4$  or  $R_2$  is equivalent to one of  $Z_0$  or  $Z_1$ , Lemmas 3.1 and 3.3 imply the existence of a second proper triangulation of  $R$ .

Suppose that  $R_1$  is equivalent to  $Z_2$ .

Then  $\beta - \delta = 1$  and  $\delta + \gamma - \beta = 0$ , so  $\gamma = 1$ .

Thus  $\alpha = 0$  and  $R_2$  is equivalent to  $Z_0$ . In this case Lemma 3.1 applied to  $R_3$  in the tessellation  $\{FT_\delta^{(1,0)}, R_3 = (1, 0) \rightarrow (1, \delta - 1) \rightarrow (0, \delta) \rightarrow (0, 0)\}$  verifies the existence of a second proper triangulation of  $R$ .

Otherwise suppose  $R_1$  is equivalent to  $Z_4$ . Then  $\gamma = 2, \beta - \gamma = 2, \beta - \delta = 1$  and  $\delta + \gamma - \beta = 1$ . Hence  $\beta = 4, \delta = 3$  and since  $\delta + \gamma - \beta \leq \alpha$  and  $\alpha < \gamma$  it follows that  $\alpha = 1$ . In which case  $\{FT_2^{(0,1)}, FT_2^{(2,0)}, FT_1^{(1,0)}, FT_1^{(2,2)}, FT_1^{(3,1)}, BT_1^{(1,1)}, BT_1^{(2,1)}, BT_1^{(3,2)}, BT_1^{(4,1)}\}$  is a second distinct proper triangulation of  $R$ .

**Subcase B1:**  $\gamma < \delta, \gamma + 2 \leq \beta, \gamma \leq \alpha$  and  $\alpha - \gamma \geq \gamma + \delta - \beta$ .

From the conditions for this subcase and the assumption that  $0 < \gamma$  it follows that  $\gamma + \delta - \beta < \alpha$  or equivalently  $\gamma + \delta - \alpha < \beta$ .

Consider the tessellation  $\{FT_{\delta+\gamma-\alpha}^{(\gamma, \alpha-\gamma)}, R_1 = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (2\gamma + \delta - \alpha, \alpha - \gamma) \rightarrow (\gamma, \alpha - \gamma) \rightarrow (\alpha, 0), R_2 = (\gamma, \alpha - \gamma) \rightarrow (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a proper triangulation of  $R$ .

By Lemmas 3.1 and 3.3 either there exists a second proper triangulation of  $R$  or  $R_1$  is equivalent to an element of  $\{Z_0, Z_1, Z_2, Z_4\}$  and  $R_2$  is equivalent to  $Z_1$ . Henceforth, assume the latter.

Suppose that  $\delta = 2$ . Then  $\gamma = \alpha = 1$  and so  $R$  is equivalent to  $X_2$  and by inspection there does not exist a second distinct proper triangulation of  $R$ . Thus,  $\delta > 2$ .

Since  $R_2$  is equivalent to  $Z_1$ , either  $\gamma = 1$  or  $\delta - \alpha = 1$ .

First suppose  $\gamma = 1$  and  $\delta - \alpha = 1$ . The fact that  $\delta > 2$  implies  $\alpha > 1$ . In this case consider the tessellation  $\{FT_1^{(0, \delta-1)}, FT_1^{(1, \delta-1)}, BT_2^{(2, \delta-1)}, R_3 = (\beta, 0) \rightarrow (\beta, \delta + \gamma - \beta) \rightarrow (2, \delta - 1) \rightarrow (2, \alpha - 2) \rightarrow (\alpha, 0)\}$ .

Secondly, suppose  $\gamma = 1$  and  $\delta - \alpha > 1$  and so  $\delta \geq \alpha + 2$ . Furthermore  $\delta - \alpha > 1$  implies that  $\gamma + \delta - \alpha > 2$ , so  $R_1$  is not equivalent to  $Z_4$ ; thus  $\alpha \leq 2$ . If  $\alpha = 1$ , then  $R_1$  is equivalent to  $Z_0$ , and the conditions  $0 \leq \delta + \gamma - \beta \leq \alpha - \gamma$  imply  $\delta + \gamma - \beta = 0$ . Here we take the tessellation  $\{FT_{\delta-1}^{(0,1)}, FT_1^{(1, \delta-1)}, R_4 = (\beta, 0) \rightarrow (2, \delta - 1) \rightarrow (1, \delta - 1) \rightarrow (\delta - 1, 1) \rightarrow (0, 1) \rightarrow (1, 0)\}$  of  $R$ . Note  $R_4$  is equivalent to  $Z_3$  and hence there exists a proper triangulation of  $R_4$ . Otherwise  $\alpha = 2$  and we take the tessellation  $\{FT_{\delta-2}^{(1,2)}, BT_2^{(2,2)}, R_5 = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\delta - 1, 2) \rightarrow (2, 2) \rightarrow (2, 0), R_6 = (1, 2) \rightarrow (1, \delta - 1) \rightarrow (0, \delta) \rightarrow (0, 2)\}$ .

Thirdly, suppose  $\gamma > 1$  and  $\delta - \alpha = 1$ . Thus  $\gamma + \delta - \alpha > 2$ , so  $R_2$  is not equivalent to  $Z_4$ . If  $\alpha \neq \gamma$ ,  $R_1$  is not equivalent to  $Z_0$ , thus  $\delta - \gamma = (\delta - \alpha) + (\alpha - \gamma) = 2$ ; take the tessellation  $\{FT_\gamma^{(\gamma,2)}, BT_2^{(\gamma+1,2)}, R_7 = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\delta + \gamma - 2, 2) \rightarrow (\gamma + 1, 2) \rightarrow (\alpha, 0), R_8 = (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha) \rightarrow (\gamma - 1, \delta - \gamma)\}$ . Otherwise  $\alpha = \gamma$ . When  $\beta = \gamma + 2$ , from the conditions for this subcase  $\gamma + \delta - \beta \leq \alpha - \gamma = 0$ , so,  $\delta \leq 2$ , hence, as  $\gamma < \delta$  for this case,  $\gamma \leq 1$ , a contradiction. Thus  $\beta \geq \gamma + 3$ ; take the tessellation  $\{FT_2^{(\alpha,0)}, R_2, R_9 = (\beta, 0) \rightarrow (\gamma, \delta) \rightarrow (\gamma, 2) \rightarrow (\gamma + 2, 0)\}$ .

In each of the above cases the given tessellation, together with Lemmas 3.1, 3.2 and 3.3, verify the existence of a second distinct proper triangulation of  $R$ .

**Subcase B2.1:**  $\gamma < \delta, \gamma + 2 \leq \beta, \gamma \leq \alpha, \alpha - \gamma < \gamma + \delta - \beta$  and  $\beta - \gamma > \delta - \alpha$ .

Note that  $\gamma > 0$  and  $\alpha \geq \gamma$  imply  $\alpha > 0$ . Also since  $\alpha - \gamma < \gamma + \delta - \beta$ , then  $\gamma + \delta - \beta \geq 1$ .

Consider the tessellation  $\{FT_{\beta-\gamma}^{(\gamma, \alpha-\gamma)}, R_1 = (\beta, \alpha - \gamma) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \alpha + \beta - 2\gamma), R_2 = (\gamma, \alpha - \gamma) \rightarrow (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha), R_3 = (\beta, 0) \rightarrow (\beta, \alpha - \gamma) \rightarrow (\gamma, \alpha - \gamma) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemma 3.1, there exists a proper triangulation of  $R$ .

Lemma 3.1 verifies the existence of a second proper triangulation of  $R$  unless  $R_1$  is equivalent to  $Z_1$ ,  $R_2$  is equivalent to  $Z_1$  and  $R_3$  is equivalent to one of  $Z_0$  or  $Z_2$ .

Suppose that  $R_1$  is equivalent to  $Z_1$ . This supposition together with the condition  $\gamma + 2 \leq \beta$  imply that  $\alpha - \gamma + 1 = \gamma + \delta - \beta$ .

If  $\alpha > \gamma$ , then take the tessellation  $\{FT_{\beta-\gamma}^{(\gamma, \alpha-\gamma+1)}, BT_2^{(\gamma+1, \alpha-\gamma+1)} R_4 = (\gamma, \alpha - \gamma + 1) \rightarrow (\gamma, \delta - \gamma) \rightarrow (0, \delta) \rightarrow (0, \alpha) \rightarrow (\gamma - 1, \alpha - \gamma + 1), R_5 = (\beta, 0) \rightarrow (\beta, \alpha - \gamma + 1) \rightarrow (\gamma + 1, \alpha - \gamma + 1) \rightarrow (\gamma + 1, \alpha - \gamma - 1) \rightarrow (\alpha, 0)\}$ .

If  $\alpha = \gamma$ , then  $R_3 = Z_0$  and  $\delta + \gamma - \beta = 1$ . In this case further suppose that  $R_2$  is equivalent to  $Z_1$ . Under this supposition either  $\gamma = 1$  or  $\delta - \alpha = 1$ . When  $\gamma = 1$ ,  $\alpha = 1$  and so the condition  $\delta + \gamma - \beta = 1$  implies  $\beta = \delta$ , or equivalently  $\beta - \gamma = \beta - 1 = \delta - 1 = \delta - \alpha$ , contradicting the condition that  $\beta - \gamma > \delta - \alpha$ . Hence  $\gamma > 1$ . If  $\delta - \alpha = 1$  and  $\gamma + 2 = \beta$ , the condition  $\delta + \gamma - \beta = 1$  implies  $\delta = 3$  and  $\alpha = 2$ , which in turn implies  $\gamma = 2$ ; in this case  $R$  is equivalent to  $X_4$  and there exists only one proper triangulation of  $R$ . So it is left to check the case where  $\alpha = \gamma$ ,  $\delta - \alpha = 1$  and  $\gamma + 2 < \beta$ . Under these conditions take the tessellation  $\{FT_{\beta-\gamma-1}^{(\gamma, 0)}, R_2, R_6 = (\beta, 0) \rightarrow (\beta, 1) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \beta - \gamma - 1) \rightarrow (\beta - 1, 0)\}$ .

In each of the above cases the given tessellation together with Lemmas 3.1 and 3.3, verify the existence of a second distinct proper triangulation of  $R$ .

**Subcase B2.2.1:**  $\gamma < \delta$ ,  $\gamma + 2 \leq \beta$ ,  $\gamma \leq \alpha$ ,  $\beta - \gamma = \delta - \alpha$  and  $\alpha \geq 2$ .

Note that  $\delta + \gamma - \beta = \alpha$ .

If  $2 \leq \gamma$ , then, as  $\gamma = \alpha + \beta - \delta$ ,  $2 \leq (\gamma + \delta - \beta) - (\alpha - \gamma)$ , so,  $\alpha - \gamma + 2 \leq \gamma + \delta - \beta$ . So, consider the tessellation  $\{FT_{\beta-\gamma}^{(\gamma-1, \alpha-\gamma+1)}, BT_2^{(\beta, \alpha-\gamma+2)}, R_1 = (\beta, \alpha - \gamma + 2) \rightarrow (\beta, \delta + \gamma - \beta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - \gamma) \rightarrow (\beta - 2, \alpha - \gamma + 2), R_2 = (\beta, 0) \rightarrow (\beta, \alpha - \gamma) \rightarrow (\beta - 1, \alpha - \gamma + 1) \rightarrow (\gamma - 1, \alpha - \gamma + 1) \rightarrow (\alpha, 0), R_3 = (\gamma - 1, \alpha - \gamma + 1) \rightarrow (\gamma - 1, \delta - \gamma + 1) \rightarrow (0, \delta) \rightarrow (0, \alpha)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a proper triangulation of  $R$ . As  $2 \leq \gamma$  and  $\gamma + 2 \leq \beta$ , there exists a second distinct proper triangulation of  $R_1$  and hence there exists a second distinct proper triangulation of  $R$ .

If  $\gamma = 1$ , then the subcase condition  $\alpha = \gamma + \delta - \beta$  implies  $\alpha - \gamma + 1 = \gamma + \delta - \beta$ . Consider the tessellation  $\{FT_{\beta-\gamma}^{(0, \alpha)}, BT_2^{(\beta, \alpha)}, R_4 = (\beta, \alpha) \rightarrow (1, \delta) \rightarrow (1, \delta - 1) \rightarrow (\beta - 1, \alpha), R_5 = (\beta, 0) \rightarrow (\beta, \alpha - 2) \rightarrow (\beta - 2, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a proper triangulation of  $R$ .

Note that  $R_4$  is equivalent to  $Z_2$ , so unless  $R_5$  is equivalent to  $Z_1$ , Lemmas 3.1 and 3.3 verify the existence of a second distinct proper triangulation of  $R$ .

Suppose  $R_5$  is equivalent to  $Z_1$ ; then  $\alpha = 2$  and  $\beta = 3$ . Since  $\delta + \gamma - \beta = \alpha$ ,  $\delta = 4$ . In this case  $R = (3, 0) \rightarrow (3, 2) \rightarrow (1, 4) \rightarrow (1, 3) \rightarrow (0, 4) \rightarrow (0, 2) \rightarrow (2, 0)$  and a second distinct proper triangulation of  $R$  is  $\{FT_2^{(1,2)}, FT_1^{(0,3)}, FT_1^{(0,2)}, FT_1^{(2,0)}, FT_1^{(2,1)}, BT_2^{(2,2)}, BT_1^{(1,3)}, BT_1^{(3,2)}, BT_1^{(3,1)}\}$ .

**Subcase B2.2.2:**  $\gamma < \delta$ ,  $\gamma + 2 \leq \beta$ ,  $\gamma \leq \alpha$ ,  $\beta - \gamma = \delta - \alpha$  and  $\alpha \leq 1$ .

From the conditions for this subcase,  $0 < \gamma \leq \alpha$ ; it follows that  $\alpha = \gamma = 1$ . In addition,  $\gamma + 2 \leq \beta$ , so  $3 \leq \beta = \delta$ . Consider the tessellation  $\{FT_{\delta-2}^{(0,2)}, BT_2^{(\beta-1,2)}, R_1 = (\beta - 1, 2) \rightarrow (1, \delta) \rightarrow (1, \delta - 1) \rightarrow (\beta - 2, 2), R_2 = (\beta, 0) \rightarrow (\beta, 1) \rightarrow (\beta - 1, 2) \rightarrow (\beta - 1, 0), R_3 = (\beta - 1, 0) \rightarrow (\beta - 3, 2) \rightarrow (0, 2) \rightarrow (0, 1) \rightarrow (1, 0)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a proper triangulation of  $R$ .

Note that both  $R_1$  and  $R_2$  are equivalent to  $Z_2$ . So unless  $R_3$  is equivalent to  $Z_2$  or  $Z_4$ , by Lemmas 3.1 and 3.3, there exists a second proper triangulation of  $R$ .

Suppose  $R_3$  is equivalent to  $Z_2$ ; then  $\beta = 3$  and so  $R$  is equivalent to  $X_3$ . In this case by inspection there does not exist a second distinct proper triangulation of  $R$ .

Suppose  $R_3$  is equivalent to  $Z_4$ ; then  $\beta = 4$ . Then  $R = (4, 0) \rightarrow (4, 1) \rightarrow (1, 4) \rightarrow (1, 3) \rightarrow (0, 4) \rightarrow (0, 1) \rightarrow (1, 0)$  and  $\{FT_2^{(2,0)}, FT_2^{(1,2)}, FT_1^{(0,1)}, FT_1^{(0,2)}, FT_1^{(0,3)}, FT_1^{(1,0)}, FT_1^{(3,1)}, BT_2^{(2,2)}, BT_1^{(1,1)}, BT_1^{(1,3)}, BT_1^{(3,2)}, BT_1^{(4,1)}\}$  is a second distinct proper triangulation of  $R$ .

**Subcase C:**  $\gamma < \delta$  and  $\beta = \gamma + 1$ .

Since  $\delta + \gamma - \beta \leq \alpha$ , it follows, from the conditions for this subcase, that  $\delta - \alpha \leq 1$ . However, from the conditions for this lemma,  $\alpha < \delta$ , so  $\delta = \alpha + 1$ .

Now, from the conditions for this subcase,  $0 < \delta - \gamma$  and  $\beta - 1 = \gamma$ , so  $0 < (\alpha + 1) - (\beta - 1)$ ; thus  $\beta - 2 < \alpha$ . From the conditions for this lemma,  $\alpha < \beta$ , so it follows that  $\alpha = \beta - 1$ .

Thus,  $R = X_1$  and by inspection there does not exist a proper triangulation of  $R$ .

**Subcase D:**  $\delta = \gamma$ .

In this case the region  $R$  is the union of the region  $R_1 = (\gamma, 0) \rightarrow (0, \gamma) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$  and the region  $R_2 = (\beta, 0) \rightarrow (\beta, 2\gamma - \beta) \rightarrow (\gamma, \gamma) \rightarrow (\gamma, 0)$ . Furthermore,  $R_1 \cap R_2 = \{(0, \gamma)\}$ . By Lemma 3.1, there exists a proper triangulation of  $R$  and if either one of  $R_1$  or  $R_2$  is not equivalent to  $Z_2$ , then there exists a second distinct proper triangulation. If  $R_1$  and  $R_2$  are both equivalent to  $Z_2$ , then  $R$  is equivalent to  $X_5$  and by inspection has only one possible proper triangulation. □

Now that we have established Lemma 3.4 we can use it in conjunction with Lemmas 3.1 and 3.2 to establish when a convex region with six sides has a proper triangulation and when it has at least two distinct proper triangulations.

**Lemma 3.5.** *Let  $R$  be a convex region with six corners (sides). Whenever  $R$  is not equivalent to  $Z_6$  then  $R$  has a proper triangulation; moreover, whenever  $R$  is not equivalent to  $Z_5$  or  $Z_6$  then  $R$  has a second distinct proper triangulation.*

PROOF: Recall that if a proper triangulation for some region,  $R$ , exists, then a proper triangulation exists for all regions equivalent to  $R$ .

The region  $R$  is equivalent to  $(\beta, 0) \rightarrow (\beta, \gamma - \delta) \rightarrow (\beta - \delta, \gamma) \rightarrow (0, \gamma) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$ , where  $0 < \alpha, \delta < \beta, \gamma$ .

Two cases are considered; Case  $A$  where  $\gamma - \delta \leq \alpha + \delta - \beta$  and Case  $B$  where  $\gamma - \delta > \alpha + \delta - \beta$ .

Case  $B$  has several additional subcases which are summarized in the following table.

|    |   |  |
|----|---|--|
| B1 | The condition $\alpha = \beta - \delta = \gamma - \delta = 1$ does not hold |  |
| B2 | $\alpha = \beta - \delta = \gamma - \delta = 1$                             | B2.1 The condition $\delta = \beta - \alpha = \gamma - \alpha = 1$ does not hold |
|    |   | B2.2 $\delta = \beta - \alpha = \gamma - \alpha = 1$                             |

**Subcase A:**  $\gamma - \delta \leq \alpha + \delta - \beta$ .

Consider the tessellation  $\{FT_{\beta+\gamma-\alpha-\delta}^{(\beta-\delta, \alpha+\delta-\beta)}, R_1 = (\alpha, 0) \rightarrow (\beta, 0) \rightarrow (\beta, \gamma - \delta) \rightarrow (2\beta + \gamma - \alpha - 2\delta, \alpha + \delta - \beta) \rightarrow (\beta - \delta, \alpha + \delta - \beta), R_2 = (\beta - \delta, \alpha + \delta - \beta) \rightarrow (\beta - \delta, \gamma) \rightarrow (0, \gamma) \rightarrow (0, \alpha)\}$  of  $R$ . By Lemmas 3.1 and 3.3 there exists a proper triangulation of  $R$ . Moreover if neither  $R_1$  nor  $R_2$  is equivalent to an element of  $\{Z_0, Z_2, Z_4\}$ , then there exists a second distinct proper triangulation of  $R$ .

As  $\alpha, \delta < \beta, \gamma$ , both  $R_1$  and  $R_2$  are not equivalent to  $Z_0$ . Let  $R_1$  and  $R_2$  both be equivalent to  $Z_2$ . Then  $1 = \beta - \delta = \gamma - \delta = \alpha + \delta - \beta$ , thus,  $\beta = \gamma = \delta + 1$  and  $\alpha = 2$ .

From the condition for this case  $\gamma + \beta - \alpha \leq 2\delta$ . Recall that  $0 < \alpha < \beta, \gamma$ , so  $1 \leq \beta - \alpha$  and  $2 \leq \gamma$ . Hence,  $3 \leq 2\delta$ , thus,  $2 \leq \delta$ . Consider the tessellation  $\{FT_{\beta-2}^{(2,1)}, BT_2^{(2,2)}, R_3 = (2, 0) \rightarrow (\beta, 0) \rightarrow (\beta, 1) \rightarrow (2, 1), R_4 = (0, 2) \rightarrow (2, 2) \rightarrow (2, \gamma - 1) \rightarrow (1, \gamma) \rightarrow (0, \gamma)\}$  of  $R$ . By Lemmas 3.1 and 3.3 there exists a second distinct proper triangulation of  $R$ .

As  $R_2$  has four corners it is not equivalent to  $Z_4$ . Let  $R_1$  be equivalent to  $Z_2$  and  $R_2$  be equivalent to  $Z_4$  then  $(\alpha, \beta, \gamma, \delta) = (3, 4, 4, 3)$  and,  $\{FT_2^{(2,1)}, FT_1^{(3,0)}, FT_1^{(0,3)}, FT_1^{(1,3)}, BT_2^{(2,3)}, BT_1^{(3,1)}, BT_1^{(4,1)}, BT_1^{(1,4)}\}$  is a second distinct proper triangulation of  $R$ .

**Subcase B1:**  $\gamma - \delta > \alpha + \delta - \beta$  and the condition  $\alpha = \beta - \delta = \gamma - \delta = 1$  does not hold.

Consider the tessellation  $\{FT_\delta^{(\beta-\delta, \gamma-\delta)}, R_1 = (\alpha, 0) \rightarrow (\beta, 0) \rightarrow (\beta, \gamma - \delta) \rightarrow (\beta - \delta, \gamma - \delta) \rightarrow (\beta - \delta, \gamma) \rightarrow (0, \gamma) \rightarrow (0, \alpha)\}$  of  $R$ . As the condition  $\alpha = \beta - \delta = \gamma - \delta = 1$  does not hold,  $R_1$  is not equivalent to  $X_1$ . By Lemma 3.4, there exists a proper triangulation of  $R$  and if  $R_1 \notin \mathcal{X}$ , there exists a second distinct proper triangulation of  $R$ .

Let  $R_1 \in \mathcal{X}$ . As  $\gamma - \delta > \alpha + \delta - \beta$ ,  $R_1$  is not equivalent to  $X_5$ . Recall that  $\alpha < \gamma, \beta$ , so,  $R_1$  is not equivalent to  $X_2$ . Thus,  $R_1$  is equivalent to either  $X_3$  or  $X_4$ . If  $R_1$  is equivalent to  $X_4$ , then, by inspection, there exists a second distinct

proper triangulation of  $R$ . If  $R_1$  is equivalent to  $X_3$ , then  $R$  is equivalent to  $Z_5$  and by inspection there exists precisely one proper triangulation of  $R$ .

**Subcase B2.1:**  $\gamma - \delta > \alpha + \delta - \beta$ ,  $\alpha = \beta - \delta = \gamma - \delta = 1$  and the condition  $\delta = \beta - \alpha = \gamma - \alpha = 1$  does not hold.

Consider the linear transformation  $R\lambda + (\gamma, \beta)$  where  $\lambda = \begin{bmatrix} 0 & -1 \\ -1 & 0 \end{bmatrix} \in G$ . This transformation interchanges  $\beta$  with  $\gamma$  and  $\alpha$  with  $\delta$ . Thus, in this subcase the region  $R$  is equivalent to a region in Subcase B1.

**Subcase B2.2:**  $\gamma - \delta > \alpha + \delta - \beta$ ,  $\alpha = \beta - \delta = \gamma - \delta = \delta = \beta - \alpha = \gamma - \alpha = 1$ .

In this subcase  $(\alpha, \beta, \gamma, \delta) = (1, 2, 2, 1)$ , hence,  $R$  is equivalent to  $Z_6$  and, by inspection,  $R$  has no possible proper triangulation.  $\square$

We can now state the first major result of this paper.

**Theorem 3.6.** *Let  $R$  be any convex region. Then if  $R$  is not equivalent to  $Z_6$ , it has a proper triangulation. Moreover, whenever  $R \notin \mathcal{Z}$  then  $R$  has a second distinct proper triangulation.*

PROOF: This follows immediately from Lemmas 3.1, 3.3 and 3.5.  $\square$

#### 4. Distinct proper triangulations containing a fixed triangle

We now move on to the question of establishing when it is possible to find two distinct proper triangulations of a region when some triangle is forced to occur in both proper triangulations. We answer this question completely for three-sided regions, and also for non-rectangular four-sided regions. We deal with the four-sided regions first; the two choices for the direction of the internal triangle yield two theorems.

**Theorem 4.1.** *Let  $1 \leq \alpha \leq \beta$ ,  $0 < \chi \leq \alpha$ ,  $0 < \gamma \leq \beta - \chi$ ,  $0 \leq \delta \leq \alpha - \chi$  and  $\alpha \leq \gamma + \delta$ ; then there exists a proper triangulation of the region  $R = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$  which contains  $FT_\chi^{(\gamma, \delta)}$  if and only if  $(\alpha, \beta, \gamma, \delta, \chi) \neq (2, 3, 1, 1, 1)$ .*

*If  $\alpha = 1$  or  $(\alpha, \beta) = (2, 2)$  or  $(\alpha, \beta, \gamma, \delta, \chi) \in \{(2, 3, 2, 1, 1), (2, 3, 2, 0, 1), (2, 4, 2, 1, 1)\}$ , then there exists precisely one proper triangulations of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .*

*Otherwise there exists a second distinct proper triangulations of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .*


PROOF: If  $(\alpha, \beta, \gamma, \delta, \chi) = (2, 3, 1, 1, 1)$ , then, by inspection, there does not exist a proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .

If  $\alpha = 1$  or  $(\alpha, \beta) = (2, 2)$  or  $(\alpha, \beta, \gamma, \delta, \chi) \in \{(2, 3, 2, 1, 1), (2, 3, 2, 0, 1), (2, 4, 2, 1, 1)\}$ , then, by inspection, there exists precisely one proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .

By inspection, the other cases where  $\alpha + \beta \leq 6$  have two distinct proper triangulations of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .

Henceforth, assume that  $1 < \alpha$  and  $6 < \alpha + \beta$ . Two subcases are considered: Case *A* where  $\alpha \leq \gamma$  and Case *B* where  $\alpha > \gamma$  (see Figure 5 for illustrations of these cases and their subcases).

FIGURE 5. Tessellations for cases of Lemma 4.1.



Let  $m_1 = \min\{\gamma + \delta + \chi, \beta\}$  and  $m_2 = \min\{\gamma + \chi, \alpha\}$ .

**Case A:**  $\alpha \leq \gamma$ .

Note that for this case  $m_2 = \alpha$ .

Consider the tessellation  $\{BT_\alpha^{(\gamma, \alpha)}, BT_{m_1 - \gamma}^{(m_1, \delta + \chi)}, FT_\chi^{(\gamma, \delta)}, R_1 = (\beta, 0) \rightarrow (m_1, \gamma + \delta + \chi - m_1) \rightarrow (\gamma + \chi, \delta) \rightarrow (\gamma, \delta) \rightarrow (\gamma, 0), R_2 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (\gamma, \alpha) \rightarrow (\gamma, \delta + \chi) \rightarrow (m_1, \delta + \chi) \rightarrow (m_1, 0), R_3 = (\gamma, 0) \rightarrow (\gamma - \alpha, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, there exists a proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ , and if  $\{R_1, R_2, R_3\} \not\subset \mathcal{Z}$ , then there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ . So, assume that  $\{R_1, R_2, R_3\} \subset \mathcal{Z}$ .


Consider the tessellation  $\{BT_{\delta + \chi}^{(\gamma, \delta + \chi)}, BT_{\alpha + m_1 - \gamma - \delta - \chi}^{(m_1, \alpha)}, FT_\chi^{(\gamma, \delta)}, R_1, R_4 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (m_1, \alpha) \rightarrow (m_1, 0), R_5 = (\gamma, \delta + \chi) \rightarrow (\gamma + \delta + \chi - \alpha, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0) \rightarrow (\gamma, 0) \rightarrow (\gamma - \delta - \chi, \delta + \chi)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, unless  $\delta + \chi = \alpha$ , there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .

If  $\delta + \chi = \alpha$  and  $\chi \neq 1$ , consider the tessellation  $\{BT_{\alpha - 1}^{(\gamma, \alpha - 1)}, BT_{m_1 - \gamma}^{(m_1, \alpha)}, BT_1^{(\gamma, \alpha)}, FT_\chi^{(\gamma, \delta)}, R_1, R_4, R_6 = (\gamma, \alpha - 1) \rightarrow (\gamma - 1, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0) \rightarrow (\gamma, 0) \rightarrow (\gamma - \alpha + 1, \alpha - 1)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma, \delta)}$ .

If  $\delta + \chi = \alpha$ ,  $\chi = 1$  and  $\gamma + \chi < \beta$ , recall that  $\{R_1, R_2, R_3\} \subset \mathcal{Z}$ , hence,  $\alpha \leq 3$  and  $\beta \leq 6$ . For  $6 < \alpha + \beta \leq 9$  a pair of proper triangulations of  $R$  both containing  $FT_\chi^{(\gamma, \delta)} = FT_1^{(\gamma, \delta)}$  is shown in Figure 6.



FIGURE 6. Proper triangulations for Subcase A of Lemma 4.1.



If  $\delta + \chi = \alpha$ ,  $\chi = 1$  and  $\gamma + \chi = \beta$ , consider the tessellation  $\{BT_{\alpha-1}^{(\beta,\delta)}, BT_1^{(\gamma,\alpha)}, BT_1^{(\beta,\alpha)}, FT_\chi^{(\gamma,\delta)}, R_7 = (\beta, 0) \rightarrow (\beta - \alpha + 1, \delta) \rightarrow (\gamma, \delta) \rightarrow (\gamma - 1, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemma 3.2, there exists a second distinct proper triangulation of  $R$ .

**Case B:**  $\alpha > \gamma$ .

Consider the tessellation  $\{BT_{m_2+\delta-\alpha}^{(m_2,\delta)}, BT_{\alpha-\delta}^{(\gamma,\alpha)}, BT_{m_1-\gamma}^{(m_1,\delta+\chi)}, FT_\chi^{(\gamma,\delta)}, R_2, R_8 = (\beta, 0) \rightarrow (m_1, \gamma + \delta + \chi - m_1) \rightarrow (\gamma + \chi, \delta) \rightarrow (m_2, \delta) \rightarrow (m_2, \alpha - m_2) \rightarrow (\alpha, 0), R_9 = (\gamma, \delta) \rightarrow (\gamma + \delta - \alpha, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha - \delta, \delta)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, there exists a proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$ , and if  $\{R_2, R_8, R_9\} \not\subset \mathcal{Z}$ , then there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$ .

So, assume that  $\{R_2, R_8, R_9\} \subset \mathcal{Z}$ .


Recall the above assumption that  $6 < \alpha + \beta$ . Consider the tessellation  $\{BT_{m_2+\delta-\alpha}^{(m_2,\delta)}, BT_\chi^{(\gamma,\delta+\chi)}, BT_{\alpha+m_1-\gamma-\delta-\chi}^{(m_1,\alpha)}, FT_\chi^{(\gamma,\delta)}, R_4, R_8, R_{10} = (\gamma, \delta + \chi) \rightarrow (\gamma + \delta + \chi - \alpha, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha - \delta, \delta) \rightarrow (\gamma, \delta) \rightarrow (\gamma - \chi, \delta + \chi)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, unless  $\delta + \chi = \alpha$ , there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$ .

If  $\delta + \chi = \alpha$  and  $\chi \neq 1$ , consider the tessellation  $\{BT_{m_2+\delta-\alpha}^{(m_2,\delta)}, BT_{\chi-1}^{(\gamma,\alpha-1)}, BT_{m_1-\gamma}^{(m_1,\alpha)}, BT_1^{(\gamma,\alpha)}, FT_\chi^{(\gamma,\delta)}, R_4, R_8, R_{11} = (\gamma, \alpha - 1) \rightarrow (\gamma - 1, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha - \delta, \delta) \rightarrow (\gamma, \delta) \rightarrow (\gamma - \chi + 1, \alpha - 1)\}$  of  $R$ . By Lemmas 3.1, 3.2 and 3.3, there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$ .

Otherwise  $\delta + \chi = \alpha$  and  $\chi = 1$ . Since  $\alpha > \gamma$  for this subcase,  $m_2 = \gamma + 1$ . Observe that  $R_8$  is equivalent to one of  $Z_0, Z_1, Z_2$  or  $Z_4$ .

If  $R_8$  is equivalent to  $Z_4$ , then  $\alpha = \beta - 1 = 4$  and a pair of proper triangulations of  $R$  both containing  $FT_\chi^{(\gamma,\delta)} = FT_1^{(\gamma,\delta)}$  is shown in Figure 7.

FIGURE 7. Proper triangulations for Subcase  $B$  of Lemma 4.1.



Otherwise  $R_8$  is equivalent to  $Z_0, Z_1$  or  $Z_2$ . Thus either  $\gamma = 1$  or  $\beta - \gamma \leq 2$ .

When  $R_8$  is equivalent to  $Z_0$  and  $\gamma = 1$  then  $\beta \leq 2$ . But from the conditions of this lemma,  $\alpha \leq \beta$  and from above  $6 < \alpha + \beta$ , creating a contradiction.

When  $R_8$  is equivalent to  $Z_1$  or  $Z_2$  and  $\gamma = 1$  consider the tessellation  $\{BT_\delta^{(\alpha,\delta)}, BT_1^{(1,\alpha)}, BT_1^{(2,\alpha)}, FT_\chi^{(\gamma,\delta)} = FT_1^{(1,\alpha-1)}, R_{11} = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (2, \alpha) \rightarrow (2, \alpha - 1) \rightarrow (\alpha, \delta) \rightarrow (\alpha, 0)\}$  of  $R$ . As  $R_8$  is equivalent to  $Z_1$  or  $Z_2$  and  $R_2 \in \mathcal{Z}$ ,  $\beta \leq \alpha + 2$ . So, if  $\alpha \leq 2$ , then  $\alpha + \beta \leq 6$  contradicting our assumption that  $6 < \alpha + \beta$ . Thus, by Lemmas 3.1 and 3.2, we have a proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$  which is distinct to the one above.

Otherwise  $1 < \gamma$  and  $\beta - \gamma \leq 2$ . Note that  $\alpha = \delta + \chi < \delta + \gamma$ . Suppose that  $\gamma - 1 \geq \beta - \gamma$ . (Equivalently, as  $\chi = 1, \gamma + \delta - \beta \geq \alpha - \gamma$ .) Then consider the tessellation  $\{BT_\gamma^{(\gamma,\alpha)}, BT_{\beta-\gamma}^{(\beta,\alpha)}, FT_{\beta-\gamma}^{(\gamma,\gamma+\delta-\beta)}, FT_\chi^{(\gamma,\delta)} = FT_1^{(\gamma,\alpha-1)}, R_{13} = (\beta, 0) \rightarrow (\beta, \gamma + \delta - \beta) \rightarrow (\gamma, \gamma + \delta - \beta) \rightarrow (\gamma, \alpha - \gamma) \rightarrow (\alpha, 0), R_{14} = (\beta, \gamma + \delta - \beta) \rightarrow (\beta, \alpha + \gamma - \beta) \rightarrow (\gamma + 1, \delta) \rightarrow (\gamma, \delta)\}$  of  $R$ . By Lemmas 3.1 and 3.3, there exists a second distinct proper triangulation of  $R$  containing  $FT_\chi^{(\gamma,\delta)}$ .

Otherwise  $\gamma - 1 < \beta - \gamma$ . Then  $2 = \gamma < \alpha \leq \beta \leq 4$ . A pair of proper triangulations of  $R$  both containing  $FT_\chi^{(\gamma,\delta)} = FT_1^{(\gamma,\delta)}$  is shown in Figure 7.  $\square$

**Theorem 4.2.** Let  $0 < \chi \leq \delta \leq \alpha \leq \beta; \gamma \leq \beta; \alpha \leq \gamma + \delta - \chi$  and  $R = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0)$ .

If  $1 = \beta - \gamma = \beta - \alpha = \delta - \chi$ , then there does not exist a proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

Otherwise, if  $1 = \alpha$  or  $(\alpha, \beta, \gamma, \delta, \chi) = (2, 3, 3, 2, 1)$  or  $\alpha - \delta, \beta - \gamma, \gamma + \delta - \alpha - \chi \leq 1$ , then there exists precisely one proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

In cases other than the above, there exists at least two distinct proper triangulations of  $R$  both containing  $BT_\chi^{(\gamma,\delta)}$ .

PROOF: If  $1 = \beta - \gamma = \beta - \alpha = \delta - \chi$ , then, by inspection, there does not exist a proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

Similarly, if  $1 = \alpha$  or  $(\alpha, \beta, \gamma, \delta, \chi) = (2, 3, 3, 2, 1)$  or  $\alpha - \delta, \beta - \gamma, \gamma + \delta - \alpha - \chi \leq 1$ , then, by inspection, there exists precisely one proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .


By inspection, the other cases where  $\alpha + \beta \leq 6$  have two distinct proper triangulations of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

Henceforth, assume that  $1 < \alpha$ ,  $6 < \alpha + \beta$  and either  $1 < \alpha - \delta$  or  $1 < \beta - \gamma$  or  $1 < \gamma + \delta - \alpha - \chi$ .

Let  $m_1 = \min\{\gamma + \delta, \beta\}$ ,  $m_2 = \min\{\delta + \chi, \alpha\}$  and  $m_3 = \max\{\alpha + \chi - \gamma, 0\}$ .

Consider the tessellation  $\{BT_{\alpha+m_1-\gamma-\delta}^{(m_1,\alpha)}, BT_{m_2-m_3}^{(\gamma-\chi,m_2)}, BT_\chi^{(\gamma,\delta)}, R_1 = (m_1, 0) \rightarrow (m_1, \gamma + \delta - m_1) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - \chi) \rightarrow (\gamma - \chi, \delta) \rightarrow (\gamma - \chi, m_3) \rightarrow (\alpha, 0), R_2 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (m_1, \alpha) \rightarrow (m_1, 0), R_3 = (\gamma - \chi, m_3) \rightarrow (\gamma + m_3 - m_2 - \chi, m_2) \rightarrow (\gamma - \chi, m_2) \rightarrow (\gamma + \delta - \alpha, \alpha) \rightarrow (0, \alpha) \rightarrow (\alpha, 0), R_4 = (\gamma, \delta) \rightarrow (\gamma + \delta - m_2, m_2) \rightarrow (\gamma - \chi, m_2) \rightarrow (\gamma - \chi, \delta)\}$  of  $R$  (see Figure 8 for an illustration of these cases). Since the condition  $1 = \beta - \gamma = \beta - \alpha = \delta - \chi$  does not hold, the region  $R_1$  is not equivalent to the region  $X_1$ . By Lemmas 3.1, 3.2, 3.3 and 3.4, there exists a proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ . Furthermore if  $\{R_1, R_2, R_3, R_4\} \not\subset \mathcal{X} \cup \mathcal{Z}$ , then there exists a second distinct proper triangulation.

FIGURE 8. Tessellations for cases of Lemma 4.2.



Otherwise  $\{R_1, R_2, R_3, R_4\} \subset \mathcal{X} \cup \mathcal{Z}$  and we wish to establish the existence of a second distinct proper triangulation when  $1 < \alpha$ . For  $6 < \alpha + \beta \leq 12$  there are a small number of cases; these are dealt with individually in [11]. Henceforth

we assume that  $12 < \alpha + \beta$ . Three cases are considered: Case *A* where  $\chi = 1$ , Case *B* where  $\chi = 2$  and Case *C* where  $3 \leq \chi$ .

**Case A:**  $\chi = 1$ .

Suppose first that  $\gamma \neq \beta$  and  $\chi = 1 < \gamma + \delta - \alpha$ .

From Figure 8,  $R_3$  is equivalent to  $Z_0$  or  $Z_1$  or  $Z_3$ . Thus  $\alpha - \delta \leq 2$  and  $\gamma - \alpha - \chi \leq 1$ . Since  $R_2$  is equivalent to  $Z_0$  or  $Z_1$ ,  $\beta - m_1 \leq 1$ . Since  $R_1 \in Z \cup \mathcal{X}$  and  $\chi = 1 < \gamma + \delta - \alpha$ , the region  $R_1$  is equivalent to either  $Z_3$  or  $X_2$  or  $X_3$  or  $X_5$ .

If  $R_1$  is equivalent to  $Z_3$ , then  $\alpha = \beta$  and  $\delta = 3$ . Thus,  $\alpha = \beta \leq \delta + 2 = 5$ , so  $\alpha + \beta \leq 10$ , a contradiction.

If  $R_1$  is equivalent to  $X_2$ , then  $\delta = 2$  and either  $m_1 - \gamma = 1$ , in which case  $m_1 = \beta$  and  $m_1 - \alpha \leq 3$ , or  $m_1 - \gamma = 2$ , in which case  $\beta - m_1 \leq 1$  and  $m_1 - \alpha = 2$ . Thus,  $\alpha \leq \delta + 2 = 4$  and  $\beta - \alpha \leq 3$ , so,  $\beta \leq 7$ . Hence  $\alpha + \beta \leq 11$ , a contradiction.

If  $R_1$  is equivalent to  $X_3$ , then  $\delta = 3$ ,  $\alpha + \chi - \gamma = 1$  and  $\beta + \chi - \gamma = 3$ . Thus,  $\alpha \leq \delta + 2 = 5$  and  $\beta = 2 + \alpha \leq 7$ . So  $\alpha + \beta \leq 12$ , a contradiction.

If  $R_1$  is equivalent to  $X_5$ , then  $\delta = 1$ ,  $\beta - m_1 \leq 1$ ,  $0 \leq \gamma - \alpha - \chi \leq 1$  and  $m_1 - \gamma = 1$ . Thus,  $\alpha \leq \delta + 2 = 3$  and  $\beta \leq \alpha + 4 \leq 7$ . So,  $\alpha + \beta \leq 10$ , a contradiction.

Secondly, suppose that  $\gamma \neq \beta$  and  $\gamma + \delta - \alpha = \chi = 1$ . Hence,  $R_1$  is equivalent to  $Z_1$  or  $Z_2$ .

If  $3 \leq \gamma$ , then consider the tessellation  $\{BT_2^{(2,\alpha)}, BT_{m_1-2}^{(m_1,\alpha-1)}, BT_\chi^{(\gamma,\delta)} = BT_1^{(\gamma,\alpha-\gamma+1)}, R_1, R_5 = (\gamma, \delta) \rightarrow (2, \alpha - 1) \rightarrow (2, \alpha - 2) \rightarrow (\gamma - 1, \delta), R_6 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (2, \alpha) \rightarrow (2, \alpha - 1) \rightarrow (m_1, \alpha - 1) \rightarrow (m_1, 0)\}$  of  $R$ . By Lemmas 3.1 and 3.2, there exists a second distinct proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

If  $\gamma \leq \alpha - 2$ , consider the tessellation  $\{BT_2^{(\alpha,2)}, BT_{\alpha-2}^{(\alpha-1,\alpha)}, BT_\chi^{(\gamma,\delta)} = BT_1^{(\gamma,\alpha-\gamma+1)}, R_7 = (\alpha - 1, 2) \rightarrow (\gamma, \delta) \rightarrow (\gamma, \delta - 1) \rightarrow (\alpha - 2, 2), R_8 = (\gamma, \delta) \rightarrow (1, \alpha) \rightarrow (0, \alpha) \rightarrow (\gamma - 1, \delta), R_9 = (\beta, 0) \rightarrow (\beta, \alpha) \rightarrow (\alpha - 1, \alpha) \rightarrow (\alpha - 1, 2) \rightarrow (\alpha, 2) \rightarrow (\alpha, 0)\}$  of  $R$ . By Lemmas 3.1 and 3.2, there exists a second distinct proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

Otherwise  $\alpha - 2 < \gamma < 3$ , so  $\alpha \leq 3$ . As  $R_2$  is equivalent to  $Z_0$  or  $Z_1$  and  $\gamma + \delta - \alpha = \chi = 1$  it follows that  $\beta \leq \alpha + 2$ . Thus,  $\alpha + \beta \leq 8$ , a contradiction.

Thirdly, suppose that  $\gamma = \beta$  and  $\gamma + \delta - \alpha = \chi = 1$ ; then  $\delta - 1 = \alpha - \beta$ . From the conditions for this lemma  $0 \leq \delta - 1$  and  $\alpha - \beta \leq 0$ . Thus  $\alpha = \beta$  and  $\delta = 1$ . Note that  $6 < \alpha + \beta$ , so,  $4 \leq \alpha = \beta$ . Consider the tessellation  $\{BT_{\alpha-1}^{(\alpha-1,\alpha)}, BT_\chi^{(\gamma,\delta)} = BT_1^{(\alpha,1)}, R_{10} = (\alpha, 1) \rightarrow (\alpha, \alpha) \rightarrow (\alpha - 1, \alpha) \rightarrow (\alpha - 1, 1)\}$  of  $R$ . By Lemma 3.1, there exists a second distinct proper triangulation of  $R$  containing  $BT_\chi^{(\gamma,\delta)}$ .

Finally assume  $\gamma = \beta$  and  $\chi = 1 < \gamma + \delta - \alpha$ . Recall that  $R_3$  is equivalent to  $Z_0$ ,  $Z_1$  or  $Z_3$ , so,  $\alpha - \delta \leq 2$  and  $\gamma - \alpha - \chi \leq 1$ . Hence, as  $\gamma = \beta$  and  $\chi = 1$ , it follows that  $\beta - 2 \leq \alpha$ . If  $\alpha \leq 2$ , then  $\beta \leq 4$  and thus  $\alpha + \beta \leq 6$ ; therefore,  $3 \leq \alpha$ .

If  $\beta - 2 < \alpha + \chi - \delta$ , then  $\beta - 2 < 2 + \chi$ , hence,  $\beta < 5$ , thus,  $6 < \alpha + \beta \leq 8$ , a contradiction.

Hence, assume  $\alpha + \chi - \delta \leq \beta - 2$ .

If  $\alpha - 2 - (\alpha - (\beta - 2)) < 0$ , then  $\beta < 4$ , so,  $\alpha + \beta \leq 6$ , a contradiction. Hence,  $\alpha - 2 - (\alpha - (\beta - 2)) \geq 0$ .

Let  $m_4 = \min\{\alpha - 2, \delta - 1\}$ . Consider the tessellation  $\{FT_2^{(\beta-2, m_4)}, BT_{\beta-2}^{(\beta-2, \alpha)}, BT_\chi^{(\gamma, \delta)} = BT_1^{(\beta, \delta)}, R_{11} = (\beta, 0) \rightarrow (\beta, m_4) \rightarrow (\beta - 2, m_4) \rightarrow (\beta - 2, \alpha - \beta + 2) \rightarrow (\alpha, 0), R_{12} = (\beta, m_4) \rightarrow (\beta, \delta - 1) \rightarrow (\beta - 1, \delta) \rightarrow (\beta, \delta) \rightarrow (\beta, \alpha) \rightarrow (\beta - 2, \alpha) \rightarrow (\beta - 2, m_4 + 2)\}$  of  $R$ . By Lemmas 3.1 and 3.3 there exists a second distinct proper triangulation of  $R$  containing  $BT_\chi^{(\gamma, \delta)}$ .

**Case B:**  $\chi = 2$ .

As  $R_3$  is equivalent to  $Z_0, Z_1$  or  $Z_3$  and  $R_4$  is equivalent to  $Z_0$  or  $Z_2, \alpha - \delta \leq 1$  (so,  $m_2 = \alpha$ ) and  $\gamma - \alpha - \chi \leq 1$ . Since  $R_1$  is equivalent to  $Z_0, Z_1, Z_2, Z_3, Z_4, X_4$  or  $X_5, m_1 = \beta, \beta - \alpha \leq 2, m_1 + \chi - \gamma \leq 4$  and  $\delta \leq 4$ . If  $\delta = 4$ , then  $R_1$  is equivalent to  $Z_3$ , so,  $\alpha = \beta$  and  $\alpha - \delta, \beta - \gamma, \gamma + \delta - \alpha - \chi \leq 1$  contradicting our assumption that either  $1 < \alpha - \delta$  or  $1 < \beta - \gamma$  or  $1 < \gamma + \delta - \alpha - \chi$ . So,  $\delta \leq 3$ . Thus,  $\alpha \leq 4$  and  $\beta \leq 2 + \alpha$ , so,  $\alpha + \beta \leq 10$ , a contradiction.

**Case C:**  $3 \leq \chi$ .

As  $R_i \in \mathcal{X} \cup \mathcal{Z}$  for all  $1 \leq i \leq 4$  it follows that  $\beta - \gamma, \alpha - \delta, \gamma + \delta - \alpha - \chi \leq 1$ , a contradiction. □

Letting  $\alpha = \beta$  in Theorem 4.1 and 4.2 describes precisely when at least one or two proper triangulations exist for a three-sided region containing any fixed triangle.

Note that, this result is described in terms of finding proper triangulations of a backward triangle. However, the equivalences of regions (and indeed proper triangulations) discussed earlier mean that a similar result holds for forward triangles.

REFERENCES

- [1] Cavenagh N.J., *Latin trades and critical sets in latin squares*, PhD Thesis, University of Queensland, Australia, 2003.
- [2] Cavenagh N.J., Donovan D.M., Khodkar A., Lefevre J.G., McCourt T.A., *Identifying flaws in the security of critical sets in latin squares via triangulations*, Australas. J. Combin. **52** (2012), 243–268.
- [3] Drápal A., *On a planar construction of quasigroups*, Czechoslovak Math. J. **41** (1991), no. 3, 538–548.
- [4] Drápal A., *Hamming distances of groups and quasi-groups*, Discrete Math. **235** (2001), no. 1–3, 189–197.
- [5] Drápal A., Hämäläinen C., *An enumeration of equilateral triangle dissections*, Discrete Applied Math. **158** (2010), no. 14, 1479–1495.
- [6] Drápal A., Hämäläinen C., Kala V., *Latin bitrades, dissections of equilateral triangles and abelian groups*, J. Combin. Des. **18** (2010), no. 1, 1–24.
- [7] Keedwell A.D., *Critical sets in latin squares and related matters: an update*, Util. Math. **65** (2004), 97–131.


- [8] Laczkovich M., *Tilings of polygons with similar triangles*, *Combinatorica* **10** (1990), no. 3, 281–306.
- [9] Laczkovich M., *Tilings of triangles*, *Discrete Math.* **140** (1995), no. 1–3, 79–94.
- [10] Laczkovich M., *Tilings of polygons with similar triangles, II*, *Discrete Comput. Geom.* **19** (1998), no. 3, Special Issue, 411425, dedicated to the memory of Paul Erdős.
- [11] McCourt T.A., *On defining sets in latin squares and two intersection problems, one for latin squares and one for Steiner triple systems*, PhD Thesis, University of Queensland, Australia, 2010.
- [12] Tutte W.T., *The dissection of equilateral triangles into equilateral triangles*, *Proc. Cambridge Philos. Soc.* **44** (1948), 463–482.

### Appendix


Regions equivalent to  $Z_1$




Regions equivalent to  $Z_2$



Regions equivalent to  $Z_3$



Regions equivalent to  $Z_4$




Regions equivalent to  $Z_5$




A Region equivalent to  $Z_6$



Regions equivalent to  $X_1$




Regions equivalent to  $X_2$




Regions equivalent to  $X_3$



Regions equivalent to  $X_4$



Regions equivalent to  $X_5$



Diane M. Donovan, James G. Lefevre, Thomas A. McCourt:  
SCHOOL OF MATHEMATICS AND PHYSICS, UNIVERSITY OF QUEENSLAND, BRISBANE,  
QLD, 4072, AUSTRALIA

Nicholas J. Cavenagh:  
DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WAIKATO, PRIVATE BAG 3105,  
HAMILTON, NEW ZEALAND

(Received June 1, 2011, revised January 27, 2012)