

Hanbiao Yang

Metrization of function spaces with the Fell topology

Commentationes Mathematicae Universitatis Carolinae, Vol. 53 (2012), No. 2, 307--318

Persistent URL: <http://dml.cz/dmlcz/142891>

Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2012

Institute of Mathematics of the Academy of Sciences of the Czech Republic provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This paper has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://project.dml.cz>

Metrization of function spaces with the Fell topology

HANBIAO YANG

Abstract. For a Tychonoff space X , let $\downarrow C_F(X)$ be the family of hypographs of all continuous maps from X to $[0, 1]$ endowed with the Fell topology. It is proved that X has a dense separable metrizable locally compact open subset if $\downarrow C_F(X)$ is metrizable. Moreover, for a first-countable space X , $\downarrow C_F(X)$ is metrizable if and only if X itself is a locally compact separable metrizable space. There exists a Tychonoff space X such that $\downarrow C_F(X)$ is metrizable but X is not first-countable.

Keywords: space of continuous maps, Fell topology, hyperspace, metrizable, hypograph, separable, first-countable

Classification: 54C35, 54E45, 54B20

1. Introduction and main results

For a topological space X , let $C(X)$ denote the set of all continuous maps from X to the unit closed interval $\mathbf{I} = [0, 1]$ with the usual topology. Then we can endow $C(X)$ with various topologies. For example, the topology of uniform convergence, the topology of pointwise convergence and the compact-open topology are well known. In [4]–[10], $C(X)$ is endowed with other natural topologies inherited from the spaces $\text{Cld}(X \times \mathbf{I})$ of nonempty closed sets in $X \times \mathbf{I}$.

For a space Y , let $\text{Cld}(Y)$ be the set of all nonempty closed sets in Y . For an open set U in Y , let

$$U^- = \{A \in \text{Cld}(Y) : A \cap U \neq \emptyset\} \quad \text{and} \quad U^+ = \{A \in \text{Cld}(Y) : A \subset U\}.$$

The most well-known topology of $\text{Cld}(Y)$, called the *Vietoris topology*, is generated by

$$\{U^-, U^+ : U \text{ is open in } Y\}.$$

In this paper, we consider the *Fell topology* of $\text{Cld}(Y)$, which is generated by

$$\{U^-, (Y \setminus K)^+ : U \text{ is open and } K \text{ is compact in } Y\}.$$

The hyperspaces $\text{Cld}(Y)$ with the above two topologies are denoted by $\text{Cld}_V(Y)$ and $\text{Cld}_F(Y)$, respectively. It is well-known that $\text{Cld}_V(Y)$ (resp. $\text{Cld}_F(Y)$) is metrizable if and only if Y is a compact (resp. locally compact and separable) metrizable space. Obviously, when Y is compact, the Fell topology of $\text{Cld}(Y)$ is equal to the Vietoris topology.

For every $f \in C(X)$, let

$$\downarrow f = \{(x, s) \in X \times \mathbf{I} : s \leq f(x)\} \in \text{Cld}(X \times \mathbf{I}),$$

which is called the *hypograph* of f . By identifying each $f \in C(X)$ with $\downarrow f \in \text{Cld}_V(X \times \mathbf{I})$, we can regard $C(X)$ as the subset

$$\downarrow C(X) = \{\downarrow f : f \in C(X)\} \subset \text{Cld}(X \times \mathbf{I}).$$

Let $\downarrow C_V(X)$ and $\downarrow C_F(X)$ be the spaces with the topologies inherited from $\text{Cld}_V(X \times \mathbf{I})$ and $\text{Cld}_F(X \times \mathbf{I})$, respectively. These topologies are different from the three topologies mentioned previously (see [4, Corollary 1]). In [9, Theorem 1], it was proved that, for a Tychonoff space X , $\downarrow C_V(X)$ is metrizable if and only if $\downarrow C_V(X)$ is second-countable if and only if X is compact and metrizable. The following theorem is our main result.

Theorem 1. *For a Tychonoff space X , the following conditions are equivalent:*

- (a) $\downarrow C_F(X)$ is separable metrizable;
- (b) $\downarrow C_F(X)$ is metrizable.

In case X is first-countable, the above two conditions are equivalent to

- (c) X is a locally compact and separable metrizable space.

We also prove the following theorem.

Theorem 2. *Let $\bigoplus_{s \in S} Y_s$ be the topological sum of Tychonoff spaces Y_s , $s \in S$, and $a_s \in Y_s$ a non-isolated point for every $s \in S$. Let, further, Y be the quotient space of $\bigoplus_{s \in S} Y_s$ with the set $\{a_s : s \in S\}$ identified to a point. Then $\downarrow C_F(Y)$ is homeomorphic to a subspace of the product space $\prod_{s \in S} \downarrow C_F(Y_s)$.*

Applying this theorem, we show the following.

Corollary 1. *There exists a Tychonoff space X such that $\downarrow C_F(X)$ is separable metrizable but X is not first-countable.*

The above corollary shows that the first-countability of X is essential for the equivalence between (a) and (c) in Theorem 1. The following Theorem 3 tells us that, the non-compact case is very different from the compact one.

Theorem 3. *There exists a countable Tychonoff space X such that $\downarrow C_F(X)$ is Hausdorff and second-countable but not regular.*

In [1, 5.1.2 Proposition], it was proved that, for a Tychonoff space X , the following conditions are equivalent: (a) $\text{Cld}_F(X)$ is Hausdorff, (b) $\text{Cld}_F(X)$ is regular, (c) $\text{Cld}_F(X)$ is Tychonoff, and (d) X is locally compact. Theorem 3 shows that we cannot replace $\text{Cld}_F(X)$ by $\downarrow C_F(X)$ in [1, 5.1.2 Proposition].

The following Theorem 4 states that, even for a compact space X , the regularity and the first-countability of $\downarrow C_F(X)$ do not imply the metrizability of it.

Theorem 4. *There exists a compact space X such that $\downarrow C_F(X)$ is Tychonoff, separable and first-countable but not metrizable.*

Finally, we will give a necessary condition for the metrizability of $\downarrow C_F(X)$.

Theorem 5. *For a Tychonoff space X , if $\downarrow C_F(X)$ is metrizable, then there exists a dense, locally compact, open and separable metrizable subspace of X . But the converse is not true.*

2. Preparatory results

In the following, we always assume that X is a Tychonoff space and $p : X \times \mathbf{I} \rightarrow X$ is the projection. For $s \in \mathbf{I}$, we use \underline{s} to denote the constant function from X to \mathbf{I} which maps all elements to s . By \mathbb{R} and \mathbb{Q} , we denote the sets of all real numbers and of all rational numbers, respectively. Let cl_Y and int_Y be the closure-operator and the interior-operator in a space Y . If $Y = X$, the subscript in the above operators will be omitted. And, for a closed set F in Y , let

$$F^* = (Y \setminus F)^+ = \{A \in \text{Cl}_d(Y) : A \cap F = \emptyset\}.$$

By the definition, the topology of $\downarrow C_F(X)$ is generated, as a base, by the following sets:

$$\bigcap_{i=1}^n G_i^- \cap K^* \cap \downarrow C(X),$$

where G_1, G_2, \dots, G_n are open sets in $X \times (0, 1]$ and K is a compact set in $X \times (0, 1]$. In particular,

$$\left\{ \bigcap_{i=1}^n G_i^- \cap \downarrow C(X) : G_1, \dots, G_n \text{ are nonempty open in } X \times (0, 1] \right\}$$

and $\{K^* \cap \downarrow C(X) : K \text{ is compact in } X \times (0, 1]\}$

are neighborhood bases at $\downarrow \underline{1}$ and $\downarrow \underline{0}$ in $\downarrow C_F(X)$, respectively.

To prove our theorems, we need some lemmas. At first, we show the following lemma.

Lemma 1. *For a space X , the following hold:*

- (1) $\downarrow C_F(X)$ is T_1 ;
- (2) $\downarrow C_F(X)$ is Hausdorff if and only if there exists a dense open subset U of X which is locally compact.

PROOF: (1): Let $f \neq g \in C(X)$. We may assume that $f(x_0) < g(x_0)$ for some $x_0 \in X$. Then x_0 has an open neighborhood W such that $f(x) < a < g(x)$ for every $x \in W$, where $a = \frac{f(x_0)+g(x_0)}{2}$. Thus $\downarrow f \in (\{x_0\} \times [a, 1])^* \not\preceq \downarrow g$ and $\downarrow g \in (W \times (a, 1])^- \not\preceq \downarrow f$.

(2): The “if” part: Take $f, g \in C(X)$, $x_0 \in W$ and $a \in \mathbf{I}$ as the same as in (1). Since f and g are continuous, we assume that $x_0 \in U$. Because U is locally compact, we have an open set V in X such that $x_0 \in V \subset \text{cl}V \subset U \cap W$ and $\text{cl}V$ is compact. Since $f(x) < a < g(x)$ for $x \in \text{cl}V$, $(\text{cl}V \times [a, 1])^* \cap \downarrow C(X)$ and $(V \times (a, 1])^- \cap \downarrow C(X)$ are disjoint neighborhoods of $\downarrow f$ and $\downarrow g$, respectively.

The “only if” part: We define an open set

$$U = \bigcup \{ \text{int } K : K \text{ is compact in } X \} \subset X.$$

Then U is locally compact. We show that U is dense in X . Assume that U is not dense in X . Then there exists a nonempty open set V in X such that the interior of every compact subset of V is empty. Because X is Tychonoff, we can choose $f \in C(X)$ such that $f(X \setminus V) \subset \{1\}$ and $f(x_0) = 0$ for some $x_0 \in V$. Since $\downarrow C_F(X)$ is Hausdorff, there exist disjoint open sets \mathcal{U} and \mathcal{V} in $\downarrow C_F(X)$ such that $\downarrow 1 \in \mathcal{U}$ and $\downarrow f \in \mathcal{V}$. Then we can find nonempty open sets $G_1, G_2, \dots, G_n, \dots, G_m \subset X \times (0, 1]$ and a compact set $K \subset X \times (0, 1]$ such that

$$\begin{aligned} \downarrow 1 &\in G_1^- \cap G_2^- \cap \dots \cap G_n^- \cap \downarrow C(X) \subset \mathcal{U} \quad \text{and} \\ \downarrow f &\in G_{n+1}^- \cap \dots \cap G_m^- \cap K^* \cap \downarrow C(X) \subset \mathcal{V}. \end{aligned}$$

Since $f(X \setminus V) \subset \{1\}$, it follows that $p(K) \subset V$, which implies that $\text{int } p(K) = \emptyset$. For every $i \leq m$, $p(G_i) \setminus p(K) \neq \emptyset$ since $p(G_i)$ is a nonempty open set in X . Take $x_i \in p(G_i) \setminus p(K)$. Because X is Tychonoff, we have $g \in C(X)$ satisfying

$$g(x_i) = 1 \text{ for } i \leq m \text{ and } g(p(K)) = \{0\}.$$

Then $\downarrow g \in \mathcal{U} \cap \mathcal{V}$, which contradicts that $\mathcal{U} \cap \mathcal{V} = \emptyset$. □

Lemma 2. *If $\downarrow C_F(X)$ is first-countable, then there exist compact sets $C_1 \subset C_2 \subset \dots$ in X such that every compact set in X is contained in some C_n . In particular, $X = \bigcup_{n=1}^\infty C_n$.*

PROOF: Because $\downarrow C_F(X)$ is first-countable, we can find compact sets $K_1 \subset K_2 \subset \dots$ in $X \times (0, 1]$ such that $\{K_n^* \cap \downarrow C(X) : n = 1, 2, \dots\}$ is a neighborhood base of $\downarrow 0$ in $\downarrow C_F(X)$. Then $C_n = p(K_n)$, $n = 1, 2, \dots$, are the desired compact sets in X . We verify that every compact set C in X is contained in some C_n . Otherwise, for every n , we can choose $x_n \in C \setminus C_n$ and define $f_n \in C(X)$ such that $f_n(x_n) = 1$ and $f_n(C_n) = \{0\}$. Then $\downarrow f_n \in K_n^*$ for every n and hence $\downarrow f_n \rightarrow \downarrow 0$ in $\downarrow C_F(X)$. But every $\downarrow f_n$ is not contained in the neighborhood $(C \times \{1\})^*$ of $\downarrow 0$, which is a contradiction. □

Lemma 3. *If X and $\downarrow C_F(X)$ are first-countable, then X is locally compact.*

PROOF: Suppose there exists $x_0 \in X$, which has no compact neighborhood. Because X is first-countable, x_0 has a countable open neighborhood base $\{U_n : n = 1, 2, \dots\}$, where $U_n \supset U_{n+1}$ for every n . Since $\downarrow C_F(X)$ is also first-countable, we can find compact sets $K_1 \subset K_2 \subset \dots$ in $X \times (0, 1]$ such that $\{K_n^* \cap \downarrow C(X) : n = 1, 2, \dots\}$ is a neighborhood base at $\downarrow 0$ in $\downarrow C(X)$. By the assumption, $p(K_n) \not\subset U_n$ for every $n = 1, 2, \dots$, hence we can take $x_n \in U_n \setminus p(K_n)$. Then $x_n \rightarrow x_0$ in X . Since X is Tychonoff, we have $f_n \in C(X)$ such that

$$f_n(x_n) = 1 \text{ and } f_n(p(K_n) \cup (X \setminus U_n)) = \{0\}.$$

Then $\downarrow f_n \in K_n^*$ and hence $\downarrow f_n \rightarrow \downarrow 0$. On the contrary,

$$(\{x_n : n = 0, 1, 2, \dots\} \times \{1\})^* \cap \downarrow C(X)$$

is a neighborhood of $\downarrow 0$ in $\downarrow C_F(X)$ which does not contain any $\downarrow f_n$. □

When X is locally compact and non-compact, let $\alpha X = X \cup \{\infty\}$ be the one-point compactification of X . Using Lemmas 2 and 3, we may prove the following

Proposition 1. *If X and $\downarrow C_F(X)$ are first-countable, then*

- (1) *X is locally compact and αX is also first-countable;*
- (2) *$\downarrow C_F(\alpha X)$ is first-countable;*
- (3) *$\downarrow C_F(\alpha X)$ is second-countable if $\downarrow C_F(X)$ is second-countable.*

PROOF: The assertion (1) directly follows from Lemmas 2 and 3. To show (2) and (3), we only consider the case that X is not compact. Let $\{U_n : n = 1, 2, \dots\}$ be a countable open neighborhood base at ∞ in αX , and let $\phi : C(\alpha X) \rightarrow C(X)$ be the restriction, that is,

$$\phi(f) = f|X \text{ for every } f \in C(\alpha X).$$

Then it is not hard to verify that $\downarrow \phi : \downarrow C_F(\alpha X) \rightarrow \downarrow C_F(X)$ is a continuous injection. Unfortunately, it is not an embedding. However, the following \mathcal{S} is a subbase of $\downarrow C_F(\alpha X)$:

$$\begin{aligned} \mathcal{S} = & \{(\downarrow \phi)^{-1}(G) : G \in \mathcal{G}\} \\ & \cup \{(\text{cl}_{\alpha X} U_n \times [r, 1])^* \cap \downarrow C(\alpha X) : r \in \mathbb{Q} \cap (0, 1], n = 1, 2, \dots\}, \end{aligned}$$

where \mathcal{G} is an open base for $\downarrow C_F(X)$. Obviously, \mathcal{S} is a subfamily of the topology of $\downarrow C_F(\alpha X)$. For every open set V in $\alpha X \times \mathbf{I}$, $V \cap (X \times \mathbf{I})$ is open in $X \times \mathbf{I}$ and

$$V^- \cap \downarrow C(\alpha X) = (\downarrow \phi)^{-1}((V \cap (X \times \mathbf{I}))^- \cap \downarrow C(\alpha X)).$$

For every compact set K in $\alpha X \times (0, 1]$, if $K \cap (\{\infty\} \times \mathbf{I}) = \emptyset$, then K is also compact in $X \times \mathbf{I}$ and

$$K^* \cap \downarrow C(\alpha X) = (\downarrow \phi)^{-1}(K^* \cap \downarrow C(X)).$$

If $K \cap (\{\infty\} \times \mathbf{I}) \neq \emptyset$, then for every $\downarrow f \in K^* \cap \downarrow C(\alpha X)$, using the Wallace's Theorem, there exist n and a rational number $r \in (0, 1]$ such that

$$\begin{aligned} & (\text{cl}_{\alpha X} U_n \times [r, 1]) \cap \downarrow f = \emptyset \text{ and} \\ & K \cap (\text{cl}_{\alpha X} U_n \times \mathbf{I}) \subset \text{cl}_{\alpha X} U_n \times [r, 1]. \end{aligned}$$

Let

$$K_1 = (K \cap ((\alpha X \setminus U_n) \times \mathbf{I})) \cup (\text{cl}_{\alpha X} U_n \times [r, 1]).$$

Then K_1 is compact in $\alpha X \times (0, 1]$, $K_1 \supset K$ and $K_1 \cap \downarrow f = \emptyset$. Thus, $\downarrow f \in K_1^* \subset K^*$. Note that

$$K_1^* \cap \downarrow C_F(\alpha X) = (\downarrow \phi)^{-1}((K \cap ((\alpha X \setminus U_n) \times \mathbf{I}))^* \cap (\text{cl}(U_n) \times [r, 1])^* \cap \downarrow C_F(\alpha X)),$$

that is, $K_1^* \cap \downarrow C_F(\alpha X)$ is an intersection of two elements of \mathcal{S} .

As a conclusion, \mathcal{S} is a subbase for $\downarrow C_F(\alpha X)$. Therefore, $\downarrow C_F(\alpha X)$ is first-countable. Moreover, $\downarrow C_F(\alpha X)$ is second-countable if $\downarrow C_F(X)$ is second-countable. Hence (2) and (3) hold. \square

Lemma 4. *We consider the following statements.*

- (a) $\downarrow C_F(X)$ is first-countable.
- (b) $\downarrow C_F(X)$ has a countable neighborhood base at $\downarrow \mathbf{1}$.
- (c) There exists a countable family \mathcal{U} of nonempty open sets in X such that every nonempty open set in X includes an element of \mathcal{U} , that is, \mathcal{U} is a countable π -base for X .
- (d) $\downarrow C_F(X)$ is separable.

Then the implications (a) \Rightarrow (b) \Rightarrow (c) \Rightarrow (d) hold.

Furthermore, when X is compact, the implication (c) \Rightarrow (a) holds and hence (a), (b) and (c) are equivalent.

PROOF: The implication (a) \Rightarrow (b) is trivial.

(b) \Rightarrow (c): We may assume that

$$\{(G_1^n)^- \cap (G_2^n)^- \cap \dots \cap (G_{k(n)}^n)^- \cap \downarrow C(X) : n = 1, 2, \dots\}$$

is a countable neighborhood base at $\downarrow \mathbf{1}$ in $\downarrow C_F(X)$. Let

$$\mathcal{U} = \{p(G_i^n) : i = 1, 2, \dots, k(n), n = 1, 2, \dots\}.$$

Then \mathcal{U} is a countable family of nonempty open sets in X . We show that every nonempty open set U in X includes an element of \mathcal{U} . Take $f \in C(X)$ such that $f(X \setminus U) \subset \{1\}$ and $f(x_0) = 0$ for some point $x_0 \in U$. Because $\downarrow C_F(X)$ is T_1 by Lemma 1(1), $\downarrow f \notin \bigcap_{i=1}^{k(n)} (G_i^n)^-$ for some n , hence $\downarrow f \notin (G_i^n)^-$ for some $i \leq k(n)$. Then $\downarrow f \cap G_i^n = \emptyset$. Since $f(X \setminus U) \subset \{1\}$, we have $U \supset p(G_i^n)$, as required.

(c) \Rightarrow (d): Let \mathcal{U} be a countable π -base for X . For every $U \in \mathcal{U}$ and $r \in \mathbb{Q} \cap (0, 1]$, we can take a continuous map $f_{(U,r)} : X \rightarrow [0, r]$ such that $f_{(U,r)}(X \setminus U) \subset \{0\}$ and $f_{(U,r)}(x) = r$ for some $x \in U$. Let

$$D = \{\max\{f_{(U,r)} : U \in \mathcal{F}, r \in F\} : \mathcal{F} \text{ and } F \text{ are finite subsets of } \mathcal{U} \text{ and } \mathbb{Q} \cap (0, 1], \text{ resp.}\}.$$

Then $\downarrow D = \{\downarrow f : f \in D\}$ is a countable subset of $\downarrow C(X)$. It remains to verify that $\downarrow D$ is dense in $\downarrow C_F(X)$. Let $f \in C(X)$, K be compact in $X \times (0, 1]$ and G_i ,

$i \leq k$, open in $X \times (0, 1]$, such that

$$\downarrow f \in G_1^- \cap G_2^- \cap \dots \cap G_k^- \cap K^* \cap \downarrow C(X).$$

We have $x_1, \dots, x_k \in X$ such that $\{x_i\} \times [0, f(x_i)] \cap G_i \neq \emptyset$ for each $i \leq k$. Because $\{x_i\} \times [0, f(x_i)] \cap K = \emptyset$, we have an open neighborhood W_i of x_i in X and $s_i < t_i$ such that $W_i \times (s_i, t_i) \subset G_i$ and $W_i \times [0, t_i] \cap K = \emptyset$. Thus, by (c), choose $r_i \in \mathbb{Q} \cap (s_i, t_i)$ and $U_i \in \mathcal{U}$ such that $U_i \subset W_i$. Then $\downarrow f_{(U_i, r_i)} \in G_i^- \cap K^*$ and hence

$$\downarrow \max\{f_{(U_i, r_i)} : i \leq k\} \in \downarrow D \cap G_1^- \cap G_2^- \cap \dots \cap G_k^- \cap K^*.$$

Now, we show (c) \Rightarrow (a) under the assumption that X is compact. Let \mathcal{U} be a countable π -base of X . Then, $X \times \mathbf{I}$ has the following countable π -base:

$$\mathcal{G} = \{U \times (s, t) : U \in \mathcal{U}, s < t \in \mathbb{Q} \cap (0, 1)\}.$$

For every $f \in C(X)$ and $n = 1, 2, \dots$, let

$$\mathcal{G}(f) = \{G \in \mathcal{G} : \downarrow f \in G^-\}, \quad K_n(f) = \{(x, t) \in X \times \mathbf{I} : t \geq f(x) + n^{-1}\}.$$

For every open set H in $X \times (0, 1]$ with $H^- \ni \downarrow f$, there exists $x_0 \in X$ such that $\{x_0\} \times [0, f(x_0)] \cap H \neq \emptyset$. Since $f(x_0) > 0$, we can find an open neighborhood V of x_0 in X and $s < t \in \mathbb{Q} \cap (0, 1)$ such that $s < f(x_0)$, $V \times (s, t) \subset H$ and $s < f(x)$ for every $x \in V$. Since \mathcal{U} is a π -base for X , V contains some $U \in \mathcal{U}$. Then we have $G = U \times (s, t) \in \mathcal{G}$ and $\downarrow f \in G^- \subset H^-$. Moreover, for every compact set K in $X \times \mathbf{I}$ with $K^* \ni \downarrow f$, by the compactness of X , there exists n such that $K_n(f) \supset K$ and hence $\downarrow f \in K_n(f)^* \subset K^*$. Therefore,

$$\{G_1^- \cap \dots \cap G_k^- \cap K_n(f)^* \cap \downarrow C(X) : G_i \in \mathcal{G}(f) \text{ for } i \leq k, k, n = 1, 2, \dots\}$$

is a countable neighborhood base at $\downarrow f$ in $\downarrow C_F(X)$. □

As a consequence of Lemma 4, we have the equivalence between (a) and (b) in Theorem 1, that is,

Proposition 2. *The space $\downarrow C_F(X)$ is metrizable if and only if it is separable metrizable.* □

We need the following two lemmas which were proved in [8], [9], respectively.

Lemma 5. *If V is open in X such that $\text{cl} V$ is compact, then the restriction $\phi : \downarrow C_F(X) \rightarrow \downarrow C_F(\text{cl} V)$ defined by $\phi(\downarrow f) = \downarrow f|_{\text{cl} V}$ is a continuous open surjection.* □

Lemma 6. *If X is compact and $\downarrow C_F(X) = \downarrow C_V(X)$ is second-countable, then X is metrizable.* □

3. Proofs of main results

In this section, we show our main results.

PROOF OF THEOREM 1: The equivalence between (a) and (b) is Proposition 2. If X is first-countable, then X is locally compact by Proposition 1(1). Using Proposition 1(3), the condition (b) implies that $\downarrow C(\alpha X)$ is second-countable. It follows from Lemma 6 that αX is metrizable. Hence the condition (c) holds. That is, the implication (b) \Rightarrow (c) holds under the assumption that X is first-countable. The condition (c) implies that $\text{Cld}_F(X \times \mathbf{I})$ is metrizable ([1, 5.1.5 Theorem]), hence so is $\downarrow C_F(X)$, i.e., (b) holds. Therefore, the implication (c) \Rightarrow (b) holds. \square

PROOF OF THEOREM 2: We may think that every Y_s is a subspace of Y . Define $\phi : C(Y) \rightarrow \prod_{s \in S} C(Y_s)$ by

$$\phi(f) = (f|_{Y_s})_{s \in S} \text{ for each } f \in C(Y).$$

Evidently, ϕ is an injection and its image is

$$\phi(C(Y)) = \left\{ g \in \prod_{s \in S} C(Y_s) : g(s)(a_s) = g(s')(a_{s'}) \text{ for } s, s' \in S \right\}.$$

Now we show that $\downarrow \phi : \downarrow C_F(Y) \rightarrow \prod_{s \in S} \downarrow C_F(Y_s)$ is an embedding. Let $p_s : \prod_{s \in S} \downarrow C_F(Y_s) \rightarrow \downarrow C_F(Y_s)$ be the projection.

To show the continuity of $\downarrow \phi$, it is sufficient to verify that $p_s \circ \downarrow \phi$ is continuous for every $s \in S$. For every open set G in $Y_s \times (0, 1]$, $G \setminus (\{a_s\} \times \mathbf{I})$ is open in $Y \times (0, 1]$. Since a_s is a non-isolated point in Y_s ,

$$(p_s \circ \downarrow \phi)^{-1}(G^- \cap \downarrow C(Y_s)) = (G \setminus (\{a_s\} \times \mathbf{I}))^- \cap \downarrow C(Y).$$

For each compact set K in $Y_s \times (0, 1]$,

$$(p_s \circ \downarrow \phi)^{-1}(K^* \cap \downarrow C(Y_s)) = K^* \cap \downarrow C(Y).$$

Hence, $p_s \circ \downarrow \phi : \downarrow C_F(Y) \rightarrow \downarrow C_F(Y_s)$ is continuous for every $s \in S$.

Moreover, for every open set H in $Y \times (0, 1]$, if $\downarrow f \in H^- \cap \downarrow C_F(Y)$, then there exists $s \in S$ such that $\downarrow f|_{Y_s} \in (H \cap (Y_s \times \mathbf{I}))^-$. Hence

$$\downarrow \phi(H^- \cap \downarrow C_F(Y)) = \bigcup_{s \in S} \left((H \cap (Y_s \times \mathbf{I}))^- \times \prod_{t \in S \setminus \{s\}} \downarrow C(Y_t) \right) \cap \downarrow \phi(\downarrow C(Y)).$$

It shows that $\downarrow \phi(H^- \cap \downarrow C_F(Y))$ is open in $\downarrow \phi(\downarrow C_F(Y))$. For every compact set K in $Y \times (0, 1]$, there exists a finite subset S_0 of S such that $K \subset \bigcup_{s \in S_0} Y_s \times (0, 1]$. Then $K \cap Y_s \times (0, 1]$ is compact for every $s \in S_0$ and

$$\downarrow \phi(K^* \cap \downarrow C(Y)) = \left(\prod_{s \in S_0} (K \cap Y_s \times (0, 1])^* \times \prod_{s \in S \setminus S_0} \downarrow C(Y_s) \right) \cap \downarrow \phi(\downarrow C(Y)).$$

It follows that $\downarrow\phi(K^* \cap \downarrow C(Y))$ is open in $\downarrow\phi(\downarrow C_F(Y))$. Since ϕ is one-to-one, we have that $\downarrow\phi$ maps every open set in $\downarrow C_F(Y)$ to an open set in $\downarrow\phi(\downarrow C_F(Y))$.

Therefore, $\downarrow\phi : \downarrow C_F(Y) \rightarrow \prod_{s \in S} \downarrow C_F(Y_s)$ is an embedding. \square

Remark 1. Even for a set S of two points, if a_s is an isolated point in Y_s for some s , the map $\downarrow\phi$ defined in the above proof needs not be continuous. For example, let $Y_1 = \{1\} \times (\{0\} \cup [1, 2]), Y_2 = \{2\} \times \mathbf{I}$ as subspaces of \mathbb{R}^2 . If we think that $a_1 = (1, 0), a_2 = (2, 0)$, then $p_1 \circ \downarrow\phi : \downarrow C(Y) \rightarrow \downarrow C(Y_1)$ is not continuous. In fact, choose $f_n \in C(Y)$ such that $f_n(2, 0) = f_n(1, 0) = 0$ and $f_n(x) = 1$ for every $x \in Y \setminus (\{2\} \times [0, n^{-1}])$. Then $\downarrow f_n \rightarrow \downarrow \perp$ but $(p_1 \circ \downarrow\phi)(\downarrow f_n) \not\rightarrow (p_1 \circ \downarrow\phi)(\downarrow \perp)$.

PROOF OF COROLLARY 1: Let $\{Y_n : n = 1, 2, \dots\}$ be a family of pairwise disjoint locally compact separable metrizable spaces Y_n with a non-isolated point a_n . Then, by Theorems 1 and 2, the space Y defined in Theorem 2 is as required. \square

PROOF OF THEOREM 3: Let $\beta\omega$ be the Čech-Stone compactification of the discrete space ω of non-negative integers and $q \in \beta\omega \setminus \omega$. Then the subspace $X = \omega \cup \{q\}$ of $\beta\omega$ satisfies the conditions in Theorem 3. By Lemma 1(2), $\downarrow C_F(X)$ is Hausdorff.

Before showing that $\downarrow C_F(X)$ is second-countable but not regular, we verify that every compact subset of X is finite. In fact, let C be an infinite compact subset of X . Then $q \in C$. Write $C = A \cup B \cup \{q\}$ such that A and B are disjoint infinite subsets of ω . Define a continuous map $f : \omega \rightarrow \{0, 1\}$ as $f^{-1}(0) = A$. Then there exists a continuous extension $\bar{f} : X \rightarrow \{0, 1\}$ since X is a subspace of $\beta\omega$. If $\bar{f}(q) = 0$, then B is closed in X and hence is compact. But it is impossible since B is infinite discrete. If $\bar{f}(q) = 1$, then A is closed in X and hence is compact. It is also impossible since A is also infinite discrete.

Now, we define a product space $Y = \prod_{x \in X} \mathbf{I}_x$, where \mathbf{I}_x is a copy of the unit interval $[0, 1]$ with the usual topology for $x \in \omega$ and \mathbf{I}_q is $[0, 1]$ with the topology generated by $\{[0, r) : r \in [0, 1] \cap \mathbb{Q}\} \cup \{[0, 1]\}$. Then Y is second-countable. We may regard $\downarrow C(X) \subset Y$ by identifying $\downarrow f$ with $(f(x))_{x \in X}$ for every $f \in C(X)$. To show that $\downarrow C_F(X)$ is second-countable, it suffices to verify that $\downarrow C_F(X)$ is the subspace of the space Y . It is easy to see that for each $x \in X$, the map $p_x : \downarrow C_F(Y) \rightarrow \mathbf{I}_x$ defined by $p_x(\downarrow f) = f(x)$ is continuous. Hence the subspace topology is coarser than the Fell topology on $\downarrow C(X)$. Conversely, take a compact set $K \subset X \times (0, 1]$ and $f \in C(X)$. Then $p(K)$ is compact in X . Then $p(K)$ is a finite set in X and $\downarrow f \cap K = \emptyset$ if and only if $f(x) < m(x) = \min\{s : (x, s) \in K\}$ for every $x \in p(K)$. Hence we can identify

$$K^* \cap \downarrow C(X) = \left(\prod_{x \in p(K)} [0, m_x) \times \prod_{x \in X \setminus p(K)} \mathbf{I}_x \right) \cap \downarrow C(X)$$

is open in the subspace topology of Y . For every open set G in $X \times (0, 1]$ and

$f \in C(X)$, $\downarrow f \cap G \neq \emptyset$ if and only if $\downarrow f \cap G \setminus (\{q\} \times \mathbf{I}) \neq \emptyset$ if and only if $f(n) > s_n$ for some $n \in p(G) \cap \omega$, where $s_n = \inf\{s : (n, s) \in G\}$. Hence

$$G^- \cap \downarrow C(X) = \left(\bigcup_{n \in p(G) \cap \omega} p_n^{-1}(s_n, 1] \right) \cap \downarrow C(X),$$

where $p_n : Y \rightarrow \mathbf{I}_n$ is the projection, is open in the subspace topology of Y . Therefore, $\downarrow C_F(X)$ is the subspace of Y .

To show that $\downarrow C_F(X)$ is not regular, we consider an open neighborhood $\mathcal{U} = (\{q\} \times [\frac{1}{2}, 1])^* \cap \downarrow C(X)$ of $\downarrow 0$. For every compact set K in $X \times (0, 1]$, $p(K)$ is finite. Define $f \in C(X)$ such that $f^{-1}(0) = p(K) \cap \omega$ and $f^{-1}(1) = X \setminus (p(K) \cap \omega)$. Then $\downarrow f \in \text{cl}_{\downarrow C_F(X)}(K^* \cap \downarrow C_F(X)) \setminus \mathcal{U}$. In fact, every neighborhood of $\downarrow f$ in $\downarrow C_F(X)$ contains the following neighborhood of $\downarrow f$:

$$\mathcal{G} = G_1^- \cap \dots \cap G_k^- \cap G^- \cap C^* \cap \downarrow C_F(X),$$

where $G_i = \{n_i\} \times (s_i, t_i)$ for $1 \leq i \leq k$ and $G = (A \cup \{q\}) \times (s, t)$ are open and C is compact in $X \times (0, 1]$. Then A is an infinite subset of ω and hence we may choose $n_0 \in A \setminus p(K \cup C)$. Now, define $g \in C(X)$ as

$$g(x) = \begin{cases} 0 & \text{if } x \in A \cup \{q\} \setminus \{n_i : 0 \leq i \leq k\}; \\ 1 & \text{if } x = n_0; \\ f(x) & \text{otherwise.} \end{cases}$$

Then it is easy to verify that $\downarrow g \in \mathcal{G} \cap K^*$. This shows that $\downarrow f \in \text{cl}_{\downarrow C_F(X)}(K^* \cap \downarrow C_F(X))$. Because $f(q) = 1$, we have $\downarrow f \notin \mathcal{U}$. Hence, $\text{cl}_{\downarrow C_F(X)}(K^* \cap \downarrow C_F(X)) \not\subset \mathcal{U}$ for any compact K in $X \times (0, 1]$. Note that the family of all of such $K^* \cap \downarrow C_F(X)$ is a neighborhood base at $\downarrow 0$ in $\downarrow C_F(X)$. Therefore, $\downarrow C_F(X)$ is not regular. \square

PROOF OF THEOREM 4: Choose a compact Hausdorff non-metrizable space X satisfying (c) in Lemma 4, for example, $\beta\omega$ or Helly space (see [2, Problem 5.M]). Then, by Lemma 4, $\downarrow C_F(X)$ is separable and first-countable. By [3] (cf. [1, 5.1.2 Proposition]), $\text{Cld}_F(X \times \mathbf{I}) = \text{Cld}_V(X \times \mathbf{I})$ is Tychonoff and hence so is $\downarrow C_F(X)$. Since X is compact and non-metrizable, $\downarrow C_F(X)$ is not second-countable because of Lemma 6. According to Proposition 2, if $\downarrow C_F(X)$ is metrizable, then $\downarrow C_F(X)$ is separable metrizable, hence second-countable. Therefore, $\downarrow C_F(X)$ is not metrizable. \square

PROOF OF THEOREM 5: Assume that $\downarrow C_F(X)$ is metrizable, which means that $\downarrow C_F(X)$ is separable metrizable by Proposition 2. Then $\downarrow C_F(X)$ is second-countable. By Lemma 1(2), there exists a dense open set U in X such that U is locally compact. To complete the proof, it remains to verify that U is separable metrizable. By Lemma 2, there exists a countable family $\mathcal{C} = \{C_1, C_2, \dots\}$ of compact sets in X such that every compact set in X is contained in some C_n . For each n , let $U_n = \text{int}(U \cap C_n)$. Then, $\text{cl}U_n$ is compact because $\text{cl}U_n \subset C_n$. By

Lemma 5, there exists a continuous open surjection from $\downarrow C_F(X)$ onto $\downarrow C_F(\text{cl } U_n)$. Therefore, $\downarrow C_F(\text{cl } U_n)$ is second-countable, hence $\text{cl } U_n$ is compact and metrizable by Lemma 6. Thus every U_n is also separable metrizable, hence it is second-countable. Moreover, for every $x \in U$, there exists an open set V such that $x \in V$, $\text{cl } V$ is compact and $\text{cl } V \subset U$. Hence there exists n such that $\text{cl } V \subset C_n$. Then, $x \in V \subset \text{int}(U \cap C_n) = U_n$. It follows that $U = \bigcup_{n=1}^{\infty} U_n$. Therefore, U is second-countable, hence it is separable metrizable.

As mentioned in proof of Theorem 4, $\beta\omega$ is a compact space and $\downarrow C_F(\beta\omega)$ is not metrizable but ω is a dense, locally compact, open and separable metrizable subspace of $\beta\omega$. Namely, the converse is not true. \square

Remark 2. The referee pointed out that McCoy and Ntantu [11] obtained analogous results in 1992. For example, Theorem 4.12 in [11] is similar to our Theorem 1. Our Theorem 3 for $\downarrow C_F(X, \mathbf{I})$ is true for $\uparrow C_F(X, \mathbb{R})$ using Theorems 3.5, 3.7, 4.11 and Example 3.3 in [11], where $\uparrow C_F(X, \mathbb{R})$ is the subspace of $\text{Cld}_F(X \times \mathbb{R})$ consisting of the epigraphs

$$\uparrow f = \{(x, s) \in X \times \mathbb{R} : f(x) \leq s\} \in \text{Cld}(X \times \mathbb{R}),$$

of all $f \in C(X, \mathbb{R})$. However our arguments are quite different from their arguments in [11].

Acknowledgment. The author is so grateful to Professors K. Sakai and K. Mine for their guidance of revising the original manuscript. He also appreciates the referee's comments on the results of McCoy and Ntantu in [11].

REFERENCES

- [1] Beer G., *Topologies on Closed and Closed Convex Sets*, MIA 268, Kluwer Acad. Publ., Dordrecht, 1993.
- [2] Kelly J.L., *General Topology*, GTM 27, Springer, New York; Reprint of the 1955 ed. published by Van Nostrand, 1955.
- [3] Michael E., *Topologies on spaces of subsets*, Trans. Amer. Math. Soc. **71** (1951), 152–182.
- [4] Yang Z., *The hyperspace of the regions below of continuous maps is homeomorphic to c_0* , Topology Appl. **153** (2006), 2908–2921.
- [5] Yang Z., Fan L., *The hyperspace of the regions below of continuous maps from the converging sequence*, Northeast Math. J. **22** (2006), 45–54.
- [6] Yang Z., Wu N., *The hyperspace of the regions below of continuous maps from S^*S to I* , Questions Answers Gen. Topology **26** (2008), 29–39.
- [7] Yang Z., Wu N., *A topological position of the set of continuous maps in the set of upper semicontinuous maps*, Science in China, Ser. A: Math. **52** (2009), 1815–1828.
- [8] Yang Z., Zhang B., *The hyperspace of the regions below continuous maps with the Fell topology is homeomorphic to c_0* , Acta Math. Sinica, English Ser. **28** (2012), 57–66.
- [9] Yang Z., Zhou X., *A pair of spaces of upper semi-continuous maps and continuous maps*, Topology Appl. **154** (2007), 1737–1747.
- [10] Zhang Y., Yang Z., *Hyperspaces of the regions below of upper semi-continuous maps on non-compact metric spaces*, Advances in Math. in China **39** (2010), 352–360 (Chinese).

- [11] McCoy R.A., Ntanyu I., *Properties $C(X)$ with the epi-topology*, Bollettion U.M.I. (7)**6-B**(1992), 507–532.

DOCTORAL PROGRAM IN MATHEMATICS, GRADUATE SCHOOL OF PURE AND APPLIED SCIENCES, UNIVERSITY OF TSUKUBA, TSUKUBA, 305-8571, JAPAN

E-mail: hongsejulebu@sina.com

(Received April 3, 2012, revised May 8, 2012)