

Tao Xu; Heguo Liu

On groups of automorphisms of nilpotent p -groups of finite rank

Czechoslovak Mathematical Journal, Vol. 70 (2020), No. 4, 1161–1165

Persistent URL: <http://dml.cz/dmlcz/148419>

Terms of use:

© Institute of Mathematics AS CR, 2020

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

ON GROUPS OF AUTOMORPHISMS
OF NILPOTENT p -GROUPS OF FINITE RANK

TAO XU, Handan, HEGUO LIU, Wuhan

Received June 8, 2019. Published online April 22, 2020.

Abstract. Let α and β be automorphisms of a nilpotent p -group G of finite rank. Suppose that $\langle (\alpha\beta(g))(\beta\alpha(g))^{-1} : g \in G \rangle$ is a finite cyclic subgroup of G , then, exclusively, one of the following statements holds for G and Γ , where Γ is the group generated by α and β .

- (i) G is finite, then Γ is an extension of a p -group by an abelian group.
- (ii) G is infinite, then Γ is soluble and abelian-by-finite.

Keywords: automorphism; nilpotent group; finite rank

MSC 2020: 20F18, 20F28

1. INTRODUCTION

A well-known result of Hirsch (see [7]) asserts that a polycyclic group G is nilpotent if and only if all finite images of G are nilpotent. Hirsch's result has been extended to wider classes of groups: finitely generated hyper-abelian groups (see [5]), residually finite soluble minimax groups (see [4]) and linear groups over a finitely generated integral domain (see [8]).

In particular, Dardano, Eick and Menth in [1] proved that if the automorphism subgroup Γ of a residually finite soluble minimax group G induces a nilpotent group of automorphisms on every finite characteristic quotient of G , then Γ is nilpotent, which improves Hirsch's theorem.

The research has been supported by the National Natural Science Foundation of China (11801129, 11771129), the Hebei Provincial Natural Science Foundation (A2019402211), the Program for Young Top Talent of Higher Learning Institutions of Hebei (BJ2018025), the Excellent Young and Middle-Aged Innovative Team Program of Hubei (T201601), the New Century High-Level Talents Foundation of Hubei and the Foundation of Handan (1723208068-5).

Recall that a group has finite rank r if every finitely generated subgroup can be generated by r elements and if r is the least positive integer with this property. It is natural to ask what the structure of the group generated by automorphisms is if the automorphisms of nilpotent groups of finite rank have some given property.

Liu and Zhang in [3] determined the structure of the group Γ generated by two p -automorphisms α and β acting on a finite rank nilpotent p -group G such that the subgroup I generated by the set $\{(\alpha\beta(g))(\beta\alpha(g))^{-1} : g \in G\}$ is finite cyclic. In this paper the same question is treated under the hypothesis that both α and β are automorphisms.

Theorem 1.1. *Let α and β be automorphisms of a nilpotent p -group G of finite rank. Assume that $I = \langle (\alpha\beta(g))(\beta\alpha(g))^{-1} : g \in G \rangle$ is a finite cyclic subgroup of G , then, exclusively, one of the following holds for G and Γ , where Γ is the group generated by α and β .*

- (i) G is finite, then Γ is an extension of a p -group by an abelian group.
- (ii) G is infinite, then Γ is soluble and abelian-by-finite.

Let G be a group. We shall write $\text{Aut } G$ for the automorphism group of G and ζG for its centre. Assume that $\alpha \in \text{Aut } G$ and H is an α -invariant subgroup of G . Denote the order of α by $|\alpha|$ and the restriction of α to H by $\alpha|_H$. Other notations and terminologies used throughout this paper are consistent with those in [6].

2. PROOF OF THEOREM

Lemma 2.1. *Let α and β be automorphisms of a finite vector space A . If $\text{Im}(\alpha\beta - \beta\alpha)$ is a cyclic group, then the group generated by α and β is an extension of a p -group by an abelian group.*

Proof. Suppose that $|\alpha| = m$, $|\beta| = n$, and A is a r -dimensional vector space over the finite field K . We may assume that F is an algebraic closure of K . It is easy to see that $\alpha, \beta \in \text{GL}(r, K) \leq \text{GL}(r, F)$. Since $\text{Im}(\alpha\beta - \beta\alpha)$ is finite cyclic, $\text{rank}(\alpha\beta - \beta\alpha) \leq 1$. By [2], α and β can be put in upper triangular form simultaneously. In other words, there exists $\gamma \in \text{GL}(r, F)$ such that

$$\gamma^{-1}\alpha\gamma = \begin{pmatrix} a_1 & & & \\ & a_2 & * & \\ & & \ddots & \\ 0 & & & a_r \end{pmatrix}, \quad \gamma^{-1}\beta\gamma = \begin{pmatrix} b_1 & & & \\ & b_2 & * & \\ & & \ddots & \\ 0 & & & b_r \end{pmatrix},$$

where $a_i^m = 1$, $b_i^n = 1$ for all $i \in \{1, 2, \dots, r\}$.

Since F is a finite field, the group of unitriangular matrices $U(r, F)$ is a finite p -group. Noticing

$$T(r, F) = \underbrace{(F^* \times F^* \times \dots \times F^*)}_r \rtimes U(r, F),$$

where $T(r, F)$ is a group of upper triangular matrices over F , we have that the group generated by $\gamma^{-1}\alpha\gamma$ and $\gamma^{-1}\beta\gamma$ is an extension of a finite p -group by an abelian group. Hence the group generated by α and β is an extension of a finite p -group by an abelian group. \square

Lemma 2.2. *Let G and Γ be as in Theorem 1.1 and suppose that N is a normal Γ -invariant proper subgroup with G/N a finite vector space, then N and $\Gamma|_N = \Gamma/C_\Gamma(N)$ satisfy the premises of Theorem 1.1. If $\Gamma/C_\Gamma(N)$ satisfies the conclusion of (i) or (ii), respectively, then correspondingly so does Γ .*

Proof. The epimorphism, say φ , from Γ to $\Gamma/C_\Gamma(N)$ is given by sending $\gamma \in \Gamma$ to the restriction on N . Clearly $\varphi(\alpha)$ and $\varphi(\beta)$ satisfy the premise of Theorem 1.1.

Suppose first that N and $\Gamma/C_\Gamma(N)$ satisfy (i). Then G is finite and, by assumption, $\Gamma/C_\Gamma(N)$ contains a normal p -subgroup $P/C_\Gamma(N)$ such that Γ/P is abelian. By [6], Theorem 5.3.3, $C_\Gamma(N/\text{Frat } N)$ is a p -group. Since $C_\Gamma(N) \leq C_\Gamma(N/\text{Frat } N)$, $C_\Gamma(N)$ is a p -group. It follows that Γ is an extension of a p -group by an abelian group. Thus (i) holds for Γ .

Suppose next that N and $\Gamma/C_\Gamma(N)$ satisfy (ii). Then, by assumption, $\Gamma/C_\Gamma(N)$ is soluble and abelian-by-finite. Thus there is a normal series of Γ

$$\Gamma \geq A \geq C_\Gamma(N) \geq 1,$$

where $A/C_\Gamma(N)$ is abelian and Γ/A is finite and soluble. Since

$$C_\Gamma(N) \leq C_\Gamma(N/\text{Frat } N)$$

and, as already observed during the proof of (i), the latter group is a finite p -group, it follows that Γ is soluble. Note that $A/C_\Gamma(N)$ is abelian and $C_\Gamma(N)$ is a finite p -group, and we have that the commutator subgroup of A must be finite. As A has finite index in the finitely generated group Γ we conclude that A is finitely generated. By [6], Exercise 7, ζA has a finite index in A . Since both Γ/A and $A/\zeta A$ are finite, Γ is an extension of the abelian group ζA by the finite soluble group $\Gamma/\zeta A$, establishing (ii) for Γ . \square

Lemma 2.3. *Let α and β be automorphisms of an abelian divisible p -group A of finite rank. If $\text{Im}(\alpha\beta - \beta\alpha)$ is a finite cyclic group, then $\Gamma = \langle \alpha, \beta \rangle$ is abelian.*

Proof. The endomorphism $\alpha\beta - \beta\alpha$ maps the divisible group A onto a divisible subgroup of A and, as A is infinite, its image cannot be finite. It follows that α and β commute. Hence $\Gamma = \langle \alpha, \beta \rangle$ is abelian. \square

Proof of Theorem 1.1. Since G is a finite rank p -group, it satisfies the minimal condition on normal subgroups. By [6], Theorem 5.4.23, G is a finite extension of a divisible subgroup D . Thus G/D is a finite p -group and G is finite if and only if D is trivial. We need to consider the cases (i) G being finite and (ii) G being infinite.

(i) If G is elementary abelian, the result follows from Lemma 2.1. Otherwise, by Lemma 2.2, we may replace G by any maximal Γ -invariant subgroup N with G/N elementary abelian. Hence (i) holds for G and Γ if and only if (i) holds for N and $\Gamma/C_\Gamma(N)$. Now apply induction on $|G|$ until N itself becomes elementary abelian.

(ii) Now D is nontrivial. If $G = D$, then Lemma 2.3 implies the result. Otherwise G/D is a nontrivial finite p -group. Then G/D has a normal Γ -invariant subgroup N containing D such that G/N is elementary abelian. Using Lemma 2.2, we may replace G by N . Now apply induction on $|G/D|$ until N becomes identical to D . \square

3. EXAMPLE

The following example validates the correctness of Lemma 2.1 and also illustrates that in Lemma 2.1 the group generated by α and β is not necessarily nilpotent.

Example 3.1. Let A be a 2-dimensional vector space over \mathbb{Z}_3 , then the automorphism group of A is $\text{GL}(2, \mathbb{Z}_3)$. In $\text{GL}(2, \mathbb{Z}_3)$, we take

$$\alpha = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad \beta = \begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}.$$

Clearly $|\alpha| = 3$, $|\beta| = 2$, and

$$\alpha\beta - \beta\alpha = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad \beta^{-1}\alpha\beta = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix} = \alpha^{-1}.$$

It is easy to check that $\text{Im}(\alpha\beta - \beta\alpha)$ is a group of order 3 and $\langle \alpha, \beta \rangle$ is a symmetric group of degree 3.

Acknowledgements. We thank the reviewers very much for their careful work and valuable suggestions which simplify the proof of the main results.

References

- [1] *U. Dardano, B. Eick, M. Menth*: On groups of automorphisms of residually finite groups. *J. Algebra* 231 (2000), 561–573. [zbl](#) [MR](#) [doi](#)
- [2] *R. M. Guralnick*: A note on pairs of matrices with rank one commutator. *Linear Multi-linear Algebra* 8 (1979), 97–99. [zbl](#) [MR](#) [doi](#)
- [3] *H. G. Liu, J. P. Zhang*: On p -automorphisms of a nilpotent p -group with finite rank. *Acta Math. Sin., Chin. Ser.* 50 (2007), 11–16. (In Chinese.) [zbl](#) [MR](#)
- [4] *D. J. S. Robinson*: Residual properties of some classes of infinite soluble groups. *Proc. Lond. Math. Soc., III. Ser.* 18 (1968), 495–520. [zbl](#) [MR](#) [doi](#)
- [5] *D. J. S. Robinson*: Finiteness Conditions and Generalized Soluble Groups. Part 2. *Ergebnisse der Mathematik und ihrer Grenzgebiete* 63, Springer, Berlin, 1972. [zbl](#) [MR](#) [doi](#)
- [6] *D. J. S. Robinson*: A Course in the Theory of Groups. *Graduate Texts in Mathematics* 80, Springer, New York, 1982. [zbl](#) [MR](#) [doi](#)
- [7] *D. Segal*: Polycyclic Groups. *Cambridge Tracts in Mathematics* 82, Cambridge University Press, Cambridge, 1983. [zbl](#) [MR](#) [doi](#)
- [8] *B. A. F. Wehrfritz*: Infinite Linear Groups. An Account of the Group-Theoretic Properties of Infinite Groups of Matrices. *Ergebnisse der Mathematik und ihrer Grenzgebiete* 76, Springer, Berlin, 1973. [zbl](#) [MR](#) [doi](#)

Authors' addresses: T a o X u, Department of Science, Hebei University of Engineering, No. 19 Taiji Road, Handan, 056038, Hebei, P. R. China, e-mail: gtxutao@163.com; H e g u o L i u, Department of Mathematics, Hubei University, No. 368 Youyi Road, Wuhan, 430062, Hubei, P. R. China, e-mail: ghliu@hubu.edu.cn.