

Murat Alan

Mersenne numbers as a difference of two Lucas numbers

*Commentationes Mathematicae Universitatis Carolinae*, Vol. 63 (2022), No. 3, 269–276

Persistent URL: <http://dml.cz/dmlcz/151474>

## Terms of use:

© Charles University in Prague, Faculty of Mathematics and Physics, 2022

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

# Mersenne numbers as a difference of two Lucas numbers

MURAT ALAN

*Abstract.* Let  $(L_n)_{n \geq 0}$  be the Lucas sequence. We show that the Diophantine equation  $L_n - L_m = M_k$  has only the nonnegative integer solutions  $(n, m, k) = (2, 0, 1), (3, 1, 2), (3, 2, 1), (4, 3, 2), (5, 3, 3), (6, 2, 4), (6, 5, 3)$  where  $M_k = 2^k - 1$  is the  $k$ th Mersenne number and  $n > m$ .

*Keywords:* Lucas number; Mersenne number; Diophantine equation; linear forms in logarithm

*Classification:* 11B39, 11J86, 11D61

## 1. Introduction

Let  $(F_n)_{n \geq 0}$  and  $(L_n)_{n \geq 0}$  be the Fibonacci and Lucas sequences given by  $F_0 = 0, F_1 = 1, L_0 = 2, L_1 = 1, F_{n+2} = F_{n+1} + F_n$  and  $L_{n+2} = L_{n+1} + L_n$  for  $n \geq 0$ , respectively. Because of their interesting properties, these two sequences have been investigated from various point of view in many papers. Especially over the last decade products, sums and differences of these two sequences have been investigated by a number of authors, see for example [4], [5], [7], [9], [12], [13], [14]. We refer [10] for a detailed information on these sequences.

Recall that Mersenne numbers are the numbers of the form  $2^k - 1$ , where  $k$  is a nonnegative integer. In [2], Mersenne numbers in a generalized Fibonacci numbers are found. In this study, we search the differences of two Lucas numbers which are Mersenne numbers. More precisely the aim of this paper is to prove the following theorem.

**Theorem 1.** *The equation*

$$(1) \quad L_n - L_m = 2^k - 1, \quad n > m,$$

*has only the nonnegative integer solutions  $(n, m, k) \in \{(2, 0, 1), (3, 1, 2), (3, 2, 1), (4, 3, 2), (5, 3, 3), (6, 2, 4), (6, 5, 3)\}$ .*

The proof of the above theorem mainly depends on two results one of them gives us a general lower bound for linear forms in logarithms and the other one

is a version of reduction method given in Baker–Davenport lemma, see [1]. We summarize these two results in the next chapter. It is worth to note that when it is needed we used the software Maple for all calculations and computations.

## 2. Preliminaries

It is well-known that all Lucas numbers  $L_n$  can be written as

$$L_n = \alpha^n + \beta^n, \quad n \geq 0,$$

where

$$\alpha := \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \beta := \frac{1 - \sqrt{5}}{2}$$

are the roots of the equation  $x^2 - x - 1 = 0$ . This is the Binet formula of Lucas numbers. By using the Binet formula, one can see that,

$$(2) \quad \alpha^{n-1} \leq L_n \leq 2\alpha^n$$

holds for all  $n > 0$ .

**Lemma 2.** If  $n \equiv m \pmod{3}$  then  $L_n \equiv L_m \pmod{2}$

PROOF: Assume that  $n \equiv m \pmod{3}$ . Assume also without loss of generality that  $n \geq m$  and let  $n = m + 3k$  for some nonnegative integer  $k$ . Then the proof follows by induction on  $k$ , since  $L_{m+3} = L_{m+2} + L_{m+1} = 2L_{m+1} + L_m \equiv L_m \pmod{2}$ .  $\square$

Let  $\eta$  be an algebraic number of degree  $d$  with minimal polynomial

$$a_0x^d + a_1x^{d-1} + \cdots + a_d = a_0 \prod_{i=1}^d (X - \eta^{(i)}),$$

where the  $a_i$ 's are relatively prime integers with  $a_0 > 0$  and the  $\eta^{(i)}$ 's are conjugates of  $\eta$ . The logarithmic height of  $\eta$  is defined by

$$h(\eta) = \frac{1}{d} \left( \log a_0 + \sum_{i=1}^d \log(\max\{|\eta^{(i)}|, 1\}) \right).$$

In particular, for a rational number  $p/q$  with  $\gcd(p, q) = 1$  and  $q > 0$ ,  $h(p/q) = \log \max\{|p|, q\}$ . The following properties are very useful for calculation of  $h(\eta)$ :

- $h(\eta \pm \gamma) \leq h(\eta) + h(\gamma) + \log 2$ .
- $h(\eta\gamma^{\pm 1}) \leq h(\eta) + h(\gamma)$ .
- $h(\eta^s) = |s|h(\eta)$ .

**Theorem 3** (Matveev’s theorem, [11]). *Assume that  $\gamma_1, \dots, \gamma_t$  are positive real algebraic numbers in a real algebraic number field  $\mathbb{K}$  of degree  $d_{\mathbb{K}}$ ,  $b_1, \dots, b_t$  are rational integers, and*

$$\Lambda := \gamma_1^{b_1} \cdots \gamma_t^{b_t} - 1$$

*is not zero. Then*

$$|\Lambda| > \exp \left( - 1.4 \cdot 30^{t+3} \cdot t^{4.5} \cdot d_{\mathbb{K}}^2 (1 + \log d_{\mathbb{K}}) (1 + \log B) A_1 \cdots A_t \right),$$

where

$$B \geq \max\{|b_1|, \dots, |b_t|\},$$

and

$$A_i \geq \max\{d_{\mathbb{K}} h(\gamma_i), |\log \gamma_i|, 0.16\} \quad \text{for all } i = 1, \dots, t.$$

In [6, Lemma 5 (a)], A. Dujella and A. Pethő give a version of the reduction method based on the Baker–Davenport lemma, see [1]. The following lemma is from [3, Lemma 4] which is a variation of [6, Lemma 5 (a)] and it is useful for reducing some upper bounds on the variables. For a real number  $\theta$ , we put  $\|\theta\| = \min\{|\theta - n| : n \in \mathbb{Z}\}$ , the distance from  $\theta$  to the nearest integer.

**Lemma 4.** Let  $M$  be a positive integer, let  $p/q$  be a convergent of the continued fraction of the irrational  $\gamma$  such that  $q > 6M$ , and let  $A, B, \mu$  be some real numbers with  $A > 0$  and  $B > 1$ . If  $\varepsilon := \|\mu q\| - M\|\gamma q\| > 0$ , then there is no solution to the inequality

$$0 < |u\gamma - v + \mu| < AB^{-w},$$

in positive integers  $u, v$  and  $w$  with

$$u \leq M \quad \text{and} \quad w \geq \frac{\log(Aq/\varepsilon)}{\log B}.$$

### 3. Proof of Theorem 1

PROOF OF THEOREM 1: Assume that the equation (1) holds. A quick computation with Maple shows that the equation (1) has the solutions  $(n, m, k) \in \{(2, 0, 1), (3, 1, 2), (3, 2, 1), (4, 3, 2), (5, 3, 3), (6, 2, 4), (6, 5, 3)\}$  in nonnegative integers in the range  $0 \leq m < n \leq 200$ . If  $n - m = 1$  and  $m \neq 0$  or  $n - m = 2$ , then the equation (1) turns into the equations  $L_{m-1} = 2^k - 1$  and  $L_{m+1} = 2^k - 1$ , respectively. All solutions of these two equations only come from the equalities  $L_1 + 1 = 2$ ,  $L_2 + 1 = 2^2$  and  $L_4 + 1 = 2^8$ , see [4], which corresponds already known solutions  $(n, m, k) \in \{(3, 2, 1), (4, 3, 2), (6, 5, 3)\}$  and  $(n, m, k) \in \{(2, 0, 1), (3, 1, 2),$

$(5, 3, 3)$  of (1). Also, by Lemma 2, the case  $n - m = 3$  is clearly false. So we will take  $n - m \geq 4$  and  $n > 200$ .

From (1), we may get a relation between variables  $n$  and  $k$  for future reference. Indeed from (1) and (2) we have that

$$2^{k-1} \leq 2^k - 1 = L_n - L_m < L_n \leq 2\alpha^n < 2^{n+1},$$

and hence we observe that

$$(3) \quad k < n + 2.$$

Using Binet formula for Lucas sequences we rewrite equation (1) as

$$\begin{aligned} \alpha^n + \beta^n - L_m &= 2^k - 1, \\ \alpha^n - 2^k &= L_m - \beta^n - 1. \end{aligned}$$

So

$$\left| 1 - \frac{2^k}{\alpha^n} \right| < \frac{L_m}{\alpha^n} + \frac{|\beta|^n}{\alpha^n} + \frac{1}{\alpha^n} < \frac{2\alpha^m}{\alpha^n} + \frac{1}{\alpha^{2n}} + \frac{1}{\alpha^n} < \frac{4}{\alpha^{n-m}}.$$

Hence we have that

$$(4) \quad |\Lambda_1| := \left| 1 - \frac{2^k}{\alpha^n} \right| < \frac{4}{\alpha^{n-m}}.$$

Now we may apply Theorem 3 to the left side of the inequality (4) with  $\eta_1 := 2$ ,  $\eta_2 := \alpha$  and  $b_1 := k$ ,  $b_2 := -n$ . Clearly  $\eta_1, \eta_2 \in \mathbb{K} = \mathbb{Q}(\sqrt{5})$ , and hence we take  $d_{\mathbb{K}} = 2$ , the degree of the number field  $\mathbb{K}$ . Since  $\alpha^n \neq 2^k \in \mathbb{Z}$  for  $n > 0$ , we see that  $\Lambda_1$  is nonzero. Further since  $h(\eta_1) = \log 2$  and  $h(\eta_2) = (1/2) \log \alpha$ , we take  $A_1 = 2 \log 2$ ,  $A_2 = \log \alpha$ . By (3) we know that  $k < n + 2$ , so we take

$$B := \max\{|b_i|\} = \max\{k, n\} < n + 2.$$

Now we are ready to apply Theorem 3 to the inequality (4) and we get that

$$\log |\Lambda_1| > -1.4 \times 30^5 \times 2^{4.5} \times 2^2 (1 + \log 2)(1 + \log(n + 2)) 2 \log 2 \cdot \log \alpha,$$

which implies that

$$(5) \quad \log |\Lambda_1| > 3.48 \times 10^9 (1 + \log(n + 2)).$$

On the other hand, from (4), we know that

$$(6) \quad \log |\Lambda_1| < \log 4 - (n - m) \log \alpha.$$

From (5) and (6) we find that

$$(7) \quad n - m < 7.24 \times 10^9 \times (1 + \log(n + 2)).$$

Now we rewrite equation (1) in the Binet formula as

$$\begin{aligned} \alpha^n + \beta^n - \alpha^m - \beta^m &= 2^k - 1, \\ \alpha^n - \alpha^m - 2^k &= -\beta^n + \beta^m - 1, \\ \alpha^n(1 - \alpha^{m-n}) - 2^k &= -\beta^n + \beta^m - 1. \end{aligned}$$

So from the last equation above we write

$$\left| 1 - \frac{2^k}{\alpha^n(1 - \alpha^{m-n})} \right| < \frac{1}{\alpha^n(1 - \alpha^{m-n})} (|\beta|^n + |\beta|^m + 1) < \frac{3}{\alpha^n},$$

where we used the fact that  $|\beta|^n + |\beta|^m + 1 < 2.01$  and  $(1 - \alpha^{m-n})^{-1} < 1.4$  for  $m \geq 0, n \geq 200$  and  $n - m \geq 3$ . So

$$(8) \quad |\Lambda_2| := \left| 1 - \frac{2^k}{\alpha^n(1 - \alpha^{m-n})} \right| < \frac{3}{\alpha^n}.$$

Let  $\eta_1 := 2, \eta_2 := \alpha, \eta_3 := (1 - \alpha^{m-n})^{-1}$  and  $b_1 := k, b_2 := -n, b_3 := 1$ . Since  $\eta_1, \eta_2$  and  $\eta_3$  are elements of the quadratic number field  $\mathbb{K} = \mathbb{Q}(\sqrt{5})$ , we take  $d_{\mathbb{K}} = 2$ , the degree of the number field  $\mathbb{K}$ . Since  $h(\eta_1) = \log 2, h(\eta_2) = (1/2) \log \alpha$  and

$$\begin{aligned} h(\eta_3) &= h((1 - \alpha^{m-n})^{-1}) = h(1 - \alpha^{m-n}) \\ &\leq |m - n|h(\alpha) + \log 2 = \frac{n - m}{2} \log \alpha + \log 2 \end{aligned}$$

we take

$$A_1 = 2 \log 2, \quad A_2 = \log \alpha \quad \text{and} \quad A_3 = (n - m) \log \alpha + 2 \log 2.$$

By (3),  $k < n + 2$ , so we take

$$B := \max\{|b_i|\} = \max\{n, k, 1\} < n + 2.$$

Note that  $\Lambda_2$  is nonzero. Indeed if it were zero then we would have that  $\alpha^n - \alpha^m = 2^k$ . Conjugating in  $\mathbb{K}$ , we get that  $\beta^n - \beta^m = -2^k$  and adding these two equation we get that  $L_n - L_m = 0$ , which is false since  $n > m$ . Thus we apply Theorem 3 to  $\Lambda_2$  given in (8) and we get that

$$(9) \quad \begin{aligned} \log(|\Lambda_2|) &> -1.4 \cdot 30^6 \cdot 3^{4.5} \cdot 2^2(1 + \log 2)(1 + \log(n + 2)) \\ &\quad \times 2 \log 2 \cdot \log \alpha \cdot ((n - m) \log \alpha + 2 \log 2). \end{aligned}$$

On the other hand, from (8), we know that

$$(10) \quad \log|\Lambda_2| < \log 3 - n \log \alpha.$$

Combining (9) and (10) we get that

$$(11) \quad n < 1.35 \cdot 10^{12} \times (1 + \log(n + 2)) \times ((n - m) \log \alpha + 2 \log 2).$$

From the inequality (11) together with (7), it follows that

$$(12) \quad n < 1.8 \cdot 10^{25}.$$

Now we use the theory of continued fractions to reduce the upper bound on  $n - m$ . More precisely we show that  $n - m < 137$ . To this end, assume that  $n - m \geq 137$ . Let

$$(13) \quad \Gamma_1 := k \log 2 - n \log \alpha$$

so that

$$|\Lambda_1| := |\exp(\Gamma_1) - 1| < \frac{4}{\alpha^{n-m}} < \frac{1}{2},$$

and hence we get that

$$|\Gamma_1| < \frac{8}{\alpha^{n-m}}.$$

So from (13), we write

$$\left| \frac{\log \alpha}{\log 2} - \frac{k}{n} \right| < \frac{8}{n \alpha^{n-m} \log 2}.$$

Since  $2.8 \cdot 10^{28} < \alpha^{137} \log 2$ , we have the inequality

$$16n < 16 \cdot 1.8 \cdot 10^{25} < 2.9 \cdot 10^{26} < \alpha^{137} \log 2 \leq \alpha^{n-m} \log 2.$$

Thus we write

$$\left| \frac{\log \alpha}{\log 2} - \frac{k}{n} \right| < \frac{8}{n \alpha^{n-m} \log 2} < \frac{1}{2n^2},$$

which means that  $k/n$  is a convergent of continued fractions of irrational  $\tau := \log \alpha / \log 2$ , say  $p_k/q_k$ . Since  $p_k$  and  $q_k$  are relatively prime integers we get that  $q_k \leq n < 1.8 \times 10^{25}$ . Let  $[a_0, a_1, a_2, a_3, a_4, \dots] = [0, 1, 2, 3, 1, \dots]$  be the continued fraction expansion of  $\tau$ . By Maple, we see that  $k \leq 56$  and  $\max\{a_i\} = 134$  for  $i = 1, 2, \dots, 56$ . Thus, from the well known property of continued fractions [8, Theorem 163], we write

$$\frac{1}{136n^2} < \frac{1}{(a_k + 2)n^2} < \left| \frac{\log \alpha}{\log 2} - \frac{k}{n} \right| < \frac{8}{n \alpha^{n-m} \log 2},$$

which means

$$2.7 \cdot 10^{25} < \frac{\alpha^{137} \log 2}{136 \cdot 8} \leq \frac{\alpha^{n-m} \log 2}{136 \cdot 8} < n,$$

a contradiction since  $n < 1.8 \times 10^{25}$ . So we conclude that  $n - m < 137$ . By substituting this upper bound for  $n - m$  into (11) we get that

$$(14) \quad n < 1.35 \cdot 10^{12} \times (1 + \log(n + 2)) \times (137 \log \alpha + 2 \log 2),$$

which implies that

$$(15) \quad n < 4 \cdot 10^{15}.$$

Let

$$(16) \quad \Gamma_2 := k \log 2 - n \log \alpha - \log(1 - \alpha^{m-n})$$

so that

$$|\Lambda_2| := |\exp \Gamma_2 - 1| < \frac{3}{\alpha^n} < \frac{1}{2}.$$

Then

$$|\Gamma_2| < \frac{6}{\alpha^n}.$$

So, from (16),

$$(17) \quad 0 < \left| k \frac{\log 2}{\log \alpha} - n - \frac{\log(1 - \alpha^{m-n})}{\log \alpha} \right| < \frac{6}{\alpha^n \log \alpha}.$$

Now we take

$$M := 4 \cdot 10^{15} + 2 > n + 2 > k \quad \text{and} \quad \tau := \frac{\log 2}{\log \alpha}.$$

Then, in the continued fraction expansion of  $\tau$ , we take  $q_{50}$ , the denominator of the 50th convergent of  $\tau$ , which exceeds  $6M$ . Now with the help of Maple we calculate

$$\varepsilon_{n-m} := \|\mu_{n-m} q_{50}\| - M \|\tau q_{50}\|$$

for each  $n - m \in \{4, \dots, 137\}$  where

$$\mu_{n-m} := -\frac{\log(1 - \alpha^{m-n})}{\log \alpha}, \quad q_{50} = 9041151586240430787539$$

and we get that

$$0.000012 < \varepsilon_{n-m} \quad \text{for all } n - m \in \{4, \dots, 137\}$$

except for  $n - m = 6$ , since for  $n - m = 6$ ,  $\varepsilon_6 < 0$ . So we need to handle this case separately. But, from Lemma 2, this is not the case. So  $n - m \neq 6$ .



Let  $A := 6/\log \alpha$ ,  $B := \alpha$  and  $\omega := n$ . Thus from Lemma (4) we get that the inequality (17) has no solution for

$$n > \frac{\log(Aq_{50}/0.000012)}{\log B} \geq 133.8.$$

So we get that  $n < 134$  which is a contradiction since  $n > 200$ . This completes the proof.  $\square$

#### REFERENCES

- [1] Baker A., Davenport H., *The equations  $3x^2 - 2 = y^2$  and  $8x^2 - 7 = z^2$* , Quart. J. Math. Oxford Ser. (2) **20** (1969), 129–137.
- [2] Bravo J. J., Gómez C. A., *Mersenne  $k$ -Fibonacci numbers*, Glas. Matemat. Ser. III **51(71)**, (2016), no. 2, 307–319.
- [3] Bravo J. J., Luca F., *On a conjecture about repdigits in  $k$ -generalized Fibonacci sequences*, Publ. Math. Debrecen **82** (2013), no. 3–4, 623–639.
- [4] Bravo J. J., Luca F., *Powers of two as sums of two Lucas numbers*, J. Integer Seq. **17** (2014), no. 8, Article 14.8.3, 12 pages.
- [5] Demirtürk Bitim B., *On the Diophantine equation  $L_n - L_m = 2 \cdot 3^a$* , Period. Math. Hungar. **79** (2019), no. 2, 210–217.
- [6] Dujella A., Pethő A., *A generalization of a theorem of Baker and Davenport*, Quart. J. Math. Oxford Ser. (2) **49** (1998), no. 195, 291–306.
- [7] Erduvan F., Keskin R., *Nonnegative integer solutions of the equation  $F_n - F_m = 5^a$* , Turkish J. Math. **43** (2019), no. 3, 1115–1123.
- [8] Hardy G. H., Wright E. M., *An Introduction to the Theory of Numbers*, The Clarendon Press, Oxford University Press, New York, 1979.
- [9] Kebli S., Kihel O., Larone J., Luca F., *On the nonnegative integer solutions to the equation  $F_n \pm F_m = y^a$* , J. Number Theory **220** (2021), 107–127.
- [10] Koshy T., *Fibonacci and Lucas Numbers with Applications*, Pure and Applied Mathematics (New York), Wiley-Interscience, New York, 2001.
- [11] Matveev E. M., *An explicit lower bound for a homogeneous rational linear form in the logarithms of algebraic numbers. II*, Izv. Ross. Akad. Nauk Ser. Mat. **64** (2000), no. 6, 125–180 (Russian); translation in Izv. Math. **64** (2000), no. 6, 1217–1269.
- [12] Normenyo B. V., Luca F., Togbé A., *Repdigits as sums of four Fibonacci or Lucas numbers*, J. Integer Seq. **21** (2018), no. 7, Art. 18.7.7, 30 pages.
- [13] Šiar Z., Keskin R., *On the Diophantine equation  $F_n - F_m = 2^a$* , Colloq. Math. **159** (2020), no. 1, 119–126.
- [14] Trojovský P., *On the order of appearance of the difference of two Lucas numbers*, Miskolc Math. Notes **19** (2018), no. 1, 641–648.

M. Alan:

YILDIZ TECHNICAL UNIVERSITY, FACULTY OF ARTS AND SCIENCES,  
 MATHEMATICS DEPARTMENT, DAVUTPASA CAMPUS, 34210, ESENLER, ISTANBUL,  
 TURKEY

*E-mail:* alan@yildiz.edu.tr

(Received June 12, 2021, revised September 2, 2021)