

Huanyuan Li

Weak Serrin-type finite time blowup and global strong solutions for three-dimensional density-dependent heat conducting magnetohydrodynamic equations with vacuum

Applications of Mathematics, Vol. 68 (2023), No. 5, 593–621

Persistent URL: <http://dml.cz/dmlcz/151835>

Terms of use:

© Institute of Mathematics AS CR, 2023

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

WEAK SERRIN-TYPE FINITE TIME BLOWUP AND GLOBAL
STRONG SOLUTIONS FOR THREE-DIMENSIONAL
DENSITY-DEPENDENT HEAT CONDUCTING
MAGNETOHYDRODYNAMIC EQUATIONS WITH VACUUM

HUANYUAN LI, Zhengzhou

Received June 25, 2022. Published online October 27, 2022.

Abstract. This paper is concerned with a Cauchy problem for the three-dimensional (3D) nonhomogeneous incompressible heat conducting magnetohydrodynamic (MHD) equations in the whole space. First of all, we establish a weak Serrin-type blowup criterion for strong solutions. It is shown that for the Cauchy problem of the 3D nonhomogeneous heat conducting MHD equations, the strong solution exists globally if the velocity satisfies the weak Serrin's condition. In particular, this criterion is independent of the absolute temperature and magnetic field. Then as an immediate application, we prove the global existence and uniqueness of strong solution to the 3D nonhomogeneous heat conducting MHD equations under a smallness condition on the initial data. In addition, the initial vacuum is allowed.

Keywords: heat conducting MHD; Cauchy problem; blowup criterion; global strong solution; vacuum

MSC 2020: 35Q35, 76W05

1. INTRODUCTION AND MAIN RESULTS

The time evolution of a three-dimensional nonhomogeneous incompressible and heat conducting magnetohydrodynamic (MHD for short) fluid is governed by the following nonhomogeneous heat conducting MHD system:

$$(1.1) \quad \begin{cases} \partial_t \varrho + \operatorname{div}(\varrho \mathbf{u}) = 0, \\ \partial_t(\varrho \mathbf{u}) + \operatorname{div}(\varrho \mathbf{u} \otimes \mathbf{u}) - \operatorname{div}(2\mu \mathfrak{D}(\mathbf{u})) + \nabla P = (\mathbf{H} \cdot \nabla) \mathbf{H}, \\ c_v[\partial_t(\varrho \theta) + \operatorname{div}(\varrho \mathbf{u} \theta)] - \kappa \Delta \theta = 2\mu |\mathfrak{D}(\mathbf{u})|^2 + \nu |\nabla \times \mathbf{H}|^2, \\ \partial_t \mathbf{H} + (\mathbf{u} \cdot \nabla) \mathbf{H} - (\mathbf{H} \cdot \nabla) \mathbf{u} = \nu \Delta \mathbf{H}, \\ \operatorname{div} \mathbf{u} = \operatorname{div} \mathbf{H} = 0, \end{cases}$$

This work was supported by the National Natural Science Foundation of China (Grant No. 12001495).

where $t \geq 0$ stands for the time and $x \in \mathbb{R}^3$ for the spatial coordinate. Moreover, ϱ , $\mathbf{u} = (u^1, u^2, u^3)$, P , θ and $\mathbf{H} = (H^1, H^2, H^3)$ denote the fluid density, velocity, pressure, absolute temperature and magnetic field, respectively. The positive constants μ , c_v , κ and ν are the viscosity coefficient, heat capacity, heat conductivity coefficient and magnetic diffusive coefficient, respectively.

$$\mathfrak{D}(\mathbf{u}) = \frac{1}{2}[\nabla \mathbf{u} + (\nabla \mathbf{u})^\top]$$

is the deformation tensor, where $\nabla \mathbf{u}$ is the gradient matrix $(\partial u^i / \partial x_j)_{ij}$ and $(\nabla \mathbf{u})^\top$ is its transpose, and $\nabla \times \mathbf{H}$ is the curl of the magnetic field $\mathbf{H}$.

In this paper, we will study the Cauchy problem of equation (1.1) with the initial conditions

$$(1.2) \quad (\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, 0) = (\varrho_0, \mathbf{u}_0, \mathbf{H}_0, \theta_0)(x), \quad x \in \mathbb{R}^3,$$

and the far field behavior conditions (in a weak sense)

$$(1.3) \quad (\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, t) \rightarrow (0, \mathbf{0}, \mathbf{0}, 0) \quad \text{as } |x| \rightarrow \infty.$$

Magnetohydrodynamics is the study of the interaction between magnetic field and moving conducting fluids. It is one of the important macroscopic fluid models, usually arising in science and engineering with a wide range of applications. Examples of such a magneto-fluids include hot ionised gases (plasmas), liquid metals or strong electrolytes. Because of the physical description of magneto-fluids dynamics, the nonhomogeneous incompressible MHD system (1.1) is a combination of the nonhomogeneous Navier-Stokes equations of fluid mechanics and the Maxwell equations of electromagnetism. The concept behind MHD is that the magnetic field can induce currents in a moving conducting fluid, which in turn polarizes the fluid and changes the magnetic field itself. One of the important issues is to understand the nature of this coupling between fluids and magnetic fields. We refer to [6] for more background and applications of MHD.

The mathematical studies of the nonhomogeneous incompressible fluids attract a lot of attention due to their physical importance, mathematical challenge and widespread applications. Let us briefly give a short survey on the nonhomogeneous fluids which are related to our results in this paper.

When we do not take account of equation (1.1)₃ for temperature, (1.1) reduces to the nonhomogeneous incompressible MHD equations. For this system, when the initial density has a positive lower bound, Gerbeau, Le Bris [9] and Desjardins, Le Bris [7] studied the global existence of weak solutions of finite energy in the

whole space and in the torus, respectively. Chen et al. [3] proved the existence of a global solution for the initial data belonging to critical Besov spaces. See also [1] for related improvement. Besides, Chen et al. [2] showed global well-posedness to the 3D Cauchy problem for discontinuous initial density. On the other hand, in the presence of vacuum, Chen et al. [4] obtained the local existence of strong solutions to the 3D Cauchy problem under a compatibility condition on the initial data. With the help of a Sobolev inequality of logarithmic type, Huang and Wang [14] showed the global existence of the strong solution for general initial data in dimension two.

When we study the motion in the absence of magnetic field, namely, $\mathbf{H} \equiv 0$, (1.1) reduces to the nonhomogeneous heat conducting Navier-Stokes equations. Under compatibility conditions for the initial data, Zhong [22] showed a Serrin-type blow-up criterion and proved global strong solutions with vacuum for small initial data, which extended the local result obtained by Cho and Kim [5] to a global one. By employing certain time-weighted a priori estimates, they showed that strong solutions exist globally provided that a smallness condition holds true. Meanwhile, Wang et al. [20] studied a three-dimensional initial boundary value problem with the general external force and obtained global existence of strong solutions under the assumption that the initial density is suitably small. Very recently, combining delicate energy estimates and a logarithmic interpolation inequality, the author established the global existence and uniqueness of strong solutions to the 2D Cauchy problem with large initial data and non-vacuum density at infinity.

Let us go back to the heat conducting MHD system (1.1). The local existence of a unique strong solution to system (1.1) with vacuum under some compatibility conditions was proved by Wu [21]. For the 2D problem, Zhong [24], [25] used a logarithmic interpolation inequality to prove the global well-posedness to the Cauchy problem and initial and boundary value problem for large initial data, respectively. And he also proved the global existence of a strong solution of initial and boundary problem with density-dependent viscosity in [26]. For 3D initial and boundary value problem, Zhong [23] obtained a global solution under some smallness conditions on the initial data, while Zhou [27] established a Serrin-type blowup criterion involving only the velocity field. Later, Zhu and Ou [28] extended the corresponding result in [23] to the case of density-temperature-dependent viscosity. However, the global well-posedness of system (1.1) in the unbounded domains is still unknown. In fact, this is the main purpose of this paper.

Before stating our main results, we first explain the notations and conventions used throughout this paper. We denote

$$\int \cdot dx = \int_{\mathbb{R}^3} \cdot dx.$$

And for $1 \leq r \leq \infty$ and $k \in \mathbb{N}$, the homogeneous and inhomogeneous Sobolev spaces are defined in a standard way:

$$\begin{cases} L^r = L^r(\mathbb{R}^3), & W^{k,r} = W^{k,r}(\mathbb{R}^3), & H^k = W^{k,2}, \\ D^{k,r} = \{f \in L^1_{\text{loc}} : \|\nabla^k f\|_{L^r} < \infty\}, & D^k = D^{k,2}, \\ D^1_0 = \{f \in L^6(\mathbb{R}^3) : \|\nabla u\|_{L^2} < \infty\}. \end{cases}$$

Now we give the definition of strong solutions to the Cauchy problem (1.1)–(1.3) as follows.

Definition 1.1 (Strong solutions). A pair of functions $(\varrho \geq 0, \mathbf{u}, \mathbf{H}, \theta \geq 0)$ is called a strong solution to the Cauchy problem (1.1)–(1.3) in $\mathbb{R}^3 \times (0, T)$ if for some $q_0 \in (3, \infty)$,

$$(1.4) \quad \begin{cases} \varrho \in C([0, T]; H^1 \cap W^{1, q_0}), & \varrho_t \in C([0, T]; L^{q_0}), \\ (\mathbf{u}, \mathbf{H}, \theta) \in C([0, T]; D^1_0 \cap D^2) \cap L^2(0, T; D^{2, q_0}), \\ (\sqrt{\varrho} \mathbf{u}_t, \mathbf{H}_t, \sqrt{\varrho} \theta_t) \in L^\infty(0, T; L^2), & (\mathbf{u}_t, \mathbf{H}_t, \theta_t) \in L^2(0, T; D^1_0), \end{cases}$$

and $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ satisfies (1.1) almost everywhere in $\mathbb{R}^3 \times (0, T)$.

Our main results read as follows:

Theorem 1.2. For a constant $q \in (3, 6]$, assume that the initial data $(\varrho_0 \geq 0, \mathbf{u}_0, \mathbf{H}_0, \theta_0 \geq 0)$ satisfy

$$(1.5) \quad \varrho \in L^1 \cap H^1 \cap W^{1, q}, \quad (\mathbf{u}_0, \mathbf{H}_0, \theta_0) \in D^1_0 \cap D^2, \quad \operatorname{div} \mathbf{u}_0 = \operatorname{div} \mathbf{H}_0 = 0,$$

and the compatibility conditions

$$(1.6) \quad -\mu \Delta \mathbf{u}_0 - \mathbf{H}_0 \cdot \nabla \mathbf{H}_0 + \nabla P_0 = \sqrt{\varrho_0} \mathbf{g}_1$$

and

$$(1.7) \quad \kappa \Delta \theta_0 + 2\mu |\mathfrak{D}(\mathbf{u}_0)|^2 + \nu |\nabla \times \mathbf{H}_0|^2 = \sqrt{\varrho_0} \mathbf{g}_2$$

for some $P_0 \in D^1$, and $\mathbf{g}_1, \mathbf{g}_2 \in L^2$. Let $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ be a strong solution in $\mathbb{R}^3 \times (0, T^*)$ as described in Definition 1.1. If $T^* < \infty$ is the maximal existence time, then

$$(1.8) \quad \lim_{T \rightarrow T^*} \|\mathbf{u}\|_{L^s(0, T; L^r_\omega)} = \infty$$

for any r and s satisfying

$$(1.9) \quad \frac{2}{s} + \frac{3}{r} \leq 1, \quad 3 < r \leq \infty,$$

where L^r_ω denotes the weak- L^r space.

Remark 1.3. The local existence of a unique strong solution to (1.1)–(1.3) with the initial data described in Theorem 1.2 can be established in a similar way as [5] (see also [21]). Hence, the maximal time T^* is well-defined.

Remark 1.4. It should be pointed out that the blowup criterion (1.8) is independent of both the temperature and magnetic field, which is the same as the weak Serrin-type blowup criterion of homogeneous Navier-Stokes equations (see the work of Sohr [18]).

Remark 1.5. The approach can be adapted to deal with the case of bounded domain in $\mathbb{R}^3$. And compared with [27] for bounded domain, some new difficulties occur in our analysis. First, the Poincaré's inequality fails for 3D Cauchy problem, which is key to estimate $\|\theta\|_{L^2}$. Furthermore, it implies the blowup criterion (1.8) is stronger than that of [27] due to $\|\mathbf{u}\|_{L^s(0,T;L^\infty_\omega)} \leq \|\mathbf{u}\|_{L^s(0,T;L^r)}$.

The proof of Theorem 1.2 will be done by contradiction. In view of the local existence result, to prove Theorem 1.2 it suffices to verify that $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ satisfy (1.5)–(1.7) at the time T^* under the assumption that the left-hand side of (1.8) is finite, then apply the local existence result to extend a local solution beyond the maximal existence time T^* , consequently leading to a contradiction.

Based on Theorem 1.2, we can establish the global existence of strong solutions to (1.1)–(1.3) under a smallness condition on the initial data.

Theorem 1.6. *Let the conditions in Theorem 1.2 hold. Then there exists a small positive constant ε_0 depending only on μ, ν , and $\|\varrho_0\|_{L^\infty}$ such that if*

$$(1.10) \quad (\|\sqrt{\varrho_0}\mathbf{u}_0\|_{L^2}^2 + \|\mathbf{H}_0\|_{L^2}^2)(\|\nabla\mathbf{u}_0\|_{L^2}^2 + \|\nabla\mathbf{H}_0\|_{L^2}^2) \leq \varepsilon_0,$$

then the Cauchy problem of system (1.1)–(1.3) admits a unique global strong solution.

We now comment on the analysis of this paper. The study of weak Serrin-type blowup criterion (Theorem 1.2) is mainly motivated by a recent work of Wang [19], which established a Serrin's blowup criterion for nonhomogeneous heat conducting Navier-Stokes equations in the whole space $\mathbb{R}^3$ using the weak Lebesgue spaces. Compared to Navier-Stokes model in [19], the mathematical analysis of nonhomogeneous heat conducting MHD system will be more complicated on the account of the coupling of the velocity and magnetic field (such as the term $\mathbf{u} \cdot \nabla \mathbf{H}$) and strong nonlinearity (such as the term $\mathbf{H} \cdot \nabla \mathbf{H}$). To overcome these difficulties, one of the key ideas is to derive an estimate of $\|\mathbf{H}\|_{L^\infty(0,T;L^q)}$ for $q > 2$ which turns out to play an important role in our analysis. It should be noted that our blowup criterion is

independent of the temperature and magnetic field, which means the temperature and magnetic field do not play a particular role when the singularity of solution $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ forms in finite time.

As an immediate application of the blowup criterion obtained in Theorem 1.2, we plan to extend the local strong solution to be a global one under a smallness condition on the initial data. Noticing that the pair $(s, r) = (4, 6)$ satisfies $2/s + 3/r \leq 1$, we conclude that the global existence of a unique strong solution can be verified if we can obtain the uniformly time independent estimate on the $L^2(0, T; L^2)$ -norm of the gradient of the velocity. To this end, we multiply the momentum equations by $\mathbf{u}_t$ and make good use of the smallness of initial data to obtain the desired estimate.

The remainder of this paper is arranged as follows. In Section 2, we give some auxiliary lemmas which will be useful in our later analysis. The proof of Theorem 1.2 will be done by combining the contradiction argument with the estimates derived in Section 3. Finally, we give the proof of Theorem 1.6 in Section 4.

2. PRELIMINARIES

In this section, we will recall some known facts and analytic inequalities that will be used in the later analysis.

We begin with the following local existence and uniqueness of strong solutions when the initial data is allowed vacuum, which can be proved in a similar way as [5] (see also [21]).

Lemma 2.1. *Assume that the initial data $(\varrho_0, \mathbf{u}_0, \mathbf{H}_0, \theta_0)$ satisfy (1.5)–(1.7). Then there exist positive time T_1 and a unique strong solution to the Cauchy problem (1.1)–(1.3) on $\mathbb{R}^3 \times (0, T_1]$.*

Next, we will introduce the well-known Gagliardo-Nirenberg inequality which will be frequently used later. See [10], Chapter 6 for the proof and more details.

Lemma 2.2. *For $p \in [2, 6]$, $q \in (1, \infty)$ and $r \in (3, \infty)$, there exists a generic constant C which may depend only on p, q and r , such that for $f \in H^1$, $g \in L^q \cap D^{1,r}$, the following inequalities hold:*

$$(2.1) \quad \|f\|_{L^p} \leq C \|f\|_{L^2}^{(6-p)/(2p)} \|\nabla f\|_{L^2}^{(3p-6)/(2p)},$$

$$(2.2) \quad \|g\|_{L^\infty} \leq C \|g\|_{L^q} + C \|\nabla g\|_{L^r}.$$

Since our blowup criterion (1.8) involves a weak Lebesgue space, it is necessary to give a short introduction and state related inequalities. Denote the Lorentz space

and its norm by $L^{p,q}$ and $\|\cdot\|_{L^{p,q}}$, respectively, where $1 < p < \infty$ and $1 \leq q \leq \infty$. And we recall the weak- L^p space L^p_ω which is defined as follows:

$$L^p_\omega := \{f \in L^1_{\text{loc}} : \|f\|_{L^p_\omega} = \sup_{\lambda > 0} \lambda |\{ |f(x)| > \lambda \}|^{1/p} < \infty\}.$$

It should be noted that

$$L^p \subsetneq L^p_\omega, \quad L^\infty = L^\infty, \quad L^p_\omega = L^{p,\infty}, \quad L^{p,p} = L^p, \quad \|f\|_{L^p_\omega} \leq \|f\|_{L^p}.$$

For the details of Lorentz space, we refer to the monograph by Grafakos [11]. In particular, we introduce the following Hölder's inequality in Lorentz space, whose proof can be found in [16].

Lemma 2.3. *Let $p_1, p_2 \in (0, \infty)$, $q_1, q_2 \in [1, \infty]$ satisfying $1/p = 1/p_1 + 1/p_2 < 1$ and $q = \min\{q_1, q_2\}$. Then for $f \in L^{p_1, q_1}$ and $g \in L^{p_2, q_2}$, there exists a positive constant C depending on p_1, p_2, q_1 and q_2 such that $f \cdot g \in L^{p, q}$ satisfying*

$$(2.3) \quad \|f \cdot g\|_{L^{p,q}} \leq C \|f\|_{L^{p_1, q_1}} \|g\|_{L^{p_2, q_2}}.$$

Based on Lemma 2.3, we have the following result involving the weak Lebesgue spaces, which will play an important role in the subsequent analysis.

Lemma 2.4. *Assume $g \in H^1$ and $f \in L^r_\omega$ with $r \in (3, \infty]$. Then $f \cdot g \in L^2$. Furthermore, for any $\varepsilon > 0$ we have*

$$(2.4) \quad \|f \cdot g\|_{L^2}^2 \leq \varepsilon \|\nabla g\|_{L^2}^2 + C(\varepsilon)(\|f\|_{L^r_\omega}^s + 1)\|g\|_{L^2}^2,$$

where C is a positive constant depending only on ε and r .

Proof. Modifying the proof in [15] for bounded domains slightly, it follows from (2.3) and the interpolation inequality that

$$\begin{aligned} (2.5) \quad \|f \cdot g\|_{L^2}^2 &= \|f \cdot g\|_{L^{2,2}}^2 \leq C \|f\|_{L^{r,\infty}} \|g\|_{L^{2r/(r-2),2}} \\ &\leq C \|f\|_{L^r_\omega} \|g\|_{L^{2r_1/(r_1-2)}} \|g\|_{L^{2r_2/(r_2-2)}} \\ &\leq C \|f\|_{L^r_\omega} \|g\|_{L^2}^{(r_1-3)/r_1} \|g\|_{L^6}^{3/r_1} \|g\|_{L^2}^{(r_2-3)/r_2} \|g\|_{L^6}^{3/r_2} \\ &\leq C \|f\|_{L^r_\omega} \|g\|_{L^2}^{(2r-6)/r} \|g\|_{L^6}^{6/r} \\ &\leq C \|f\|_{L^r_\omega} \|g\|_{L^2}^{(2r-6)/r} \|\nabla g\|_{L^2}^{6/r} \\ &\leq \varepsilon \|\nabla g\|_{L^2}^2 + C(\varepsilon)(\|f\|_{L^r_\omega}^s + 1)\|g\|_{L^2}^2, \end{aligned}$$

where r_1, r_2 and r satisfy $3 < r_1 < r < r_2 < \infty$, $2/r = 1/r_1 + 1/r_2$ and $2/s + 3/r \leq 1$. This completes the proof of Lemma 2.4. $\square$

Finally, we give classical regularity results for the Stokes system in the whole space $\mathbb{R}^3$, which have been proved in [12].

Lemma 2.5. *For any $r \in (1, \infty)$, if $\mathbf{F} \in L^r$, there exists a positive constant C depending only on r such that the unique weak solution $(\mathbf{u}, P) \in D^1 \times L^2$ to the Stokes system*

$$(2.6) \quad \begin{cases} -\Delta \mathbf{u} + \nabla P = \mathbf{F} & \text{in } \mathbb{R}^3, \\ \operatorname{div} \mathbf{u} = 0 & \text{in } \mathbb{R}^3, \\ \mathbf{u}(x) \rightarrow 0 & \text{as } |x| \rightarrow \infty, \end{cases}$$

satisfies

$$(2.7) \quad \|\nabla^2 \mathbf{u}\|_{L^r} + \|\nabla P\|_{L^r} \leq C \|\mathbf{F}\|_{L^r}.$$

3. PROOF OF THEOREM 1.2

This section is devoted to giving a proof of Theorem 1.2 using the contradiction argument. To do this, let $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ be a strong solution to the Cauchy problem (1.1)–(1.3) as described in Lemma 2.1, and T^* be the maximal existence time of the strong solution. Suppose that (1.8) in Theorem 1.2 were false, that is to say, there exists a positive constant M_0 such that

$$(3.1) \quad \lim_{T \rightarrow T^*} \|\mathbf{u}\|_{L^s(0, T; L_\omega^r)} \leq M_0 < \infty.$$

Under condition (3.1), we will extend the existence time of the strong solution beyond T^* , which contradicts the definition of the maximum of T^* .

Before proceeding, it is easy to rewrite system (1.1) in the following form if we assume the solution $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ is regular enough:

$$(3.2) \quad \begin{cases} \partial_t \varrho + \mathbf{u} \cdot \nabla \varrho = 0, \\ \varrho \partial_t \mathbf{u} + \varrho \mathbf{u} \cdot \nabla \mathbf{u} - \mu \Delta \mathbf{u} + \nabla P = (\mathbf{H} \cdot \nabla) \mathbf{H}, \\ c_v (\varrho \partial_t \theta + \varrho \mathbf{u} \cdot \nabla \theta) - \kappa \Delta \theta = 2\mu |\mathfrak{D}(\mathbf{u})|^2 + \nu |\nabla \times \mathbf{H}|^2, \\ \partial_t \mathbf{H} + (\mathbf{u} \cdot \nabla) \mathbf{H} - (\mathbf{H} \cdot \nabla) \mathbf{u} = \nu \Delta \mathbf{H}, \\ \operatorname{div} \mathbf{u} = \operatorname{div} \mathbf{H} = 0. \end{cases}$$

In this section, the symbol C denotes a generic constant which may depend on $M_0, \mu, \nu, c_v, \kappa, T^*$, and the initial data.

Now we establish some a priori estimates which will be used to prove Theorem 1.2 at the end of this section.

3.1. Lower-order estimates. In this subsection, we will derive a series of key lower-order estimates to $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$

First, it follows from the transport equation $(3.2)_1$ for the density and incompressibility condition $\operatorname{div} \mathbf{u} = 0$ that the following result holds.

Lemma 3.1. *There exists a positive constant C satisfying*

$$(3.3) \quad \sup_{0 \leq t \leq T} \|\varrho\|_{L^1 \cap L^\infty} \leq C, \quad 0 \leq T < T^*.$$

Next, the standard energy estimates read as follows.

Lemma 3.2. *It holds that for any $0 \leq T < T^*$,*

$$(3.4) \quad \sup_{0 \leq t \leq T} (\|\sqrt{\varrho} \mathbf{u}\|_{L^2}^2 + \|\mathbf{H}\|_{L^2}^2 + \|\varrho \theta\|_{L^1}) + \int_0^T (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) dt \leq C.$$

Proof. It follows from the standard maximum principle to $(1.1)_3$ together with $\theta_0 \geq 0$ that (see [8] for the proof)

$$\inf_{\mathbb{R}^3 \times [0, T]} \theta(x, t) \geq 0.$$

Moreover, multiplying $(1.1)_2$ by $\mathbf{u}$, $(3.2)_4$ by $\mathbf{H}$, and integrating the resulting equations over $\mathbb{R}^3$, it follows from integrating by parts that

$$(3.5) \quad \frac{1}{2} \frac{d}{dt} \int (\varrho |\mathbf{u}|^2 + |\mathbf{H}|^2) dx + \int (\mu |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2) dx = 0.$$

Integrating $(1.1)_3$ with respect to the spatial variable over $\mathbb{R}^3$ and performing integration by parts, we obtain that

$$(3.6) \quad c_v \frac{d}{dt} \int \varrho \theta dx = \int (2\mu |\mathfrak{D}(\mathbf{u})|^2 + \nu |\nabla \times \mathbf{H}|^2) dx.$$

By the definition of $\mathfrak{D}(\mathbf{u})$ and integration by parts, we get

$$(3.7) \quad \begin{aligned} 2\mu \int |\mathfrak{D}(\mathbf{u})|^2 dx &= \frac{\mu}{2} \int (\partial_i u^j + \partial_j u^i)^2 dx \\ &= \mu \int |\partial_i u^j|^2 dx + \mu \int \partial_i u^j \partial_j u^i dx = \mu \int |\nabla \mathbf{u}|^2 dx, \end{aligned}$$

and it follows from $-\Delta \mathbf{H} = \nabla \times (\nabla \times \mathbf{H})$ (since $\operatorname{div} \mathbf{H} = 0$) that

$$(3.8) \quad \|\nabla \mathbf{H}\|_{L^2}^2 = \|\nabla \times \mathbf{H}\|_{L^2}^2.$$

Substituting (3.7) and (3.8) into (3.7) gives

$$(3.9) \quad c_v \frac{d}{dt} \int \varrho \theta \, dx = \int (\mu |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2) \, dx.$$

Therefore, adding (3.5) multiplied by 2 to (3.9), we have

$$(3.10) \quad \frac{d}{dt} \int (\varrho |\mathbf{u}|^2 + |\mathbf{H}|^2 + c_v \varrho \theta) \, dx + \int (\mu |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2) \, dx = 0.$$

Integrating (3.10) with respect to t over $[0, T]$ leads to the desired (3.4), which completes the proof of Lemma 3.2. $\square$

Before deriving the key estimates of $\|\nabla \mathbf{u}\|_{L^\infty(0, T; L^2)}$ and $\|\nabla \mathbf{H}\|_{L^\infty(0, T; L^2)}$, we insert an important estimate on magnetic field $\mathbf{H}$ initiated by He and Xin [13], which will be stated in the following lemma.

Lemma 3.3. *Under condition (3.1), it holds that for $q \in [2, 12]$ and $0 \leq T < T^*$,*

$$(3.11) \quad \sup_{0 \leq t \leq T} \|\mathbf{H}\|_{L^q}^q + \int_0^T \int |\mathbf{H}|^{q-2} |\nabla \mathbf{H}|^2 \, dx \, dt \leq C.$$

Proof. Multiplying (3.2)₄ by $q|\mathbf{H}|^{q-2}\mathbf{H}$ and integrating the resulting equation over $\mathbb{R}^3$, it follows from (2.4) in Lemma 2.4 that

$$\begin{aligned} (3.12) \quad & \frac{d}{dt} \int |\mathbf{H}|^q \, dx + \nu \int (q|\mathbf{H}|^{q-2} |\nabla \mathbf{H}|^2 + q(q-2)|\mathbf{H}|^{q-2} |\nabla |\mathbf{H}||^2) \, dx \\ &= - \int q|\mathbf{H}|^{q-2} \left(\mathbf{H} \cdot \nabla \mathbf{H} \cdot \mathbf{u} - \frac{q-1}{2} \mathbf{u} \cdot \nabla |\mathbf{H}|^2 \right) \, dx \\ &\quad - \frac{q(q-2)}{2} \int |\mathbf{H}|^{q-4} (\mathbf{H} \cdot \nabla |\mathbf{H}|^2) (\mathbf{u} \cdot \mathbf{H}) \, dx \\ &\leq \frac{\nu}{2} \int q|\mathbf{H}|^{q-2} |\nabla \mathbf{H}|^2 \, dx + Cq^2 \int |\mathbf{u}|^2 |\mathbf{H}|^q \, dx \\ &= \frac{\nu}{2} \int q|\mathbf{H}|^{q-2} |\nabla \mathbf{H}|^2 \, dx + Cq^2 \|\mathbf{u}\| |\mathbf{H}|^{q/2} \| \mathbf{H} \|_{L^2}^2 \\ &\leq \frac{\nu}{2} \int q|\mathbf{H}|^{q-2} |\nabla \mathbf{H}|^2 \, dx + \varepsilon \|\nabla |\mathbf{H}|^{q/2}\|_{L^2}^2 + C(\varepsilon) (1 + \|\mathbf{u}\|_{L_w^s}^s) \|\mathbf{H}\|_{L^q}^q. \end{aligned}$$

Choosing ε suitably small in (3.12), we obtain the desired (3.11) after applying Gronwall's inequality and (3.1). Therefore, the proof of Lemma 3.3 is completed. $\square$

With the help of Lemma 3.3, we can now derive key time-independent estimates on the $L^\infty(0, T; L^2)$ -norm of the gradients of velocity and magnetic field.

Lemma 3.4. *Under assumption (3.1), it holds for all $0 \leq T < T^*$ that*

$$(3.13) \quad \sup_{0 \leq t \leq T} (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) + \int_0^T (\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2 + \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) dt \leq C.$$

Proof. Multiplying (3.2)₂ by $\mathbf{u}_t$ and integrating the resulting equation over $\mathbb{R}^3$ lead to

$$(3.14) \quad \frac{\mu}{2} \frac{d}{dt} \int |\nabla \mathbf{u}|^2 dx + \int \varrho |\mathbf{u}_t|^2 dx = - \int \varrho \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx + \int \mathbf{H} \cdot \nabla \mathbf{H} \cdot \mathbf{u}_t dx.$$

It follows from equation (3.2)₄ that $\mathbf{H}_t - \nu \Delta \mathbf{H} = \mathbf{H} \cdot \nabla \mathbf{u} - \mathbf{u} \cdot \nabla \mathbf{H}$. Then we have

$$(3.15) \quad \int |\mathbf{H}_t - \nu \Delta \mathbf{H}|^2 dx = \int |\mathbf{u} \cdot \nabla \mathbf{H} - \mathbf{H} \cdot \nabla \mathbf{u}|^2 dx.$$

For the LHS of (3.15), it is easy to get

$$(3.16) \quad \begin{aligned} \int |\mathbf{H}_t - \nu \Delta \mathbf{H}|^2 dx &= \int (|\mathbf{H}_t|^2 + \nu^2 |\Delta \mathbf{H}|^2 - 2\nu \mathbf{H}_t \cdot \Delta \mathbf{H}) dx \\ &= \nu \frac{d}{dt} \int |\nabla \mathbf{H}|^2 dx + \int (|\mathbf{H}_t|^2 + \nu^2 |\Delta \mathbf{H}|^2) dx. \end{aligned}$$

Substituting (3.16) into the LHS of (3.15), we obtain that

$$(3.17) \quad \nu \frac{d}{dt} \int |\nabla \mathbf{H}|^2 dx + \int (|\mathbf{H}_t|^2 + \nu^2 |\Delta \mathbf{H}|^2) dx = \int |\mathbf{u} \cdot \nabla \mathbf{H} - \mathbf{H} \cdot \nabla \mathbf{u}|^2 dx.$$

Notice that the standard L^2 -estimate of elliptic system gives

$$(3.18) \quad \|\nabla^2 \mathbf{H}\|_{L^2}^2 \leq K \|\Delta \mathbf{H}\|_{L^2}^2$$

with a positive constant K . Adding (3.14) to (3.17), we derive from the Cauchy-Schwarz inequality and (3.18) that

$$(3.19) \quad \begin{aligned} &\frac{d}{dt} \left(\frac{\mu}{2} |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2 \right) dx + \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2 + \frac{\nu^2}{K} |\nabla^2 \mathbf{H}|^2) dx \\ &\leq \int \mathbf{H} \cdot \nabla \mathbf{H} \cdot \mathbf{u}_t dx - \int \varrho \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx + \int |\mathbf{u} \cdot \nabla \mathbf{H} - \mathbf{H} \cdot \nabla \mathbf{u}|^2 dx \\ &= - \frac{d}{dt} \int (\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H} dx + \int (\mathbf{H}_t \cdot \nabla) \mathbf{u} \cdot \mathbf{H} dx + \int (\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H}_t dx \\ &\quad - \int \varrho \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx + \int |\mathbf{u} \cdot \nabla \mathbf{H} - \mathbf{H} \cdot \nabla \mathbf{u}|^2 dx \\ &\leq - \frac{d}{dt} \int (\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H} dx + \frac{1}{2} \int \varrho |\mathbf{u}_t|^2 dx + \frac{1}{2} \int |\mathbf{H}_t|^2 dx \\ &\quad + C \int |\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}|^2 dx + C \int |\mathbf{u} \cdot \nabla \mathbf{H}|^2 dx + C \int |\mathbf{H} \cdot \nabla \mathbf{u}|^2 dx. \end{aligned}$$

Thus, we obtain that

$$\begin{aligned}
(3.20) \quad & \frac{d}{dt} \int (\mu |\nabla \mathbf{u}|^2 + 2\nu |\nabla \mathbf{H}|^2 + 2(\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H}) \, dx \\
& + \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2 + \frac{2\nu^2}{K} |\nabla^2 \mathbf{H}|^2) \, dx \\
& \leq C \int |\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}|^2 \, dx + C \int |\mathbf{u} \cdot \nabla \mathbf{H}|^2 \, dx + C \int |\mathbf{H} \cdot \nabla \mathbf{u}|^2 \, dx.
\end{aligned}$$

Recall that $(\mathbf{u}, P)$ satisfies the following Stokes system:

$$(3.21) \quad \begin{cases} -\mu \Delta \mathbf{u} + \nabla P = -\varrho \mathbf{u}_t - \varrho \mathbf{u} \cdot \nabla \mathbf{u} + \mathbf{H} \cdot \nabla \mathbf{H}, & x \in \mathbb{R}^3, \\ \operatorname{div} \mathbf{u} = 0, & x \in \mathbb{R}^3, \\ \mathbf{u}(x) \rightarrow 0, & |x| \rightarrow \infty. \end{cases}$$

Applying Lemma 2.5 with $\mathbf{F} \triangleq -\varrho \mathbf{u}_t - \varrho \mathbf{u} \cdot \nabla \mathbf{u} + \mathbf{H} \cdot \nabla \mathbf{H}$, we obtain from (2.7) that

$$\begin{aligned}
(3.22) \quad & \|\nabla^2 \mathbf{u}\|_{L^2}^2 \leq C(\|\varrho \mathbf{u}_t\|_{L^2}^2 + \|\varrho \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2) \\
& \leq L(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2),
\end{aligned}$$

where L is a positive constant depending only on μ and $\|\varrho_0\|_{L^\infty}$.

Adding (3.22) multiplied by $1/(2L)$ to (3.20), we have

$$\begin{aligned}
(3.23) \quad & \frac{d}{dt} \int (\mu |\nabla \mathbf{u}|^2 + 2\nu |\nabla \mathbf{H}|^2 + 2(\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H}) \, dx \\
& + \int \left(\frac{1}{2} \varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2 + \frac{2\nu^2}{K} |\nabla^2 \mathbf{H}|^2 + \frac{1}{2L} \|\nabla^2 \mathbf{u}\|_{L^2}^2 \right) \, dx \\
& \leq C \int (|\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}|^2 + |\mathbf{u} \cdot \nabla \mathbf{H}|^2 + |\mathbf{H} \cdot \nabla \mathbf{u}|^2 + |\mathbf{H} \cdot \nabla \mathbf{H}|^2) \, dx \\
& \leq C \|\varrho\|_{L^\infty} \|\mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + C \|\mathbf{u} \cdot \nabla \mathbf{H}\|_{L^2}^2 + C \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla \mathbf{u}\|_{L^6} \\
& \quad + C \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla \mathbf{H}\|_{L^6} \\
& \leq \frac{\varepsilon}{2} (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) + C(\varepsilon)(1 + \|\mathbf{u}\|_{L_\omega^s}^s) (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) \\
& \quad + C \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla^2 \mathbf{u}\|_{L^2} + C \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla^2 \mathbf{H}\|_{L^2} \\
& \leq \varepsilon (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) + C(\varepsilon)(1 + \|\mathbf{u}\|_{L_\omega^s}^s) (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2),
\end{aligned}$$

due to Lemma 2.4, (3.11) and the Cauchy-Schwarz inequality.

Moreover, applying the Cauchy-Schwarz inequality and (3.11) gives rise to

$$\begin{aligned}
(3.24) \quad & 2 \int (\mathbf{H} \cdot \nabla) \mathbf{u} \cdot \mathbf{H} \, dx \leq C \|\mathbf{H}\|_{L^4}^2 \|\nabla \mathbf{u}\|_{L^2} \leq \frac{\mu}{2} \|\nabla \mathbf{u}\|_{L^2}^2 + C \|\mathbf{H}\|_{L^4}^4 \\
& \leq \frac{\mu}{2} \|\nabla \mathbf{u}\|_{L^2}^2 + C.
\end{aligned}$$

Together with Gronwall's inequality, (3.23) implies

$$(3.25) \quad \sup_{0 \leq t \leq T} (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) + \int_0^T (\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2 + \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) dt \leq C.$$

Therefore, the proof of Lemma 3.4 is completed. $\square$

The following lemma concerns the higher regularity of the temperature θ .

Lemma 3.5. *Under assumption (3.1), it holds that for $0 \leq T < T^*$,*

$$(3.26) \quad \sup_{0 \leq t \leq T} \|\sqrt{\varrho} \theta\|_{L^2}^2 + \int_0^T \|\nabla \theta\|_{L^2}^2 dt \leq C.$$

Proof. Multiplying (3.2)₃ by θ and integrating the resulting equation over $\mathbb{R}^3$ implies that

$$(3.27) \quad c_v \frac{d}{dt} \int \varrho \theta^2 dx + 2\kappa \int |\nabla \theta|^2 dx \leq C \int |\nabla \mathbf{u}|^2 \theta dx + C \int |\nabla \mathbf{H}|^2 \theta dx.$$

We estimate each term of the RHS of (3.27) as follows. Applying Hölder's and the Cauchy-Schwarz inequalities gives

$$(3.28) \quad \begin{aligned} \int |\nabla \mathbf{u}|^2 \theta dx &\leq C \|\nabla \mathbf{u}\|_{L^{12/5}}^2 \|\theta\|_{L^6} \leq C \|\nabla \mathbf{u}\|_{L^2}^{3/2} \|\nabla \mathbf{u}\|_{L^6}^{1/2} \|\nabla \theta\|_{L^2} \\ &\leq \frac{\kappa}{2} \|\nabla \theta\|_{L^2}^2 + C \|\nabla \mathbf{u}\|_{L^2}^3 \|\nabla^2 \mathbf{u}\|_{L^2} \leq \frac{\kappa}{2} \|\nabla \theta\|_{L^2}^2 + C \|\nabla^2 \mathbf{u}\|_{L^2}^2 + C, \end{aligned}$$

due to (3.13). And in a similar way, we have

$$(3.29) \quad \int |\nabla \mathbf{H}|^2 \theta dx \leq \frac{\kappa}{2} \|\nabla \theta\|_{L^2}^2 + C \|\nabla^2 \mathbf{H}\|_{L^2}^2 + C.$$

Substituting (3.28) and (3.29) into (3.27), we obtain

$$(3.30) \quad c_v \frac{d}{dt} \int \varrho \theta^2 dx + \kappa \int |\nabla \theta|^2 dx \leq C \|\nabla^2 \mathbf{u}\|_{L^2}^2 + C \|\nabla^2 \mathbf{H}\|_{L^2}^2 + C.$$

Integrating inequality (3.30) with respect to the time variable over $(0, t)$, we get from (3.13) that

$$(3.31) \quad \sup_{0 \leq t \leq T} \|\sqrt{\varrho} \theta\|_{L^2}^2 + \int_0^T \|\nabla \theta\|_{L^2}^2 dt \leq C.$$

Therefore, the proof of Lemma 3.5 is completed. $\square$

At the end of this subsection, we give the following remark which will be used later.

Remark 3.6. In view of Lemma 3.4, we deduce from classical L^2 -estimates for elliptic and Stokes system that

$$\begin{aligned}
 (3.32) \quad \|\nabla^2 \mathbf{u}\|_{L^2}^2 &\leq C(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2) \\
 &\leq C(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{u}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^3}^2 + \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^3}^2) \\
 &\leq C(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^2}^3 \|\nabla^2 \mathbf{u}\|_{L^2} + \|\nabla \mathbf{H}\|_{L^2}^3 \|\nabla^2 \mathbf{H}\|_{L^2}) \\
 &\leq \frac{1}{4}(\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) + C\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 \\
 &\quad + C\|\nabla \mathbf{u}\|_{L^2}^6 + C\|\nabla \mathbf{H}\|_{L^2}^6,
 \end{aligned}$$

and

$$\begin{aligned}
 (3.33) \quad \|\nabla^2 \mathbf{H}\|_{L^2}^2 &\leq C(\|\mathbf{H}_t\|_{L^2}^2 + \|\mathbf{u} \cdot \nabla \mathbf{H}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{u}\|_{L^2}^2) \\
 &\leq C(\|\mathbf{H}_t\|_{L^2}^2 + \|\mathbf{u}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^3}^2 + \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^3}^2) \\
 &\leq C(\|\mathbf{H}_t\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^2}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla^2 \mathbf{H}\|_{L^2} \\
 &\quad + \|\nabla \mathbf{H}\|_{L^2}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla^2 \mathbf{u}\|_{L^2}) \\
 &\leq \frac{1}{4}(\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) + C\|\mathbf{H}_t\|_{L^2}^2 \\
 &\quad + C\|\nabla \mathbf{u}\|_{L^2}^6 + C\|\nabla \mathbf{H}\|_{L^2}^6.
 \end{aligned}$$

Adding (3.32) to (3.33), it follows from (3.13) that

$$(3.34) \quad \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2 \leq C(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2) + C.$$

3.2. Higher-order estimates. In this subsection, we will derive a series of higher-order estimates of $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$.

Firstly, we will estimate the $L^\infty(0, T; L^2)$ -norm of $\sqrt{\varrho} \mathbf{u}_t$, $\mathbf{H}_t$ and $\nabla \theta$.

Lemma 3.7. *Under assumption (3.1), it holds that for any $0 \leq T < T^*$,*

$$\begin{aligned}
 (3.35) \quad &\sup_{0 \leq t \leq T} (\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2 + \|\nabla \theta\|_{L^2}^2) \\
 &\quad + \int_0^T (\|\nabla \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{H}_t\|_{L^2}^2 + \|\sqrt{\varrho} \theta_t\|_{L^2}^2) dt \leq C.
 \end{aligned}$$

P r o o f. Differentiating (3.2)₂ with respect to t yields

$$(3.36) \quad \varrho \mathbf{u}_{tt} + \varrho \mathbf{u} \cdot \nabla \mathbf{u}_t - \mu \Delta \mathbf{u}_t = -\nabla P_t - \varrho_t \mathbf{u}_t - (\varrho \mathbf{u})_t \cdot \nabla \mathbf{u} + (\mathbf{H}_t \cdot \nabla) \mathbf{H} + (\mathbf{H} \cdot \nabla) \mathbf{H}_t.$$

Multiplying equality (3.36) by $\mathbf{u}_t$ and integrating the resulting equation over $\mathbb{R}^3$, one obtains

$$(3.37) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} \int \varrho |\mathbf{u}_t|^2 dx + \mu \int |\nabla \mathbf{u}_t|^2 dx \\ &= - \int \varrho_t |\mathbf{u}_t|^2 dx - \int (\varrho \mathbf{u})_t \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx \\ & \quad + \int (\mathbf{H}_t \cdot \nabla) \mathbf{H} \cdot \mathbf{u}_t dx + \int (\mathbf{H} \cdot \nabla) \mathbf{H}_t \cdot \mathbf{u}_t dx. \end{aligned}$$

Next, differentiating (3.2)₄ with respect to t and multiplying the resulting equation by $\mathbf{H}_t$, we obtain from integration by parts that

$$(3.38) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} \int |\mathbf{H}_t|^2 dx + \nu \int |\nabla \mathbf{H}_t|^2 dx \\ &= - \int \mathbf{u}_t \cdot \nabla \mathbf{H} \cdot \mathbf{H}_t dx + \int (\mathbf{H}_t \cdot \nabla) \mathbf{u} \cdot \mathbf{H}_t dx + \int (\mathbf{H} \cdot \nabla) \mathbf{u}_t \cdot \mathbf{H}_t dx. \end{aligned}$$

Adding (3.38) into (3.37), we get

$$(3.39) \quad \begin{aligned} & \frac{1}{2} \frac{d}{dt} \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2) dx + \int (\mu |\nabla \mathbf{u}_t|^2 + \mu |\nabla \mathbf{H}_t|^2) dx \\ &= - \int \varrho_t |\mathbf{u}_t|^2 dx - \int (\varrho \mathbf{u})_t \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx \\ & \quad + \int (\mathbf{H}_t \cdot \nabla) \mathbf{H} \cdot \mathbf{u}_t dx - \int \mathbf{u}_t \cdot \nabla \mathbf{H} \cdot \mathbf{H}_t dx + \int (\mathbf{H}_t \cdot \nabla) \mathbf{u} \cdot \mathbf{H}_t dx \\ & \triangleq I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned}$$

The terms I_1 – I_5 are estimated as follows. Indeed, it follows from (2.1), (2.2), (3.13), integrating by parts and Hölder's inequality that

$$(3.40) \quad \begin{aligned} I_1 &= \int \operatorname{div}(\varrho \mathbf{u}) |\mathbf{u}_t|^2 dx = - \int \varrho \mathbf{u} \cdot \nabla |\mathbf{u}_t|^2 dx \\ &\leq 2 \int \varrho |\mathbf{u}| |\mathbf{u}_t| |\nabla \mathbf{u}_t| dx \leq C \|\varrho\|_{L^\infty}^{1/2} \|\mathbf{u}\|_{L^6} \|\sqrt{\varrho} \mathbf{u}_t\|_{L^3} \|\nabla \mathbf{u}_t\|_{L^2} \\ &\leq C \|\nabla \mathbf{u}\|_{L^2} \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^{1/2} \|\sqrt{\varrho} \mathbf{u}_t\|_{L^6}^{1/2} \|\nabla \mathbf{u}_t\|_{L^2} \\ &\leq C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^{1/2} \|\nabla \mathbf{u}_t\|_{L^6}^{3/2} \leq \frac{\mu}{8} \|\nabla \mathbf{u}_t\|_{L^2}^2 + C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2, \end{aligned}$$

and

$$\begin{aligned}
(3.41) \quad I_2 &= - \int \varrho \mathbf{u}_t \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t \, dx + \int \operatorname{div}(\varrho \mathbf{u}) \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t \, dx \\
&\leq \int (\varrho |\nabla \mathbf{u}| |\mathbf{u}_t|^2 + \varrho |\mathbf{u}| |\nabla \mathbf{u}|^2 |\mathbf{u}_t| + \varrho |\mathbf{u}|^2 |\nabla^2 \mathbf{u}| |\mathbf{u}_t| + \varrho |\mathbf{u}|^2 |\nabla \mathbf{u}| |\nabla \mathbf{u}_t|) \, dx \\
&\leq C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^4}^2 \|\nabla \mathbf{u}\|_{L^2} + C \|\mathbf{u}\|_{L^6} \|\nabla \mathbf{u}\|_{L^6}^2 \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2} \\
&\quad + C \|\mathbf{u}\|_{L^6}^2 \|\nabla^2 \mathbf{u}\|_{L^2} \|\mathbf{u}_t\|_{L^6} + C \|\mathbf{u}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^6} \|\nabla \mathbf{u}_t\|_{L^2} \\
&\leq C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^{1/2} \|\nabla \mathbf{u}_t\|_{L^2}^{3/2} + C \|\nabla \mathbf{u}\|_{L^2} \|\nabla^2 \mathbf{u}\|_{L^2}^2 \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2} \\
&\quad + C \|\nabla \mathbf{u}\|_{L^2}^2 \|\nabla^2 \mathbf{u}\|_{L^2} \|\nabla \mathbf{u}_t\|_{L^2} + C \|\nabla \mathbf{u}\|_{L^2}^2 \|\nabla^2 \mathbf{u}\|_{L^2} \|\nabla \mathbf{u}_t\|_{L^2} \\
&\leq \frac{\mu}{8} \|\nabla \mathbf{u}_t\|_{L^2}^2 + C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^4 + C \|\mathbf{H}_t\|_{L^2}^4 + C,
\end{aligned}$$

due to (3.34). Similarly,

$$\begin{aligned}
(3.42) \quad I_3 + I_4 &\leq C \|\nabla \mathbf{H}\|_{L^2} \|\mathbf{H}_t\|_{L^3} \|\mathbf{u}_t\|_{L^6} \leq C \|\mathbf{H}_t\|_{L^2}^{1/2} \|\mathbf{H}_t\|_{L^6}^{1/2} \|\nabla \mathbf{u}_t\|_{L^2} \\
&\leq \frac{\mu}{8} \|\nabla \mathbf{u}_t\|_{L^2}^2 + C \|\mathbf{H}_t\|_{L^2} \|\nabla \mathbf{H}_t\|_{L^2} \leq \frac{\mu}{8} \|\nabla \mathbf{u}_t\|_{L^2}^2 + \frac{\nu}{4} \|\nabla \mathbf{H}_t\|_{L^2}^2 + C \|\mathbf{H}_t\|_{L^2}^2,
\end{aligned}$$

and

$$\begin{aligned}
(3.43) \quad I_5 &\leq \|\nabla \mathbf{u}\|_{L^2} \|\mathbf{H}_t\|_{L^4}^2 \leq C \|\mathbf{H}_t\|_{L^2}^{3/2} \|\mathbf{H}_t\|_{L^6}^{1/2} \\
&\leq C \|\mathbf{H}_t\|_{L^2}^{3/2} \|\nabla \mathbf{H}_t\|_{L^2}^{1/2} \leq \frac{\nu}{4} \|\nabla \mathbf{H}_t\|_{L^2}^2 + C \|\mathbf{H}_t\|_{L^2}^2.
\end{aligned}$$

Substituting (3.40)–(3.43) into (3.39), one has

$$\begin{aligned}
(3.44) \quad \frac{d}{dt} \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2) \, dx &+ (\mu \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \nu \|\mathbf{H}_t\|_{L^2}^2) \leq C (\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^4 + \|\mathbf{H}_t\|_{L^2}^4) + C.
\end{aligned}$$

This together with Gronwall's inequality and (3.13) yields

$$(3.45) \quad \sup_{0 \leq t \leq T} (\|\sqrt{\varrho} \mathbf{u}_t(t)\|_{L^2}^2 + \|\mathbf{H}_t(t)\|_{L^2}^2) + \int_0^T (\|\nabla \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{H}_t\|_{L^2}^2) \, dt \leq C.$$

We refer the readers to [5], Section 2.3 for the formulation of initial data of $\|\sqrt{\varrho} \mathbf{u}(t)\|_{L^2}$ in terms of compatibility condition (1.6).

Now we will prove the boundedness of $\|\nabla \theta\|_{L^2}^2(t)$. Indeed, multiplying (3.2)₃ by θ_t and integrating the resulting equation over $\mathbb{R}^3$, one has

$$\begin{aligned}
(3.46) \quad \frac{\kappa}{2} \frac{d}{dt} \int |\nabla \theta|^2 \, dx &+ c_v \int \varrho \theta_t^2 \, dx \\
&= -c_v \int \varrho \mathbf{u} \cdot \nabla \theta \cdot \theta_t \, dx + 2\mu \int |\mathfrak{D}(\mathbf{u})|^2 \theta_t \, dx + \nu \int |\nabla \times \mathbf{H}| \theta_t \, dx \\
&\triangleq J_1 + J_2 + J_3.
\end{aligned}$$

In view of Hölder's inequality, (3.13) and (3.45), we obtain

$$(3.47) \quad \begin{aligned} J_1 &\leq C\|\mathbf{u}\|_{L^\infty}\|\sqrt{\varrho}\theta_t\|_{L^2}\|\nabla\theta\|_{L^2} \leq C\|\mathbf{u}\|_{W^{1,6}}\|\sqrt{\varrho}\theta_t\|_{L^2}\|\nabla\theta\|_{L^2} \\ &\leq C\|\nabla\mathbf{u}\|_{H^1}\|\sqrt{\varrho}\theta_t\|_{L^2}\|\nabla\theta\|_{L^2} \leq \frac{c_v}{2}\|\sqrt{\varrho}\theta_t\|_{L^2}^2 + C\|\nabla\theta\|_{L^2}^2. \end{aligned}$$

Furthermore, we get

$$(3.48) \quad \begin{aligned} J_2 &= 2\mu\frac{d}{dt}\int|\mathfrak{D}(\mathbf{u})|^2\theta\,dx - 2\mu\int(|\mathfrak{D}(\mathbf{u})|^2)_t\theta\,dx \\ &\leq 2\mu\frac{d}{dt}\int|\mathfrak{D}(\mathbf{u})|^2\theta\,dx + C\int|\nabla\mathbf{u}||\nabla\mathbf{u}_t|\theta\,dx \\ &\leq 2\mu\frac{d}{dt}\int|\mathfrak{D}(\mathbf{u})|^2\theta\,dx + C\|\nabla\mathbf{u}\|_{L^3}\|\nabla\mathbf{u}_t\|_{L^2}\|\theta\|_{L^6} \\ &\leq 2\mu\frac{d}{dt}\int|\mathfrak{D}(\mathbf{u})|^2\theta\,dx + C\|\nabla\mathbf{u}\|_{L^2}^{1/2}\|\nabla^2\mathbf{u}\|_{L^2}^{1/2}\|\nabla\mathbf{u}_t\|_{L^2}\|\nabla\theta\|_{L^2} \\ &\leq 2\mu\frac{d}{dt}\int|\mathfrak{D}(\mathbf{u})|^2\theta\,dx + C\|\nabla\mathbf{u}_t\|_{L^2}^2 + C\|\nabla\theta\|_{L^2}^2 \end{aligned}$$

and

$$(3.49) \quad \begin{aligned} J_3 &= \nu\frac{d}{dt}\int|\nabla\times\mathbf{H}|^2\theta\,dx - \nu\int(|\nabla\times\mathbf{H}|^2)_t\theta\,dx \\ &\leq \nu\frac{d}{dt}\int|\nabla\times\mathbf{H}|^2\theta\,dx + C\int|\nabla\mathbf{H}||\nabla\mathbf{H}_t|\theta\,dx \\ &\leq \nu\frac{d}{dt}\int|\nabla\times\mathbf{H}|^2\theta\,dx + C\|\nabla\mathbf{H}\|_{L^3}\|\nabla\mathbf{H}_t\|_{L^2}\|\theta\|_{L^6} \\ &\leq \nu\frac{d}{dt}\int|\nabla\times\mathbf{H}|^2\theta\,dx + C\|\nabla\mathbf{H}\|_{L^2}^{1/2}\|\nabla^2\mathbf{H}\|_{L^2}^{1/2}\|\nabla\mathbf{H}_t\|_{L^2}\|\nabla\theta\|_{L^2} \\ &\leq \nu\frac{d}{dt}\int|\nabla\times\mathbf{H}|^2\theta\,dx + C\|\nabla\mathbf{H}_t\|_{L^2}^2 + C\|\nabla\theta\|_{L^2}^2. \end{aligned}$$

Substituting (3.47)–(3.49) into (3.46), one has

$$(3.50) \quad \begin{aligned} &\frac{d}{dt}\int(\kappa|\nabla\theta|^2 - 4\mu|\mathfrak{D}(\mathbf{u})|^2\theta - 2\nu|\nabla\times\mathbf{H}|^2\theta)\,dx + c_v\int\varrho\theta_t^2\,dx \\ &\leq C(\|\nabla\mathbf{u}_t\|_{L^2}^2 + \|\nabla\mathbf{H}_t\|_{L^2}^2) + C\|\nabla\theta\|_{L^2}^2, \end{aligned}$$

which together with the fact that

$$(3.51) \quad \begin{aligned} &\int(4\mu|\mathfrak{D}(\mathbf{u})|^2 + 2\nu|\nabla\times\mathbf{H}|^2)\theta\,dx \\ &\leq C\|\theta\|_{L^6}(\|\nabla\mathbf{u}\|_{L^{12/5}}^2 + \|\nabla\mathbf{H}\|_{L^{12/5}}^2) \\ &\leq C\|\nabla\theta\|_{L^2}(\|\nabla\mathbf{u}\|_{L^2}^{3/2}\|\nabla^2\mathbf{u}\|_{L^2}^{1/2} + \|\nabla\mathbf{H}\|_{L^2}^{3/2}\|\nabla^2\mathbf{H}\|_{L^2}^{1/2}) \\ &\leq \frac{\kappa}{2}\|\nabla\theta\|_{L^2}^2 + C \end{aligned}$$

and Gronwall's inequality and (3.45) yields

$$(3.52) \quad \sup_{0 \leq t \leq T} \|\nabla \theta\|_{L^2}^2 + \int_0^T \|\sqrt{\varrho} \theta_t\|_{L^2}^2 dt \leq C.$$

Hence, the desired (3.35) follows from (3.45) and (3.52). Therefore, the proof of Lemma 3.7 is completed. $\square$

Remark 3.8. We can obtain from the Sobolev inequality that

$$(3.53) \quad \|\mathbf{u}\|_{L^\infty} \leq C \|\mathbf{u}\|_{W^{1,6}} \leq C \|\nabla \mathbf{u}\|_{H^1} \leq C(\|\nabla \mathbf{u}\|_{L^2} + \|\nabla^2 \mathbf{u}\|_{L^2}),$$

which implies that $\sup_{0 \leq t \leq T} \|\mathbf{u}\|_{L^\infty} \leq C$ due to (3.13), (3.34) and (3.35). And similarly, we have

$$\sup_{0 \leq t \leq T} \|\mathbf{H}\|_{L^\infty} \leq C.$$

Lemma 3.9. Under assumption (3.1), it holds that for any $0 \leq T < T^*$,

$$(3.54) \quad \sup_{0 \leq t \leq T} \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + \int_0^T \|\nabla \theta_t\|_{L^2}^2 dt \leq C.$$

Proof. Differentiating (3.2)₃ with respect to t and direct computing gives

$$(3.55) \quad \begin{aligned} c_v(\varrho \theta_{tt} + \varrho \mathbf{u} \cdot \nabla \theta_t) - \kappa \Delta \theta_t = & -c_v \varrho_t(\theta_t + \mathbf{u} \cdot \nabla \theta_t) - c_v \varrho(\mathbf{u}_t \cdot \nabla) \theta \\ & + 2\mu(|\mathfrak{D}(\mathbf{u})|^2)_t + \nu(|\nabla \times \mathbf{H}|^2)_t. \end{aligned}$$

Multiplying (3.55) by θ_t and integrating the resulting equation over $\mathbb{R}^3$ yield

$$(3.56) \quad \begin{aligned} & \frac{c_v}{2} \frac{d}{dt} \int \varrho \theta_t^2 dx + \kappa \int |\nabla \theta_t|^2 dx \\ & = c_v \int \operatorname{div}(\varrho \mathbf{u}) \theta_t^2 dx + c_v \int \operatorname{div}(\varrho \mathbf{u})(\mathbf{u} \cdot \nabla \theta) \theta_t dx \\ & \quad - c_v \int \varrho \mathbf{u}_t \cdot \nabla \theta \theta_t dx + 2\mu \int (|\mathfrak{D}(\mathbf{u})|^2)_t \theta_t dx \\ & \quad + \nu \int (|\nabla \times \mathbf{H}|^2)_t \theta_t dx \\ & \triangleq \sum_{i=1}^5 K_i. \end{aligned}$$

Next, we deal carefully with each term K_1 – K_5 as follows:

$$\begin{aligned}
 (3.57) \quad K_1 &\leq C \int \varrho |\mathbf{u}| |\theta_t| |\nabla \theta_t| \, dx \leq C \|\mathbf{u}\|_{L^\infty} \|\sqrt{\varrho} \theta_t\|_{L^2} \|\nabla \theta_t\|_{L^2} \\
 &\leq C(\|\mathbf{u}\|_{L^6} + \|\nabla \mathbf{u}\|_{L^6}) \|\sqrt{\varrho} \theta_t\|_{L^2} \|\nabla \theta_t\|_{L^2} \\
 &\leq C(\|\nabla \mathbf{u}\|_{L^2} + \|\nabla^2 \mathbf{u}\|_{L^2}) \|\sqrt{\varrho} \theta_t\|_{L^2} \|\nabla \theta_t\|_{L^2} \\
 &\leq C \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + \frac{\kappa}{10} \|\nabla \theta_t\|_{L^2}^2,
 \end{aligned}$$

$$\begin{aligned}
 (3.58) \quad K_2 &\leq C \int (\varrho |\mathbf{u}| |\nabla \mathbf{u}| |\nabla \theta| |\theta_t| + \varrho |\mathbf{u}|^2 |\nabla^2 \theta| |\theta_t| + \varrho |\mathbf{u}|^2 |\nabla \theta| |\nabla \theta_t|) \, dx \\
 &\leq C \|\mathbf{u}\|_{L^\infty} \|\nabla \mathbf{u}\|_{L^3} \|\nabla \theta\|_{L^2} \|\theta_t\|_{L^6} + C \|\mathbf{u}\|_{L^6}^2 \|\nabla^2 \theta\|_{L^2} \|\theta_t\|_{L^6} \\
 &\quad + C \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \theta\|_{L^2} \|\nabla \theta_t\|_{L^2} \\
 &\leq C \|\nabla \theta_t\|_{L^2} + C \|\nabla^2 \theta\|_{L^2} \|\nabla \theta_t\|_{L^2} \leq C \|\nabla^2 \theta\|_{L^2}^2 + \frac{\kappa}{10} \|\nabla \theta_t\|_{L^2}^2 + C,
 \end{aligned}$$

$$\begin{aligned}
 (3.59) \quad K_3 &\leq C \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2} \|\nabla \theta\|_{L^3} \|\theta_t\|_{L^6} \leq C \|\nabla \theta\|_{L^2}^{1/2} \|\nabla^2 \theta\|_{L^2}^{1/2} \|\nabla \theta_t\|_{L^2} \\
 &\leq C \|\nabla^2 \theta\|_{L^2}^2 + \frac{\kappa}{10} \|\nabla \theta_t\|_{L^2}^2 + C,
 \end{aligned}$$

$$\begin{aligned}
 (3.60) \quad K_4 &\leq C \int |\nabla \mathbf{u}| |\nabla \mathbf{u}_t| |\theta_t| \, dx \leq C \|\nabla \mathbf{u}\|_{L^3} \|\nabla \mathbf{u}_t\|_{L^2} \|\theta_t\|_{L^6} \\
 &\leq C \|\nabla \mathbf{u}\|_{L^3} \|\nabla \mathbf{u}_t\|_{L^2} \|\nabla \theta_t\|_{L^2} \leq C \|\nabla \mathbf{u}_t\|_{L^2}^2 + \frac{\kappa}{10} \|\nabla \theta_t\|_{L^2}^2,
 \end{aligned}$$

and

$$\begin{aligned}
 (3.61) \quad K_5 &\leq C \int |\nabla \mathbf{H}| |\nabla \mathbf{H}_t| |\theta_t| \, dx \leq C \|\nabla \mathbf{H}\|_{L^3} \|\nabla \mathbf{H}_t\|_{L^2} \|\theta_t\|_{L^6} \\
 &\leq C \|\nabla \mathbf{H}\|_{L^3} \|\nabla \mathbf{H}_t\|_{L^2} \|\nabla \theta_t\|_{L^2} \leq C \|\nabla \mathbf{H}_t\|_{L^2}^2 + \frac{\kappa}{10} \|\nabla \theta_t\|_{L^2}^2.
 \end{aligned}$$

Furthermore, in view of L^2 -estimate to equation (3.2)₃, we have

$$\begin{aligned}
 (3.62) \quad \|\nabla^2 \theta\|_{L^2}^2 &\leq C(\|\varrho \theta_t\|_{L^2}^2 + \|\varrho \mathbf{u} \cdot \nabla \theta\|_{L^2}^2 + \|\nabla \mathbf{u}\|^2_{L^2} + \|\nabla \mathbf{H}\|^2_{L^2}) \\
 &\leq C \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \theta\|_{L^2}^2 + C \|\nabla \mathbf{u}\|_{L^4}^4 + C \|\nabla \mathbf{H}\|_{L^4}^4 \\
 &\leq C \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + C \|\nabla \mathbf{u}\|_{L^2} \|\nabla^2 \mathbf{u}\|_{L^2}^3 + C \|\nabla \mathbf{H}\|_{L^2} \|\nabla^2 \mathbf{H}\|_{L^2}^3 + C \\
 &\leq C \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + C,
 \end{aligned}$$

due to (2.1), (3.35).

Substituting (3.57)–(3.61) into (3.56), we obtain

$$\begin{aligned}
 (3.63) \quad c_v \frac{d}{dt} \int \varrho \theta_t^2 \, dx + \kappa \int |\nabla \theta_t|^2 \, dx \\
 \leq C \|\nabla \mathbf{u}_t\|_{L^2}^2 + C \|\nabla \mathbf{H}_t\|_{L^2}^2 + C \|\sqrt{\varrho} \theta_t\|_{L^2}^2 + C,
 \end{aligned}$$

which together with Gronwall's inequality and (3.35) yields the desired (3.54). Therefore, the proof of Lemma 3.9 is completed. $\square$

Lemma 3.10. For $\tilde{q} \in (3, 6]$, under assumption (3.1), it holds that for $0 \leq T < T^*$,

$$(3.64) \quad \sup_{0 \leq t \leq T} (\|\varrho\|_{H^1 \cap W^{1, \tilde{q}}} + \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2) \\ + \int_0^T (\|\nabla^2 \mathbf{u}\|_{L^{\tilde{q}}}^2 + \|\nabla^2 \mathbf{H}\|_{L^{\tilde{q}}}^2 + \|\nabla^2 \theta\|_{L^{\tilde{q}}}^2) dt \leq C.$$

Proof. Firstly, it follows from (3.34), (3.35), (3.54) and (3.62) that

$$(3.65) \quad \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2 \leq C.$$

Next, the first order spatial derivatives $\partial_i \varrho$, $i = 1, 2, 3$, satisfy

$$\partial_t(\partial_i \varrho) + \mathbf{u} \cdot \nabla(\partial_i \varrho) + (\partial_i \mathbf{u}) \cdot \nabla \varrho = 0.$$

Therefore, for any $q \in (3, 6]$, the standard energy method gives

$$(3.66) \quad \frac{d}{dt} \|\nabla \varrho\|_{L^q} \leq C \|\nabla \mathbf{u}\|_{L^\infty} \|\nabla \varrho\|_{L^q} \leq C \|\nabla \mathbf{u}\|_{W^{1,6}} \|\nabla \varrho\|_{L^q} \\ \leq C (\|\nabla \mathbf{u}\|_{L^6} + \|\nabla^2 \mathbf{u}\|_{L^6}) \|\nabla \varrho\|_{L^q} \\ \leq C (1 + \|\varrho \mathbf{u}_t\|_{L^6} + \|\varrho \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^6} + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^6}) \|\nabla \varrho\|_{L^q} \\ \leq C (1 + \|\nabla \mathbf{u}_t\|_{L^2} + \|\mathbf{u}\|_{L^\infty} \|\nabla \mathbf{u}\|_{L^6} + \|\mathbf{H}\|_{L^\infty} \|\nabla \mathbf{H}\|_{L^6}) \|\nabla \varrho\|_{L^q} \\ \leq C (1 + \|\nabla \mathbf{u}_t\|_{L^2} + \|\nabla^2 \mathbf{u}\|_{L^2} + \|\nabla^2 \mathbf{H}\|_{L^2}) \|\nabla \varrho\|_{L^q} \\ \leq C (1 + \|\nabla \mathbf{u}_t\|_{L^2} + \|\nabla^2 \mathbf{u}\|_{L^2} + \|\nabla^2 \mathbf{H}\|_{L^2}) \|\nabla \varrho\|_{L^q} \\ \leq C (1 + \|\nabla \mathbf{u}_t\|_{L^2}^2) \|\nabla \varrho\|_{L^q},$$

due to (2.1), (2.2), (2.7), (3.13), (3.35), (3.65), Remark 3.8 and the interpolation inequality.

Together with Gronwall's inequality and taking $q = 2$, $\tilde{q}$ in (3.66), we have

$$(3.67) \quad \sup_{0 \leq t \leq T} \|\nabla \varrho\|_{L^2 \cap L^{\tilde{q}}} \leq C.$$

Next, in view of (2.1), (2.2), (2.7), (3.13), (3.35) and (3.65), we obtain

$$(3.68) \quad \int_0^T \|\nabla^2 \mathbf{u}\|_{L^{\tilde{q}}}^2 dt \\ \leq C \int_0^T (\|\varrho \mathbf{u}_t\|_{L^{\tilde{q}}}^2 + \|\varrho \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^{\tilde{q}}}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^{\tilde{q}}}^2) dt \\ \leq C \int_0^T (\|\varrho \mathbf{u}_t\|_{L^2}^2 + \|\varrho \mathbf{u}_t\|_{L^6}^2 + \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \mathbf{u}\|_{L^{\tilde{q}}}^2 + \|\mathbf{H}\|_{L^\infty}^2 \|\nabla \mathbf{H}\|_{L^{\tilde{q}}}^2) dt \\ \leq C \int_0^T (1 + \|\nabla \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) dt \\ \leq C.$$

Similarly,

$$\begin{aligned}
(3.69) \quad & \int_0^T \|\nabla^2 \mathbf{H}\|_{L^{\bar{q}}}^2 dt \\
& \leq C \int_0^T (\|\mathbf{H}_t\|_{L^{\bar{q}}}^2 + \|\mathbf{u} \cdot \nabla \mathbf{H}\|_{L^{\bar{q}}}^2 + \|\mathbf{H} \cdot \nabla \mathbf{u}\|_{L^{\bar{q}}}^2) dt \\
& \leq C \int_0^T (\|\mathbf{H}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^6}^2 + \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \mathbf{H}\|_{L^{\bar{q}}}^2 + \|\mathbf{H}\|_{L^\infty}^2 \|\nabla \mathbf{u}\|_{L^{\bar{q}}}^2) dt \\
& \leq C \int_0^T (1 + \|\nabla \mathbf{H}_t\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{u}\|_{L^2}^2) dt \\
& \leq C.
\end{aligned}$$

Furthermore, using the standard L^p -estimate to elliptic equation (3.2)₃, we obtain

$$\begin{aligned}
(3.70) \quad & \int_0^T \|\nabla^2 \theta\|_{L^{\bar{q}}}^2 dt \\
& \leq C \int_0^T (\|\varrho \theta_t\|_{L^{\bar{q}}}^2 + \|\varrho \mathbf{u} \cdot \nabla \theta\|_{L^{\bar{q}}}^2 + \| |\nabla \mathbf{u}|^2 \|_{L^{\bar{q}}}^2 + \| |\nabla \mathbf{H}|^2 \|_{L^{\bar{q}}}^2) dt \\
& \leq C \int_0^T (\|\varrho \theta_t\|_{L^2}^2 + \|\varrho \theta_t\|_{L^6}^2 + \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \theta\|_{L^{\bar{q}}}^2 \\
& \quad + \|\nabla \mathbf{u}\|_{L^\infty}^2 \|\nabla \mathbf{u}\|_{L^{\bar{q}}}^2 + \|\nabla \mathbf{H}\|_{L^\infty}^2 \|\nabla \mathbf{H}\|_{L^{\bar{q}}}^2) dt \\
& \leq C \int_0^T (1 + \|\nabla \theta_t\|_{L^2}^2 + \|\nabla^2 \theta\|_{L^2}^2 + (\|\nabla \mathbf{u}\|_{L^6}^2 + \|\nabla^2 \mathbf{u}\|_{L^6}^2)(\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^6}^2) \\
& \quad + (\|\nabla \mathbf{H}\|_{L^6}^2 + \|\nabla^2 \mathbf{H}\|_{L^6}^2)(\|\nabla \mathbf{H}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^6}^2)) dt \\
& \leq C \int_0^T (1 + \|\nabla \theta_t\|_{L^2}^2 + \|\nabla \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{H}_t\|_{L^2}^2) dt \\
& \leq C.
\end{aligned}$$

Thus, in view of (3.65), (3.67)–(3.70), we complete the proof of Lemma 3.10. $\square$

Remark 3.11. Define $\dot{f}$ as the material derivative of function f with $\dot{f} \triangleq f_t + \mathbf{u} \cdot \nabla f$. Then we can derive the regularity of the terms $\varrho \dot{\mathbf{u}}$ and $\varrho \dot{\theta}$ for later analysis. Indeed, one can deduce from (2.7), (3.13), (3.35) and (3.64) that

$$\begin{aligned}
(3.71) \quad & \|\varrho \dot{\mathbf{u}}\|_{L^2}^2 = \|\varrho(\mathbf{u}_t + \mathbf{u} \cdot \nabla \mathbf{u})\|_{L^2}^2 \\
& \leq C(\|\varrho \mathbf{u}_t\|_{L^2}^2 + \|\varrho \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2) \leq C(\|\varrho \mathbf{u}_t\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{L^2}^3 \|\nabla^2 \mathbf{u}\|_{L^2}) \leq C
\end{aligned}$$

and

$$\begin{aligned}
(3.72) \quad & \|\varrho \dot{\theta}\|_{L^2}^2 = \|\varrho(\theta_t + \mathbf{u} \cdot \nabla \theta)\|_{L^2}^2 \\
& \leq C(\|\varrho \theta_t\|_{L^2}^2 + \|\mathbf{u}\|_{L^\infty}^2 \|\nabla \theta\|_{L^2}^2) \leq C(\|\varrho \theta_t\|_{L^2}^2 + \|\nabla \mathbf{u}\|_{H^1}^2 \|\nabla \theta\|_{L^2}^2) \leq C.
\end{aligned}$$

Proof of Theorem 1.2. With the aid of the a priori estimates that are listed in Lemmas 3.1–3.10, we can now prove Theorem 1.2 as follows.

On the one hand, the functions $(\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, T^*) = \lim_{t \rightarrow T^*} (\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, t)$ satisfy the regularity condition on the initial data at time $t = T^*$. Furthermore, standard arguments yield $(\varrho \dot{\mathbf{u}}, \varrho \dot{\theta}) \in C([0, T^*]; L^2)$, which implies

$$(\varrho \dot{\mathbf{u}}, \varrho \dot{\theta})(x, T^*) = \lim_{t \rightarrow T^*} (\varrho \dot{\mathbf{u}}, \varrho \dot{\theta})(x, t) \in L^2.$$

Hence,

$$\begin{aligned} (-\operatorname{div}(2\mu \mathfrak{D}(\mathbf{u})) + \nabla P - \mathbf{H} \cdot \nabla \mathbf{H})|_{t=T^*} &= \sqrt{\varrho}(x, T^*) \tilde{\mathbf{g}}_1(x), \\ (\kappa \Delta \theta + 2\mu |\mathfrak{D}(\mathbf{u})|^2 + \nu |\nabla \times \mathbf{H}|^2)|_{t=T^*} &= \sqrt{\varrho}(x, T^*) \tilde{\mathbf{g}}_2(x), \end{aligned}$$

where

$$\tilde{\mathbf{g}}_1(x) \triangleq \begin{cases} \varrho^{-1/2}(x, T^*)(\varrho \dot{\mathbf{u}})(x, T^*) & \text{for } x \in \{x; \varrho(x, T^*) > 0\}, \\ 0 & \text{for } x \in \{x; \varrho(x, T^*) = 0\}, \end{cases}$$

and

$$\tilde{\mathbf{g}}_2(x) \triangleq \begin{cases} c_v \varrho^{-1/2}(x, T^*)(\varrho \dot{\theta})(x, T^*) & \text{for } x \in \{x; \varrho(x, T^*) > 0\}, \\ 0 & \text{for } x \in \{x; \varrho(x, T^*) = 0\}, \end{cases}$$

satisfying $\tilde{\mathbf{g}}_1, \tilde{\mathbf{g}}_2 \in L^2$. Thus, $(\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, T^*)$ satisfies the compatibility condition. Therefore, we can take $(\varrho, \mathbf{u}, \mathbf{H}, \theta)(x, T^*)$ as the initial data and apply Lemma 2.1 again to extend the local strong solutions beyond T^* , which contradicts the assumption that T^* is the maximal existence time of strong solutions. Therefore, we complete the proof of Theorem 1.2. $\square$

4. PROOF OF THEOREM 1.6

Throughout this section, we denote

$$C_0 \triangleq \|\sqrt{\varrho_0} \mathbf{u}_0\|_{L^2}^2 + \|\mathbf{H}_0\|_{L^2}^2.$$

Firstly, applying [17], Theorem 2.1 and integrating (3.5) with respect to t , we have the following results.

Lemma 4.1. *Let $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ be a strong solution to system (1.1)–(1.3) on $(0, T)$. Then for any $t \in (0, T)$, it holds that*

$$(4.1) \quad \|\varrho(t)\|_{L^\infty} = \|\varrho_0\|_{L^\infty}$$

and

$$(4.2) \quad \|\sqrt{\varrho} \mathbf{u}(t)\|_{L^2}^2 + \|\mathbf{H}(t)\|_{L^2}^2 + 2 \int_0^t (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) \, ds \leq C_0.$$

Lemma 4.2. *Let $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ be a strong solution to system (1.1)–(1.3) on $(0, T)$. Then for any $t \in (0, T)$, it holds that*

$$(4.3) \quad \sup_{0 \leq s \leq t} (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) \leq 2(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) \\ + C\sqrt{C_0} \sup_{0 \leq s \leq t} \|\nabla \mathbf{H}\|_{L^2}^3 \\ + CC_0 \sup_{0 \leq s \leq t} (\|\nabla \mathbf{u}\|_{L^2}^4 + \|\nabla \mathbf{H}\|_{L^2}^4),$$

where (and in what follows) C denotes a generic positive constant depending only on μ, ν and $\|\varrho_0\|_{L^\infty}$.

Proof. Multiplying (3.2)₂ by $\mathbf{u}_t$, (3.2)₄ by $\mathbf{H}_t$, and integrating the resulting equality over $\mathbb{R}^3$, we obtain from the Cauchy-Schwarz inequality that

$$(4.4) \quad \frac{1}{2} \frac{d}{dt} \int (\mu |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2) dx + \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2) dx \\ = \int \mathbf{H} \cdot \nabla \mathbf{H} \cdot \mathbf{u}_t dx - \int \varrho \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx + \int (\mathbf{H} \cdot \nabla \mathbf{u} - \mathbf{u} \cdot \nabla \mathbf{H}) \cdot \mathbf{H}_t dx \\ = -\frac{d}{dt} \int \mathbf{H} \cdot \nabla \mathbf{u} \cdot \mathbf{H} dx + \int \mathbf{H}_t \cdot \nabla \mathbf{u} \cdot \mathbf{H} dx + \int \mathbf{H} \cdot \nabla \mathbf{u} \cdot \mathbf{H}_t dx \\ - \int \varrho \mathbf{u} \cdot \nabla \mathbf{u} \cdot \mathbf{u}_t dx + \int (\mathbf{H} \cdot \nabla \mathbf{u} - \mathbf{u} \cdot \nabla \mathbf{H}) \cdot \mathbf{H}_t dx \\ \leq -\frac{d}{dt} \int \mathbf{H} \cdot \nabla \mathbf{u} \cdot \mathbf{H} dx + \frac{1}{2} \int (\varrho |\mathbf{u}_t|^2 + |\mathbf{H}_t|^2) dx \\ + C \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2) dx,$$

which implies that

$$(4.5) \quad \frac{d}{dt} \int (\mu |\nabla \mathbf{u}|^2 + \nu |\nabla \mathbf{H}|^2 + 2\mathbf{H} \cdot \nabla \mathbf{u} \cdot \mathbf{H}) dx + \|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2 \\ \leq C \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2) dx.$$

Integrating (4.5) with respect to the time variable over $(0, t)$ gives rise to

$$(4.6) \quad \sup_{0 \leq s \leq t} (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) + \int_0^t (\|\sqrt{\varrho} \mathbf{u}_s\|_{L^2}^2 + \|\mathbf{H}_s\|_{L^2}^2) ds \\ \leq (\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + 4 \sup_{0 \leq s \leq t} \int |\mathbf{H}|^2 |\nabla \mathbf{u}| dx \\ + C \int_0^t \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2) dx ds.$$

Recall that $(\mathbf{u}, P)$ satisfies the following Stokes system:

$$(4.7) \quad \begin{cases} -\mu \Delta \mathbf{u} + \nabla P = -\varrho \mathbf{u}_t - \varrho \mathbf{u} \cdot \nabla \mathbf{u} + \mathbf{H} \cdot \nabla \mathbf{H}, & x \in \mathbb{R}^3, \\ \operatorname{div} \mathbf{u} = 0, & x \in \mathbb{R}^3, \\ \mathbf{u}(x) \rightarrow 0, & |x| \rightarrow \infty. \end{cases}$$

We thus obtain from (2.7) that

$$(4.8) \quad \begin{aligned} \|\nabla^2 \mathbf{u}\|_{L^2}^2 &\leq C(\|\varrho \mathbf{u}_t\|_{L^2}^2 + \|\varrho \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2) \\ &\leq C(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2). \end{aligned}$$

Applying the classical L^2 -estimates for elliptic system on $\mathbf{H}$ gives

$$(4.9) \quad \|\nabla^2 \mathbf{H}\|_{L^2}^2 \leq C(\|\mathbf{H}_t\|_{L^2}^2 + \|\mathbf{u} \cdot \nabla \mathbf{H}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{u}\|_{L^2}^2),$$

which together with (4.8) leads to

$$(4.10) \quad \begin{aligned} \|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2 &\leq L(\|\sqrt{\varrho} \mathbf{u}_t\|_{L^2}^2 + \|\mathbf{H}_t\|_{L^2}^2) \\ &\quad + C(\|\sqrt{\varrho} \mathbf{u} \cdot \nabla \mathbf{u}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{H}\|_{L^2}^2 \\ &\quad + \|\mathbf{u} \cdot \nabla \mathbf{H}\|_{L^2}^2 + \|\mathbf{H} \cdot \nabla \mathbf{u}\|_{L^2}^2) \end{aligned}$$

for a positive constant L depending only on μ, ν and $\|\varrho\|_{L^\infty}$. Integrating (4.10) multiplied by $1/(2L)$ with respect to time variable over $(0, t)$ and adding the resulting inequality to (4.6), we have

$$(4.11) \quad \begin{aligned} &\sup_{0 \leq s \leq t} (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) \\ &\quad + \frac{1}{2L} \int_0^t (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) \, ds + \frac{1}{2} \int_0^t (\|\sqrt{\varrho} \mathbf{u}_s\|_{L^2}^2 + \|\mathbf{H}_s\|_{L^2}^2) \, ds \\ &\leq (\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + 4 \sup_{0 \leq s \leq t} \int |\mathbf{H}|^2 |\nabla \mathbf{u}| \, dx \\ &\quad + \bar{L} \int_0^t \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{H}|^2) \, dx \, ds. \end{aligned}$$

By Hölder's inequality, the Sobolev inequality and (4.2), we have

$$(4.12) \quad \begin{aligned} \int |\mathbf{H}|^2 |\nabla \mathbf{u}| \, dx &\leq \|\mathbf{H}\|_{L^4}^2 \|\nabla \mathbf{u}\|_{L^2} \leq \|\mathbf{H}\|_{L^2}^{1/2} \|\nabla \mathbf{H}\|_{L^2}^{3/2} \|\nabla \mathbf{u}\|_{L^2} \\ &\leq \frac{\mu}{8} \|\nabla \mathbf{u}\|_{L^2}^2 + C \|\mathbf{H}\|_{L^2} \|\nabla \mathbf{H}\|_{L^2}^3 \leq \frac{\mu}{8} \|\nabla \mathbf{u}\|_{L^2}^2 + C \sqrt{C_0} \|\nabla \mathbf{H}\|_{L^2}^3, \end{aligned}$$

which gives rise to

$$(4.13) \quad 4 \sup_{0 \leq s \leq t} \int |\mathbf{H}|^2 |\nabla \mathbf{u}| \, dx \leq \frac{\mu}{2} \sup_{0 \leq s \leq t} \|\nabla \mathbf{u}\|_{L^2}^2 + C \sqrt{C_0} \sup_{0 \leq s \leq t} \|\nabla \mathbf{H}\|_{L^2}^3.$$

In a similar way, we have

$$\begin{aligned}
(4.14) \quad & \bar{L} \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{H}|^2) dx \\
& \leq \bar{L} \|\varrho\|_{L^\infty} \|\mathbf{u}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla \mathbf{u}\|_{L^6} + \bar{L} \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla \mathbf{u}\|_{L^6} \\
& \quad + \bar{L} \|\mathbf{u}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla \mathbf{H}\|_{L^6} + \bar{L} \|\mathbf{H}\|_{L^6}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla \mathbf{H}\|_{L^6} \\
& \leq C \|\nabla \mathbf{u}\|_{L^2}^3 \|\nabla^2 \mathbf{u}\|_{L^2} + C \|\nabla \mathbf{H}\|_{L^2}^2 \|\nabla \mathbf{u}\|_{L^2} \|\nabla^2 \mathbf{u}\|_{L^2} \\
& \quad + C \|\nabla \mathbf{u}\|_{L^2}^2 \|\nabla \mathbf{H}\|_{L^2} \|\nabla^2 \mathbf{H}\|_{L^2} + C \|\nabla \mathbf{H}\|_{L^2}^3 \|\nabla \mathbf{H}\|_{L^6} \\
& \leq \frac{1}{4L} (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) \\
& \quad + C (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) (\|\nabla \mathbf{u}\|_{L^2}^4 + \|\nabla \mathbf{H}\|_{L^2}^4).
\end{aligned}$$

Integrating inequality (4.14) with respect to time variable over $(0, t)$ gives

$$\begin{aligned}
(4.15) \quad & \bar{L} \int_0^t \int (\varrho |\mathbf{u}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{u}|^2 + |\mathbf{u}|^2 |\nabla \mathbf{H}|^2 + |\mathbf{H}|^2 |\nabla \mathbf{H}|^2) dx ds \\
& \leq \frac{1}{4L} \int_0^t (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) ds \\
& \quad + C \sup_{0 \leq s \leq t} (\|\nabla \mathbf{u}\|_{L^2}^4 + \|\nabla \mathbf{H}\|_{L^2}^4) \int_0^t (\|\nabla \mathbf{u}\|_{L^2}^2 + \|\nabla \mathbf{H}\|_{L^2}^2) ds \\
& \leq \frac{1}{4L} \int_0^t (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) ds \\
& \quad + C \sup_{0 \leq s \leq t} (\|\nabla \mathbf{u}\|_{L^2}^4 + \|\nabla \mathbf{H}\|_{L^2}^4) \int_0^t (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) ds \\
& \leq \frac{1}{4L} \int_0^t (\|\nabla^2 \mathbf{u}\|_{L^2}^2 + \|\nabla^2 \mathbf{H}\|_{L^2}^2) ds + CC_0 \sup_{0 \leq s \leq t} (\|\nabla \mathbf{u}\|_{L^2}^4 + \|\nabla \mathbf{H}\|_{L^2}^4),
\end{aligned}$$

due to (4.2). Substituting (4.13) and (4.15) into (4.11) implies the desired (4.3) and therefore the proof of Lemma 4.2 is completed. $\square$

Lemma 4.3. *Let $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ be a strong solution to system (1.1)–(1.3) on $(0, T)$. Then there exists a positive constant ε_0 depending only on μ, ν and $\|\varrho_0\|_{L^\infty}$ such that for any $t \in (0, T)$ it holds that*

$$(4.16) \quad \sup_{0 \leq t \leq T} (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2) \leq 8(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2)$$

provided that

$$(4.17) \quad (\|\sqrt{\varrho_0} \mathbf{u}_0\|_{L^2}^2 + \|\mathbf{H}_0\|_{L^2}^2) (\|\nabla \mathbf{u}_0\|_{L^2}^2 + \|\nabla \mathbf{H}_0\|_{L^2}^2) \leq \varepsilon_0.$$

P r o o f. Define function $E(t)$ as

$$E(t) \triangleq \sup_{0 \leq s \leq t} (\mu \|\nabla \mathbf{u}\|_{L^2}^2 + \nu \|\nabla \mathbf{H}\|_{L^2}^2).$$

In view of the regularity of $\mathbf{u}$ and $\mathbf{H}$ as described in Lemma 2.1, it is easy to check that $E(t)$ is a continuous function on $[0, T]$. By (4.3), there is a positive constant M depending only on μ, ν and $\|\varrho_0\|_{L^\infty}$ such that

$$(4.18) \quad E(t) \leq 2(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + \sqrt{M} \sqrt{C_0} E^{3/2}(t) + MC_0 E^2(t).$$

Now suppose that

$$(4.19) \quad MC_0(\|\nabla \mathbf{u}_0\|_{L^2}^2 + \|\nabla \mathbf{H}_0\|_{L^2}^2) \leq \frac{1}{64(\mu + \nu)},$$

which implies

$$(4.20) \quad \begin{aligned} MC_0(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) &\leq MC_0(\mu + \nu)(\|\nabla \mathbf{u}_0\|_{L^2}^2 + \|\nabla \mathbf{H}_0\|_{L^2}^2) \\ &\leq \frac{1}{64(\mu + \nu)} \times (\mu + \nu) = \frac{1}{64}. \end{aligned}$$

Set

$$(4.21) \quad T_* \triangleq \max\{t \in [0, T] : E(s) \leq 16(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2), \forall s \in (0, t)\}.$$

We claim that

$$T_* = T.$$

Otherwise, we have $T_* \in (0, T)$. By the continuity of $E(t)$, it follows from (4.18) and (4.20) that

$$\begin{aligned} E(T_*) &\leq 2(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + \sqrt{MC_0} \cdot \sqrt{16(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2)} E(T_*) \\ &\quad + MC_0 \cdot 16(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) E(T_*) \\ &= 2(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + \sqrt{16MC_0(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2)} E(T_*) \\ &\quad + 16MC_0(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) E(T_*) \\ &\leq 2(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) + \frac{3}{4} E(T_*), \end{aligned}$$

and thus

$$E(T_*) \leq 8(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2),$$

which contradicts (4.21).

Choosing $\varepsilon_0 = 1/(64M(\mu + \nu))$, by virtue of the claim we have showed in the above, we derive that

$$E(t) \leq 8(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2), \quad 0 < t < T,$$

provided that (4.17) holds true. This gives the desired (4.16), which consequently completes the proof of Lemma 4.3. $\square$

Now we are ready to give a proof of Theorem 1.6.

Proof of Theorem 1.6. Let ε_0 be the constant stated in Lemma 4.3 and suppose the initial data $(\varrho_0, \mathbf{u}_0, \mathbf{H}_0, \theta_0)$ satisfy (1.5), (1.6), (1.7), and

$$(4.22) \quad (\|\sqrt{\varrho_0} \mathbf{u}_0\|_{L^2}^2 + \|\mathbf{H}_0\|_{L^2}^2)(\|\nabla \mathbf{u}_0\|_{L^2}^2 + \|\nabla \mathbf{H}_0\|_{L^2}^2) \leq \varepsilon_0.$$

According to Lemma 2.1, there is a unique strong solution $(\varrho, \mathbf{u}, \mathbf{H}, \theta)$ to system (1.1)–(1.3). Let T^* be the maximal existence time to that solution. We will show that $T^* = \infty$. Supposing, by contradiction, that $T^* < \infty$, then by (1.8), we deduce that for any (s, r) with $2/s + 3/r \leq 1, r > 3$ it holds that

$$\int_0^{T^*} \|\mathbf{u}\|_{L_\omega^r}^s dt = \infty,$$

which combined with the inequality $\|\mathbf{u}\|_{L_\omega^6}^4 \leq \|\mathbf{u}\|_{L^6}^4 \leq C \|\nabla \mathbf{u}\|_{L^2}^4$ leads to

$$(4.23) \quad \int_0^{T^*} \|\nabla \mathbf{u}\|_{L^2}^4 dt = \infty.$$

By Lemma 4.3, for any $0 < T < T^*$ we find that

$$\sup_{0 \leq t \leq T} \|\nabla \mathbf{u}\|_{L^2}^2 \leq 8(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2).$$

This together with (4.2) gives rise to

$$\begin{aligned} \int_0^{T^*} \|\nabla \mathbf{u}\|_{L^2}^4 dt &\leq \left(\sup_{0 \leq t \leq T^*} \|\nabla \mathbf{u}\|_{L^2}^2 \right) \int_0^{T^*} \|\nabla \mathbf{u}\|_{L^2}^2 dt \\ &\leq 8(\mu \|\nabla \mathbf{u}_0\|_{L^2}^2 + \nu \|\nabla \mathbf{H}_0\|_{L^2}^2) (2\mu)^{-1} C_0 < \infty, \end{aligned}$$

which contradicts (4.23). This contradiction implies that $T^* = \infty$, and thus we obtain global strong solution. Therefore the proof of Theorem 1.6 is completed.

Acknowledgments. The author would like to thank the reviewers for careful reading and helpful suggestions on the original manuscript.

References

- [1] *Q. Bie, Q. Wang, Z. Yao*: Global well-posedness of the 3D incompressible MHD equations with variable density. *Nonlinear Anal., Real World Appl.* **47** (2019), 85–105. [zbl](#) [MR](#) [doi](#)
- [2] *F. Chen, B. Guo, X. Zhai*: Global solution to the 3-D inhomogeneous incompressible MHD system with discontinuous density. *Kinet. Relat. Models* **12** (2019), 37–58. [zbl](#) [MR](#) [doi](#)
- [3] *F. Chen, Y. Li, H. Xu*: Global solution to the 3D nonhomogeneous incompressible MHD equations with some large initial data. *Discrete Contin. Dyn. Syst.* **36** (2016), 2945–2967. [zbl](#) [MR](#) [doi](#)
- [4] *Q. Chen, Z. Tan, Y. Wang*: Strong solutions to the incompressible magnetohydrodynamic equations. *Math. Methods Appl. Sci.* **34** (2011), 94–107. [zbl](#) [MR](#) [doi](#)
- [5] *Y. Cho, H. Kim*: Existence results for viscous polytropic fluids with vacuum. *J. Differ. Equations* **228** (2006), 377–411. [zbl](#) [MR](#) [doi](#)
- [6] *P. A. Davidson*: Introduction to Magnetohydrodynamics. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 2017. [zbl](#) [MR](#) [doi](#)
- [7] *B. Desjardins, C. Le Bris*: Remarks on a nonhomogeneous model of magnetohydrodynamics. *Differ. Integral Equ.* **11** (1998), 377–394. [zbl](#) [MR](#)
- [8] *E. Feireisl*: Dynamics of Viscous Compressible Fluids. Oxford Lecture Series in Mathematics and Its Applications 26. Oxford University Press, Oxford, 2004. [zbl](#) [MR](#) [doi](#)
- [9] *J.-F. Gerbeau, C. Le Bris*: Existence of solution for a density-dependent magnetohydrodynamic equation. *Adv. Differ. Equ.* **2** (1997), 427–452. [zbl](#) [MR](#)
- [10] *M.-H. Giga, Y. Giga, J. Saal*: Nonlinear Partial Differential Equations: Asymptotic Behavior of Solutions and Self-Similar Solutions. Progress in Nonlinear Differential Equations and their Applications 79. Birkhäuser, Basel, 2010. [zbl](#) [MR](#) [doi](#)
- [11] *L. Grafakos*: Classical Fourier Analysis. Graduate Texts in Mathematics 249. Springer, New York, 2008. [zbl](#) [MR](#) [doi](#)
- [12] *C. He, J. Li, B. Lü*: Global well-posedness and exponential stability of 3D Navier-Stokes equations with density-dependent viscosity and vacuum in unbounded domains. *Arch. Ration. Mech. Anal.* **239** (2021), 1809–1835. [zbl](#) [MR](#) [doi](#)
- [13] *C. He, Z. Xin*: On the regularity of weak solutions to the magnetohydrodynamic equations. *J. Differ. Equations* **213** (2005), 235–254. [zbl](#) [MR](#) [doi](#)
- [14] *X. Huang, Y. Wang*: Global strong solution to the 2D nonhomogeneous incompressible MHD system. *J. Differ. Equations* **254** (2013), 511–527. [zbl](#) [MR](#) [doi](#)
- [15] *H. Kim*: A blow-up criterion for the nonhomogeneous incompressible Navier-Stokes equations. *SIAM J. Math. Anal.* **37** (2006), 1417–1434. [zbl](#) [MR](#) [doi](#)
- [16] *H. Kozono, M. Yamazaki*: Uniqueness criterion of weak solutions to the stationary Navier-Stokes equations in exterior domains. *Nonlinear Anal., Theory Methods Appl.* **38** (1999), 959–970. [zbl](#) [MR](#) [doi](#)
- [17] *P.-L. Lions*: Mathematical Topics in Fluid Mechanics. Vol. 1. Incompressible Models. Oxford Lecture Series in Mathematics and Its Applications 3. Oxford University Press, Oxford, 1996. [zbl](#) [MR](#)
- [18] *H. Sohr*: The Navier-Stokes Equations: An Elementary Functional Analytic Approach. Birkhäuser Advanced Texts. Birkhäuser, Basel, 2001. [zbl](#) [MR](#) [doi](#)
- [19] *Y. Wang*: Weak Serrin-type blowup criterion for three-dimensional nonhomogeneous viscous incompressible heat conducting flows. *Physica D* **402** (2020), Article ID 132203, 8 pages. [zbl](#) [MR](#) [doi](#)
- [20] *W. Wang, H. Yu, P. Zhang*: Global strong solutions for 3D viscous incompressible heat conducting Navier-Stokes flows with the general external force. *Math. Methods Appl. Sci.* **41** (2018), 4589–4601. [zbl](#) [MR](#) [doi](#)
- [21] *H. Wu*: Strong solutions to the incompressible magnetohydrodynamic equations with vacuum. *Comput. Math. Appl.* **61** (2011), 2742–2753. [zbl](#) [MR](#) [doi](#)

- [22] *X. Zhong*: Global strong solution for 3D viscous incompressible heat conducting Navier-Stokes flows with non-negative density. *J. Differ. Equations* *263* (2017), 4978–4996. [zbl](#) [MR](#) [doi](#)
- [23] *X. Zhong*: Global strong solutions for 3D viscous incompressible heat conducting magnetohydrodynamic flows with non-negative density. *J. Math. Anal. Appl.* *446* (2017), 707–729. [zbl](#) [MR](#) [doi](#)
- [24] *X. Zhong*: Global well-posedness to the 2D Cauchy problem of nonhomogeneous heat conducting magnetohydrodynamic equations with large initial data and vacuum. *Calc. Var. Partial Differ. Equ.* *60* (2021), Article ID 64, 24 pages. [zbl](#) [MR](#) [doi](#)
- [25] *X. Zhong*: Global existence and large time behavior of strong solutions to the nonhomogeneous heat conducting magnetohydrodynamic equations with large initial data and vacuum. *Anal. Appl., Singap.* *20* (2022), 193–219. [zbl](#) [MR](#) [doi](#)
- [26] *X. Zhong*: Global well-posedness and exponential decay of 2D nonhomogeneous Navier-Stokes and magnetohydrodynamic equations with density-dependent viscosity and vacuum. *J. Geom. Anal.* *32* (2022), Article ID 19, 26 pages. [zbl](#) [MR](#) [doi](#)
- [27] *L. Zhou*: Serrin-type blowup criterion of three-dimensional nonhomogeneous heat conducting magnetohydrodynamic flows with vacuum. *Electron. J. Qual. Theory Differ. Equ.* *2019* (2019), Article ID 81, 16 pages. [zbl](#) [MR](#) [doi](#)
- [28] *M. Zhu, M. Ou*: Global strong solutions to the 3D incompressible heat-conducting magnetohydrodynamic flows. *Math. Phys. Anal. Geom.* *22* (2019), Article ID 8, 17 pages. [zbl](#) [MR](#) [doi](#)

Author's address: *Huanyuan Li*, School of Mathematics and Statistics, Zhengzhou University, No. 100, Science Avenue, Zhengzhou, 450001, P.R. China, e-mail: lhymaths@zzu.edu.cn.