

Davide Carlo Demaria; Garbaccio Rosanna Bogin
Inverse systems and pretopological spaces

In: Zdeněk Frolík (ed.): Proceedings of the 12th Winter School on Abstract Analysis, Section of Topology. Circolo Matematico di Palermo, Palermo, 1984. Rendiconti del Circolo Matematico di Palermo, Serie II, Supplemento No. 6. pp. 93–106.

Persistent URL: <http://dml.cz/dmlcz/701831>

Terms of use:

© Circolo Matematico di Palermo, 1984

Institute of Mathematics of the Czech Academy of Sciences provides access to digitized documents strictly for personal use. Each copy of any part of this document must contain these *Terms of use*.



This document has been digitized, optimized for electronic delivery and stamped with digital signature within the project *DML-CZ: The Czech Digital Mathematics Library* <http://dml.cz>

Davide Carlo Demaria - Rosanna Garbaccio Bogin

Given a pretopological space $S=(X,P)$, we associate to any interior covering X of S a symmetrical pf-space S_X on the set X . Precisely, to obtain the pretopology of S_X , we take for each point x of X the principal filter of base the star of x with respect to X . Taking the pf-spaces S_X as terms, we obtain the inverse system \hat{S} of the pretopological space S . Generally the inverse limit S^* of \hat{S} is different from S ; yet $S^*=S$ when S is a Tychonoff topological space.

For each dimension n , we associate to \hat{S} an inverse system of prehomotopy groups $\Pi_n(S_X, a)$ and an inverse system of singular homology groups $H_n(S_X)$. Taking the inverse limits $\varprojlim_n \Pi_n(S_X, a)$ and $\varprojlim_n H_n(S_X)$, we obtain the shape groups $\check{\Pi}_n(S, a)$ and the Čech homology groups $\check{H}_n(S)$ of the pretopological space S .

Our shape groups have the characteristic properties of the classical shape groups. Similarly we can say for our Čech groups. All proofs, except those for the homotopy conditions, are similar to the classical ones.

The relations between our groups and the classical shape groups or Čech homology groups of a compact topological space will be expounded in another paper.

1. The inverse system of a pretopological space.

Let X be a nonempty set and $P=\{F_x\}(x \in X)$ a family of filters of X such that $\overline{x} \subset F_x$ for each $x \in X$. Such a family P is called a pretopology in X , and the pair (X, P) is called a pretopological space (see [2]). Here we will denote by S the pretopological space (X, P) , since we need to consider different pretopologies on the set X .

We recall that S is a pf-space, if each filter F_x is principal, i.e. $F_x = \overline{A_x}$ with $x \in A_x$. Moreover we say that the pf-space S is symmetrical, if $y \in A_x$ implies $x \in A_y$ for any $x, y \in X$.

We also recall that (see [1]) a covering X of X is an interior covering of S , if for any $x \in X$ there is at least one element A of X such that $A \in F_x$.

Now we consider the collection $Cov(S)$ of all interior coverings of S and we preorder it by the following:

1.1 *Definition* Let $X, X' \in Cov(S)$. We write $X < X'$ iff X' is a refinement of X .

1.2 *Remark.* Clearly $(\text{Cov}(S), \leq)$ is a directed set, since $X, X' \in \text{Cov}(S)$ implies $X \wedge X' \in \text{Cov}(S)$.

1.3 *Definition* Given $X \in \text{Cov}(S)$, we denote by $P(X)$ the pretopology in X , that we obtain taking for each $x \in X$ the principal filter of base the star $\text{St}(x, X)$ of x with respect to X . Then we put $S_X = (X, P(X))$.

1.4 *Remark.* S_X is a symmetrical pf-space, and the identity $p_X: S \rightarrow S_X$ is a precontinuous map. Moreover, if $X, X' \in \text{Cov}(S)$ and $X \leq X'$, the identity $p_{XX'}: S_{X'} \rightarrow S_X$ is a precontinuous map, and $p_X = p_{XX'} \circ p_{X'}$.

1.5 *Definition* We will denote by \hat{S} the inverse system $(S_X, p_{XX'}, \text{Cov}(S))$, and we will call it the inverse system of the pretopological space S . The projection $(p_X): S \rightarrow \hat{S}$ will be denoted by \hat{p} .

1.6 *Remark.* The inverse limit $\varprojlim S_X$ is the pretopological space $S^* = (X, P^*)$, where P^* is obtained taking for each $x \in X$ the filter on X of base $\{\text{St}(x, X)\} (X \in \text{Cov}(S))$. Generally the pretopology P^* is coarser than P ; yet $S^* = S$, if S is a completely regular topological space.

2. The morphism induced by a precontinuous map.

Let us consider two pretopological spaces S and T , their inverse systems $\hat{S} = (S_X, p_{XX'}, \text{Cov}(S))$ and $\hat{T} = (T_Y, q_{YY'}, \text{Cov}(T))$, and the projections $\hat{p}: S \rightarrow \hat{S}$ and $\hat{q}: T \rightarrow \hat{T}$.


2.1 *Proposition* Any precontinuous map $f: S \rightarrow T$ induces a morphism from \hat{S} to \hat{T} .

Proof:

a) For any $Y \in \text{Cov}(T)$, the family $\{f^{-1}(Y)\} (Y \in Y)$ is an interior covering of S . So f^{-1} induces a function from $\text{Cov}(T)$ to $\text{Cov}(S)$, which preserves the preorder. We will denote also this function by f^{-1} .

b) For each $Y \in \text{Cov}(T)$ we obtain a precontinuous map $f_Y: S_{f^{-1}(Y)} \rightarrow T_Y$ putting $f_Y(x) = f(x)$ for any $x \in X$.


c) (f_Y, f^{-1}) is a morphism from \hat{S} to \hat{T} . In fact, given $Y, Y' \in \text{Cov}(T)$ such that $Y \leq Y'$, clearly $f^{-1}(Y) \leq f^{-1}(Y')$ and the following diagram commutes:



2.2 *Definition* The morphism (f_Y, f^{-1}) will be denoted by $\hat{f}: \hat{S} \rightarrow \hat{T}$, and we will call it the morphism induced by f .

2.3 *Remark.* Let us define another function $\phi: \text{Cov}(T) \rightarrow \text{Cov}(S)$, taking for each $Y \in \text{Cov}(T)$ an interior covering $\phi(Y)$ of S , such that $f^{-1}(Y) \leq \phi(Y)$. Then for each $Y \in \text{Cov}(T)$, consider the precontinuous map $f'_Y: S_{\phi(Y)} \rightarrow T_Y$ given by $f'_Y(x) = f(x)$ for any $x \in X$. It is easy to see that (f'_Y, ϕ) is a morphism from \hat{S} to \hat{T} , which is equivalent to \hat{f} .

2.4 Remark. The morphism \hat{f} induced by f makes commutative the following diagram:



Moreover, any morphism $g = (g_Y, \psi)$ from \hat{S} to \hat{T} such that $g\hat{p} = \hat{f}\hat{q}$ is equivalent to \hat{f} .

3. The morphism associated to a prehomotopy.

Let us consider two pretopological spaces S and T , the closed interval $I=[0,1]$ of the real line with the pretopology $\{U_t\}(t \in I)$ (where U_t is the neighbourhood filter of the point t), the pretopological space $Z=S \times I$, and the inverse systems $\hat{S} = (S_X, p_{XX'}, \text{Cov}(S))$, $\hat{T} = (T_Y, q_{YY'}, \text{Cov}(T))$ and $\hat{Z} = (Z_R, \pi_{RR'}, \text{Cov}(Z))$. Then let $f: S \rightarrow T$ and $g: S \rightarrow T$ be homotopic precontinuous maps, and $H: S \times I \rightarrow T$ a prehomotopy of f to g .

3.1 Theorem We can associate to the map $H: Z \rightarrow T$ a morphism $K: \hat{Z} \rightarrow \hat{T}$, which is equivalent to \hat{H} and has properties analogous with those of homotopies.

Proof:

a) Define a function $\Phi: \text{Cov}(T) \rightarrow \text{Cov}(Z)$ as follows.

Given $V \in \text{Cov}(T)$, consider $H^{-1}(V) \in \text{Cov}(Z)$. For each point $(x,t) \in Z$, take $C_x \in H^{-1}(V)$, and then $A_x^{x,t} \in F_x$ and an open interval $V_t^{x,t} \in U_t$ such that $A_x^{x,t} \times V_t^{x,t} \subseteq C_x$. $\{U_{x,t}\}((x,t) \in Z)$, where $U_{x,t} = A_x^{x,t} \times V_t^{x,t}$, is an interior covering of Z which refines $H^{-1}(V)$.

For any $x \in S$, the family $\{U_{x,t}\}(t \in I)$ is an interior covering of the subspace $\{x\} \times I$ of Z . Since $\{x\} \times I$ is compact, there is a finite number $n(x)$ of points t_h of I such that $\bigcup_{1 \leq h \leq n(x)} U_{x,t_h} \supseteq \{x\} \times I$. Now observe that $A_x = \bigcap_{1 \leq h \leq n(x)} A_x^{x,t_h}$ belongs to the filter F_x , and put $R_x = \{W_{x,t_h}\}(1 \leq h \leq n(x))$, where W_{x,t_h} is the set $A_x \times V_{t_h}^{x,t_h}$.

Then consider the family $R = \bigvee_{x \in S} R_x$.

Clearly R is a covering of Z ; moreover R refines $H^{-1}(V)$, since $W_{x,t_h} \subseteq C_x$.

Given any $(x,t) \in Z$, we have $(x,t) \in W_{x,t_h}$ for some positive integer $h \leq n(x)$. Since $V_{t_h}^{x,t_h} \in U_t$, we have $W_{x,t_h} \in F_{(x,t)}$.

So $R \in \text{Cov}(Z)$, and we put $\Phi(V) = R$.

b) For each $V \in \text{Cov}(T)$, we consider the map $K_V: Z_{\Phi(V)} \rightarrow T_Y$, given by $K_V(x,t) = H(x,t)$. K_V is a precontinuous map, since for each $(x,t) \in Z$ we obtain $H(\text{St}((x,t), \Phi(V))) \subseteq \text{St}(H(x,t), V)$.

c) $K = (K_V, \Phi)$ is a morphism from \hat{Z} to \hat{T} .


In fact, given $V, V' \in \text{Cov}(T)$ such that $V \leq V'$, consider $\Phi(V) = \bigvee_{x \in S} R_x$ and $\Phi(V') = \bigvee_{x \in S} R'_x$,

where $R_x = \{A_x \times V_{t_h}^{x,t_h}\}(1 \leq h \leq n(x))$ and $R'_x = \{A'_x \times V'_{t_k}{}^{x,t_k}\}(1 \leq k \leq m(x))$.

Then we take the family R'' of all subsets of Z of form $A'' \times V_{h,k}''$ ($1 \leq h \leq n(x), 1 \leq k \leq m(x)$) with $A'' = A \cap A'$ and $V_{h,k}'' = V_{h,k}^{x,t_h} \cap V_{h,k}^{x,t_k} \neq \emptyset$, and put $R'' = \bigvee_{x \in S} R''_x$.

Given a point $(x,t) \in Z$, we have $A'' \times V_{h,k}'' \in F(x,t)$ iff $t \in V_{h,k}$, since $A'' \in F_x$ and $V_{h,k}$ is an open subset of I . But we find two positive integers $h \leq n(x)$ and $k \leq m(x)$ such that $t \in V_{h,k}^{x,t_h}$ and $t \in V_{h,k}^{x,t_k}$. Hence $R'' \in \text{Cov}(Z)$.

Clearly R'' refines both $\Phi(V)$ and $\Phi(V')$. Moreover the following diagram commutes:



d) The morphism K is equivalent to \hat{H} , since, for each $V \in \text{Cov}(T)$, $\Phi(V)$ is an interior covering of Z which refines $H^{-1}(V)$.

e) Observe that, for each $t \in I$, the map $h_t : S \rightarrow T$ given by $h_t(x) = H(x,t)$ is precontinuous.

Then define a function $\phi : \text{Cov}(T) \rightarrow \text{Cov}(S)$ as it follows.

Given $V \in \text{Cov}(T)$, consider $\Phi(V) = \bigvee_{x \in S} R_x$ where $R_x = \{A_x \times V_{h,k}^{x,t_h} \mid (1 \leq h \leq n(x))\}$, and take $A = \{A_x \mid x \in S\}$. Clearly $A \in \text{Cov}(S)$, since $A_x \in F_x$ for any $x \in S$. Hence we put $\phi(V) = A$.

Now consider the function $h_Y^t : S_{\Phi(V)} \rightarrow T_Y$, given by $h_Y^t(x) = h_t(x)$ for any $x \in S$.

To prove that h_Y^t is precontinuous, we have to show that $h_t(\text{St}(x,A)) \subseteq \text{St}(H(x,t),V)$.

To this purpose take a point $x_0 \in S$ such that $x_0 \in A$. For any positive integer $h \leq n(x_0)$ such that $t \in V_{h,k}^{x_0,t_h}$, we have $A_{x_0} \times V_{h,k}^{x_0,t_h} \subseteq \text{St}(x_0, \Phi(V))$; therefore (see

b)) $h_t(A_{x_0}) \subseteq \text{St}(H(x_0,t),V)$.

Hence (h_Y^t, ϕ) is a morphism from \hat{S} to \hat{T} .

Moreover (h_Y^t, ϕ) is equivalent to the morphism $\hat{h}_t : \hat{S} \rightarrow \hat{T}$ induced by h_t , because the interior covering $\phi(V)$ of S is a refinement of $h_t^{-1}(V)$.

Observe that the covering $\phi(V)$, and consequently the function $\phi : \text{Cov}(T) \rightarrow \text{Cov}(S)$ do not depend on the point t of I .

For $t=0$ we have $h_t = f$; therefore the morphism (h_Y^0, ϕ) is equivalent to \hat{f} .

Then for $t=1$ we have $h_t = g$; thus the morphism (h_Y^1, ϕ) is equivalent to \hat{g} .

3.2 Remark. We proved in b) that H is a precontinuous map from $(S \times I)_{\Phi(V)}$ to T_Y , for each $V \in \text{Cov}(T)$. But generally we cannot say that H is a precontinuous map from $S_{\Phi(V)} \times I$ to T_Y .

4. Inverse systems of pairs.

Let us consider a pretopological space $S = (X,P)$ and a subset A of X .

4.1 Definition Let J' be a subset of the index set J , and for each $i \in J$ let $A_i \subseteq X$. We say that $A = \{A_i \mid i \in J, J'\}$ is an interior covering of the pair (S,A) with

(J, J') as indexing pair, if:

(1) $\{A_i\}_{i \in J} \in \text{Cov}(S)$;

(2) for each $x \in A$ there is at least one index $j \in J'$ such that $A_j \in F_x^j$.

The collection of all interior coverings of the pair (S, A) will be denoted by $\text{Cov}(S, A)$.

4.2 *Remark.* Let $A = \{A_i\}_{i \in J, J'} \in \text{Cov}(S, A)$. The families $\{A_i\}_{i \in J}$ and $\{A_i\}_{i \in J'}$ will be denoted by A_J and $A_{J'}$, respectively. A induces in X the pretopology $P(A_J) = \{\text{St}(x, A_J)\}_{x \in X}$ and in A the pretopology $P(A_{J'}) = \{\text{St}(x, A_{J'})\}_{x \in A}$.

Clearly $P(A_J)$ induces in A a pretopology $P(A_J)^*$ which is coarser than $P(A_{J'})$.

The pair $((X, P(A_J)), (A, P(A_{J'})))$ will be denoted by $(S, A)_A$.

Clearly the identity $p_A: (S, A) \rightarrow (S, A)_A$ is a precontinuous map.

4.3 *Definition* Let $A = \{A_i\}_{i \in J, J'}$ and $B = \{B_h\}_{h \in H, H'}$ be interior coverings of the pair (S, A) . We write $A \leq B$ iff:

(1) B_H is a refinement of A_J ;

(2) $B_{H'}$ is a refinement of $A_{J'}$.

4.4 *Remark.* $(\text{Cov}(S, A), \leq)$ is a directed set.

If $A, A' \in \text{Cov}(S, A)$ and $A \leq A'$, the identity $p_{AA'}: (S, A)_{A'} \rightarrow (S, A)_A$ is a precontinuous map, and $p_A = p_{AA'} p_{A'}$.

4.5 *Definition* The inverse system $((S, A)_A, p_{AA'}, \text{Cov}(S, A))$ will be called the inverse system of the pair (S, A) , and it will be denoted by $\widehat{S, A}$. $\widehat{p} = (p_A)$ will be called the projection from (S, A) to $\widehat{S, A}$.

4.6 *Proposition* Let S and T be pretopological spaces, A a subset of S , B a subset of T , $\widehat{S, A} = ((S, A)_A, p_{AA'}, \text{Cov}(S, A))$ and $\widehat{T, B} = ((T, B)_B, q_{BB'}, \text{Cov}(T, B))$. Any precontinuous map $f: (S, A) \rightarrow (T, B)$ induces a morphism $\widehat{f}: \widehat{S, A} \rightarrow \widehat{T, B}$.

Proof: Given $B = \{B_i\}_{i \in J, J'} \in \text{Cov}(T, B)$, the family $f^{-1}(B) = \{f^{-1}(B_i)\}_{i \in J, J'}$ belongs to $\text{Cov}(S, A)$. Then, for each $B \in \text{Cov}(T, B)$, we define a precontinuous map $f_B: (S, A)_{f^{-1}(B)} \rightarrow (T, B)_B$, putting $f_B(x) = f(x)$ for each $x \in S$. (f_B, f^{-1}) is a morphism from $\widehat{S, A}$ to $\widehat{T, B}$, and we will denote it by \widehat{f} .

4.7 *Remark.* For each $B \in \text{Cov}(T, B)$, let us take $\phi(B) \in \text{Cov}(S, A)$ such that $f^{-1}(B) \leq \phi(B)$. We obtain a precontinuous map $f'_B: (S, A)_{\phi(B)} \rightarrow (T, B)_B$ putting $f'_B(x) = f(x)$ for any $x \in S$. (f'_B, ϕ) is a morphism from $\widehat{S, A}$ to $\widehat{T, B}$ which is equivalent to \widehat{f} .

4.8 *Theorem* Let S and T be pretopological spaces, $A \subseteq S$ and $B \subseteq T$. Then let f and g be homotopic precontinuous maps from (S, A) to (T, B) , and let $H: (S \times I, A \times I) \rightarrow (T, B)$ be a relative prehomotopy of f to g . We can associate to the map H a morphism $K: \widehat{S \times I, A \times I} \rightarrow \widehat{T, B}$, which is equivalent to \widehat{H} and has properties analogous with those of relative homotopies.

Proof: To simplify notations, we put $S \times I = Z$ and $A \times I = C$.

a) Given $B = \{B_i\}_{i \in J, J'} \in \text{Cov}(T, B)$, observe that $H^{-1}(B) = \{H^{-1}(B_i)\}_{i \in J, J'}$ belongs to $\text{Cov}(Z, C)$. For each $(x, t) \in Z$, take $C_{x,t} \in H^{-1}(B)$ such that:

(i) $C_{x,t} \in F_{(x,t)}$;

(ii) if $x \in A$, then $C_{x,t} \in H^{-1}(B_J)$.

Afterwards, with the same process of Theorem 3.1, for each $x \in S$ we construct a finite refinement $R_x = \{W_{x,t_h}\}_{(1 \leq h \leq n(x))}$ of the family $\{C_{x,t}\}_{(t \in I)}$, such that each W_{x,t_h} is of form $A_x \times V_{t_h}^{x,t_h}$ where $A_x \in F_x$ and $V_{t_h}^{x,t_h}$ is an open interval of I containing t_h . The family $\bigvee_{x \in S, A} R_x$ belongs to $\text{Cov}(Z, C)$, and we put $\phi(B) = \bigvee_{x \in S, A} R_x$.

b) For each $B \in \text{Cov}(T, B)$, we obtain a precontinuous map $K_B: (Z, C) \xrightarrow{\phi(B)} (T, B)_B$ putting $K_B(x, t) = H(x, t)$ for any $(x, t) \in Z$.

c) $K = (K_B, \phi)$ is a morphism from Z, C to T, B , which is equivalent to \hat{h} .

d) For each $B \in \text{Cov}(T, B)$ consider $\phi(B) = \bigvee_{x \in S, A} R_x$ and put $\phi(B) = \{A_x\}_{(x \in S, A)}$.

Clearly $\phi(B) \in \text{Cov}(S, A)$. Afterwards, given $t \in I$, we obtain a precontinuous map $h_B^t: (S, A) \xrightarrow{\phi(B)} (T, B)_B$ putting $h_B^t(x) = H(x, t)$ for any $x \in S$. Then (h_B^t, ϕ) is a morphism from $\widehat{S, A}$ to $\widehat{T, B}$, which is equivalent to the morphism $\hat{h}_t: \widehat{S, A} \rightarrow \widehat{T, B}$, where $h_t: (S, A) \rightarrow (T, B)$ is the precontinuous map given by $h_t(x) = H(x, t) \quad \forall x \in S$. The function $\phi: \text{Cov}(T, B) \rightarrow \text{Cov}(S, A)$ does not depend on t . Finally (h_B^0, ϕ) is equivalent to \hat{f} and (h_B^1, ϕ) is equivalent to \hat{g} .

The proofs of b), c), d) are analogous to the corresponding ones from Theorem 3.1.

5. Shape groups and relative shape groups.

Let us consider a pretopological space S and its inverse system $\hat{S} = (S_X, p_{XX'}, \text{Cov}(S))$.

Let x be a point of S . For any $X \in \text{Cov}(S)$ and each dimension n , we can calculate (see [2]) the prehomotopy group $\Pi_n(S_X, x)$ of S_X based at x . Moreover, given $X \leq X'$ in $\text{Cov}(S)$, the precontinuous map $p_{XX'}: S_{X'} \rightarrow S_X$ induces a homomorphism $p_{XX'}^*$ from $\Pi_n(S_{X'}, x)$ to $\Pi_n(S_X, x)$. So, for each positive integer n , we obtain the inverse system $(\Pi_n(S_X, x), p_{XX'}^*, \text{Cov}(S))$.

5.1 *Definition* We put $\check{\Pi}_n(S, x) = \varprojlim \Pi_n(S_X, x)$. The group $\check{\Pi}_n(S, x)$ will be called the n -dimensional shape group of the pretopological space S based at the point x . We will write $\check{\Pi}_n(S)$ instead of $\check{\Pi}_n(S, x)$, when $\check{\Pi}_n(S, x)$ does not depend on the point x of S .

5.2 *Remark.* If $S = \{x\}$, clearly $\check{\Pi}_n(S) = 0$ for each dimension n .

Now take a subset A of the pretopological space S , and consider the inverse system $\widehat{S, A} = ((S, A)_A, p_{AA'}, \text{Cov}(S, A))$ of the pair (S, A) .

Let x be a point of A . For each dimension n , we can consider the inverse system $(\Pi_n(S, A, x)_A, p_{AA'}^*, \text{Cov}(S, A))$ of relative prehomotopy groups. (Observe that $\Pi_n(S, A, x)_A$ denotes the n -dimensional relative prehomotopy group of the pair $(S, A)_A$ based at x).

5.3 *Definition* We put $\check{\Pi}_n(S, A, x) = \varprojlim \Pi_n(S, A, x)_A$. The group $\check{\Pi}_n(S, A, x)$ will be called the n -dimensional relative shape group of the pair (S, A) based at x . We will write $\check{\Pi}_n(S, A)$ instead of $\check{\Pi}_n(S, A, x)$, when $\check{\Pi}_n(S, A, x)$ does not depend on the

point x of A .

6. Homomorphisms between shape groups.

Let S and T be pretopological spaces, $\hat{S}=(S_X, p_{XX}, Cov(S))$ and $\hat{T} = (T_Y, q_{YY}, Cov(T))$ their inverse systems, $f:S \rightarrow T$ a precontinuous map. Then consider the morphism $\hat{f}=(f_Y, f^{-1})$ from \hat{S} to \hat{T} induced by f .

For each dimension n , the precontinuous map $f_Y: S_{f^{-1}(Y)} \rightarrow T_Y$ induces a homomorphism $f_{Y,n}^*$ from $\Pi_n(S_{f^{-1}(Y)}, x)$ to $\Pi_n(T_Y, f(x))$.

6.1 *Definition* We denote by $\check{f}_n: \check{\Pi}_n(S, x) \rightarrow \check{\Pi}_n(T, f(x))$ the homomorphism $\varprojlim f_{Y,n}^*$, and we say that it is induced by the precontinuous map $f: S \rightarrow T$.

6.2 *Remark.* Similarly, given two subsets A of S and B of T , and given a point x of A , for each dimension n we obtain the homomorphism $\check{f}_n: \check{\Pi}_n(S, A, x) \rightarrow \check{\Pi}_n(T, B, f(x))$ induced by a precontinuous map $f: (S, A) \rightarrow (T, B)$.

6.3 *Proposition* If $f: (S, A) \rightarrow (S, A)$ is the identity, then $\check{f}_n: \check{\Pi}_n(S, A, x) \rightarrow \check{\Pi}_n(S, A, x)$ is the identical isomorphism.

6.4 *Proposition* Let $f: (S, A) \rightarrow (T, B)$ and $g: (T, B) \rightarrow (Z, C)$ be precontinuous maps and $h=gf$. Then $\check{h}_n = \check{g}_n \check{f}_n$.

7. The homomorphisms $\check{\delta}_n: \check{\Pi}_n(S, A, x) \rightarrow \check{\Pi}_{n-1}(A, x)$, $\check{y}_n: \check{\Pi}_n(A, x) \rightarrow \check{\Pi}_n(S, x)$ and $\check{j}_n: \check{\Pi}_n(S, x) \rightarrow \check{\Pi}_n(S, A, x)$.

Let us take a pretopological space $S=(X, P)$, a subset A of X carrying the pretopology P^* induced by P , and a point x of A . Then consider the following three functions.

- 1) $\psi: Cov(A) \rightarrow Cov(S, A)$ associates to $\{X_i\} (i \in J')$ the family $\{A_i\} (i \in J, J')$, where $J = J' \cup \{j\}$ (with $j \notin J'$), $A_j = X$, and $A_i = X_i \cup (X - A)$ for $i \in J'$.
- 2) $\bar{\psi}: Cov(S) \rightarrow Cov(A)$ associates to $\{A_i\} (i \in J)$ the family $\{A_i \cap A\} (i \in J)$.
- 3) $\tilde{\psi}: Cov(S, A) \rightarrow Cov(S)$ associates to $\{A_i\} (i \in J, J')$ the family $\{A_i\} (i \in J)$.

For any $R \in Cov(A)$ we can define a boundary homomorphism $\delta_{R,n}^*$ from $\Pi_n(S, A, x) \xrightarrow{\psi(R)}$ to $\Pi_{n-1}(A_R, x)$ in the usual way (see [2]). It is easy to prove that $(\delta_{R,n}^*, \psi)$ is a morphism from $(\Pi_n(S, A, x)_A, p_{AA}^*, Cov(S, A))$ to $(\Pi_{n-1}(A_R, x), p_{RR}^*, Cov(A))$.

Afterwards, considering the usual homomorphisms $i_{X,n}^*: \Pi_n(A_{\tilde{\psi}(X)}, x) \rightarrow \Pi_n(S_X, x)$ and $j_{A,n}^*: \Pi_n(S_{\tilde{\psi}(A)}, x) \rightarrow \Pi_n(S, A, x)_A$, we obtain the morphisms $(i_{X,n}^*, \tilde{\psi})$ from

$(\Pi_n(A_R, x), p_{RR}^*, Cov(A))$ to $(\Pi_n(S_X, x), p_{XX}^*, Cov(S))$ and $(j_{A,n}^*, \tilde{\psi})$ from $(\Pi_n(S_X, x), p_{XX}^*, Cov(S))$ to $(\Pi_n(S, A, x)_A, p_{AA}^*, Cov(S, A))$.

7.1 *Definition* We put $\check{\delta}_n = \varprojlim \delta_{R,n}^*$, $\check{y}_n = \varprojlim i_{X,n}^*$, $\check{j}_n = \varprojlim j_{A,n}^*$.

With a standard proof we obtain:

7.2 *Proposition* Let $f: (S, A) \rightarrow (T, B)$ be a precontinuous map, $g: A \rightarrow B$ the restriction of f to A , $x \in A$, $y=f(x)$. For each dimension n , the following diagram commutes:

$$\begin{array}{ccc}
 \check{\Pi}_n(S, A, x) & \xrightarrow{\check{f}_n} & \check{\Pi}_n(T, B, y) \\
 \check{\delta}_n \downarrow & & \downarrow \check{\delta}_n \\
 \check{\Pi}_{n-1}(A, x) & \xrightarrow{\check{g}_n} & \check{\Pi}_{n-1}(B, y)
 \end{array}$$

7.3 Proposition Let S be a pretopological space, $A \subseteq S$, and $x \in A$. We obtain the following 0-sequence:

$$\dots \xrightarrow{\check{\delta}_{n+1}} \check{\Pi}_n(A, x) \xrightarrow{\check{f}_n} \check{\Pi}_n(S, x) \xrightarrow{\check{g}_n} \check{\Pi}_n(S, A, x) \xrightarrow{\check{\delta}_n} \check{\Pi}_{n-1}(A, x) \xrightarrow{\check{f}_{n-1}} \dots$$

8. The homotopy condition for shape groups.

To prove the homotopy condition (i.e. Theorem 8.3), we need a definition and a lemma.

8.1 Definition Let X and Y be sets, \mathcal{C} a partition of X, $f: X \rightarrow Y$ a function. We say that f is quasiconstant with respect to \mathcal{C} , if f is constant in each element of \mathcal{C} .

8.2 Lemma Let $h: I^n \rightarrow X$ be a precontinuous map from the unit n-cube I^n to a symmetrical pf-space X. It is possible to find a finite partition \mathcal{C} of I^n in open cells (of dimensions n, n-1, ..., 0) and a precontinuous map $k: I^n \rightarrow X$, such that:

- (i) k is quasiconstant with respect to \mathcal{C} and homotopic to h;
- (ii) moreover, if h is a n-preloop of X based at a, then also k is n-preloop of X based at a.

Proof: Let $\{\overline{F}_x\} (x \in X)$ be the pretopology of X.

a) First we consider the case $n=1$.

Since $h: I \rightarrow X$ is precontinuous, for each $z \in I$ there is an open interval V_z of I such that $h(V_z) \subseteq \overline{F}_{h(z)}$.

Since I is compact, we find a finite number m of points z_i of I, such that $\{V_{z_i}\} (1 \leq i \leq m)$ is a minimal linked covering of I, where $z_1=0, z_i < z_j$ for $i < j, z_m=1$.

Then we take $y_0=0, y_m=1$, and for each positive integer $i < m$ we choose a point $y_i \in V_{z_i} \cap V_{z_{i+1}}$. Afterwards we consider the partition

$$\mathcal{C} = \{[0, y_1[, \{y_1\},]y_1, y_2[, \dots, \{y_{m-1}\},]y_{m-1}, 1]\}$$

of I, and we define a precontinuous map $k: I \rightarrow X$ putting:

$$\begin{aligned}
 k(y_i) &= h(y_i) && \text{for } i=0, 1, \dots, m; \\
 k(]y_i, y_{i+1}[) &= \{h(z_{i+1})\} && \text{for } 0 \leq i < m.
 \end{aligned}$$

Then we obtain a prehomotopy K of k to h, putting:

$$K(z, t) = \begin{cases} k(z) & \text{if } 0 \leq t < \frac{1}{2} \\ h(z) & \text{if } \frac{1}{2} < t \leq 1 \end{cases}$$

Moreover, if h is a preloop based at a, also k is a preloop of X based at a, since $k(0)=h(0)=a$ and $k(1)=h(1)=a$.

b) Now we consider the case $n > 1$, assuming that the lemma is true for $n-1$.

To use simple notations, given a point $w=(z_1, z_2, \dots, z_n)$ of I^n , we put

$$(z_1, \dots, z_{n-1})=z, \quad z_n=u, \quad (z, u)=w.$$

For each $u \in I$, the function $h_u: I^{n-1} \rightarrow X$ given by $h_u(z)=h(z, u)$ is a precontinuous

map. So we find a finite partition \mathcal{C}_u of I^{n-1} in open cells (of dimensions $n-1, n-2, \dots, 0$) and precontinuous map $k_u: I^{n-1} \rightarrow X$ for which conditions (i) and (ii) hold.

Now take a point $u \in I$. For any cell Z of the partition \mathcal{C}_u , the image $\{k_u(Z)\}$ is a point of X . For each $Z \in \mathcal{C}_u$ there exists a point $z \in Z$, such that $\{k_u(Z)\} = \{h_u(z)\}$ and moreover z has an open neighbourhood V_z which contains the closure \bar{Z} of Z . Then, since \bar{Z} is compact, we find an open interval $W_{u,Z}$ of I containing u and such that $h(V_z \times W_{u,Z}) \subseteq F_{h(z,u)}$. Put $W_u = \bigcap_{Z \in \mathcal{C}_u} W_{u,Z}$.

Since I is compact, we find a finite number m of points $u_i \in I$ such that $\{W_{u_i}\} (1 \leq i \leq m)$ is a minimal linked covering of I , where $u_1 = 0, u_i < u_j$ for $i < j, u_m = 1$. Then we take $v_0 = 0, v_m = 1$, and we choose $v_i \in W_{u_i} \cap W_{u_{i+1}}$ for each positive integer $i < m$. Afterwards we consider the following partition of I :

$$\mathcal{C}_n = \{[0, v_1], [v_1, v_2], \dots, [v_{m-1}, 1]\}.$$

By means of \mathcal{C}_n and by means of the partitions $\mathcal{C}_{u_i} (1 \leq i \leq m)$ and $\mathcal{C}_{v_i} (1 \leq i \leq m)$ of I^{n-1} , we obtain a finite partition \mathcal{C} of I^n in open cells of dimensions $n, n-1, \dots, 0$. We define a function $k: I^n \rightarrow X$, which is quasiconstant with respect to \mathcal{C} and precontinuous, putting:

$$\begin{aligned} k(z, v_i) &= k_{v_i}(z) && \text{for } i=0, 1, \dots, m; \\ k(z, u) &= k_{u_{i+1}}(z) && \text{for } u \in]v_i, v_{i+1}[\text{ and } 0 \leq i < m. \end{aligned}$$

We obtain a prehomotopy K of k to h , putting:

$$K(w, t) = \begin{cases} k(w) & \text{if } 0 \leq t \leq \frac{1}{2} \\ h(w) & \text{if } \frac{1}{2} < t \leq 1 \end{cases}$$


Moreover, if h is a n -preloop of X based at a , clearly also k is a n -preloop of X based at a .

8.3 Theorem Let S and T be pretopological spaces, $a \in S, b \in T$, and let $f: (S, a) \rightarrow (T, b)$ and $g: (S, a) \rightarrow (T, b)$ be precontinuous maps. If f and g are homotopic, then $\check{f}_n = \check{g}_n$ for each dimension n .

Proof: Let $H: (S \times I, \{a\} \times I) \rightarrow (T, b)$ be a prehomotopy of f to g .

a) Given $V \in \text{Cov}(T)$, consider the elements $\phi(V) \in \text{Cov}(S \times I)$ and $\phi(V) \in \text{Cov}(S)$ we mentioned in 3, and recall that both $f^{-1}(V) \leq \phi(V)$ and $g^{-1}(V) \leq \phi(V)$.

The theorem is proved if the following diagram commutes:



b) Observe that $S_{\phi(V)}$ is a symmetrical pf-space. Hence, by Lemma 8.2, in each class $[h]$ of $\Pi_n(S_{\phi(V)}, a)$ we find a n -preloop k , which is quasiconstant with respect to a suitable finite partition \mathcal{C} of I^n in open cells.

c) The map $k \times 1_I: I^n \times I \rightarrow (S \times I)_{\phi(V)}$ is precontinuous.

In fact, let $(w, t) \in I^n \times I$. Since the map $k: I^n \rightarrow S_{\phi(V)}$ is precontinuous, there is a

neighbourhood U_w of w such that $k(U_w) \subseteq \text{St}(k(w), \phi(V))$. But $k(U_w)$ is a finite subset of $S_{\phi(V)}$. Put $k(U_w) = \{x_1, \dots, x_m\}$, and take a positive integer $r \leq m$. The point x_r belongs to the element A_{x_r} of $\phi(V)$. Moreover, for each $t \in I$ we find a point $t_r \in I$ and an open neighbourhood $V_{t_r}^{x_r, t_r}$ of t_r such that $A_{x_r} \times_{V_{t_r}^{x_r, t_r}} \phi(V)$. Then $V_t = \bigcap_{1 \leq r \leq m} V_{t_r}^{x_r, t_r}$ is a neighbourhood of t , such that $k(U_w) \times_{V_t} \subseteq \text{St}((k(w), t), \phi(V))$.

d) The prehomotopy H of f to g is a precontinuous map from $(S \times I)_{\phi(V)}$ to T_V . By c), $K = H(k \times 1_I)$ is a precontinuous map from $I^n \times I$ to T . Moreover K is a prehomotopy of fk to gk .

Therefore the foregoing diagram commutes, because $f_{V,n}^* p_{f^{-1}(V)\phi(V)}^* [h] = [fk]$ and $g_{V,n}^* p_{g^{-1}(V)\phi(V)}^* [h] = [gk]$.

9. Čech homology groups.

Let us consider a pretopological space $S = (X, P)$ and its inverse system $\hat{S} = (S_X, p_{XX'}, \text{Cov}(S))$.

For any $X \in \text{Cov}(S)$ and each dimension n , we can calculate (see [2]) the singular homology group $H_n(S_X)$ of S_X . Moreover, given $X \in X'$, the precontinuous map $p_{XX'} : S_{X'} \rightarrow S_X$ induces a homomorphism $p_{*}^{XX'} : H_n(S_{X'}) \rightarrow H_n(S_X)$ for each dimension n . So, for each integer $n \geq 0$, we obtain the inverse system $(H_n(S_X), p_{*}^{XX'}, \text{Cov}(S))$.

9.1 Definition We put $\check{H}_n(S) = \varprojlim H_n(S_X)$. The group $\check{H}_n(S)$ will be called the n -dimensional Čech homology group of the pretopological space S .

9.2 Remark. Clearly, if $S = \{x\}$, $\check{H}_n(S) = 0$ for $n > 0$, and $\check{H}_0(S) = Z$.

Now let A be a subset of S and $\widehat{S, A} = ((S, A)_A, p_{*}^{AA'}, \text{Cov}(S, A))$ the inverse system of the pair (S, A) . For each dimension n , we can consider the inverse system $(H_n(S, A)_A, p_{*}^{AA'}, \text{Cov}(S, A))$, where $H_n(S, A)_A$ is the n -dimensional relative singular homology group of the pair $(S, A)_A$.

9.3 Definition We put $\check{H}_n(S, A) = \varprojlim H_n(S, A)_A$. The group $\check{H}_n(S, A)$ will be called the n -dimensional relative Čech homology group of the pair (S, A) .

10. Homomorphisms between Čech homology groups.

Let S and T be pretopological spaces, $\hat{S} = (S_X, p_{XX'}, \text{Cov}(S))$ and $\hat{T} = (T_Y, q_{YY'}, \text{Cov}(T))$ their inverse systems, $f : S \rightarrow T$ a precontinuous map, and $\hat{f} = (f_Y, f^{-1})$ the morphism from \hat{S} to \hat{T} induced by f .

For each dimension n , the precontinuous map $f_Y : S_{f^{-1}(Y)} \rightarrow T_Y$ induces a homomorphism $f_{*}^{Y,n}$ from $H_n(S_{f^{-1}(Y)})$ to $H_n(T_Y)$.

10.1 Definition We denote by $\check{f}_n : \check{H}_n(S) \rightarrow \check{H}_n(T)$ the homomorphism $\varprojlim f_{*}^{Y,n}$, and we say that it is induced by the precontinuous map $f : S \rightarrow T$.

10.2 Remark. Similarly, given two subsets A of S and B of T , for each dimension n we obtain the homomorphism $\check{f}_n : \check{H}_n(S, A) \rightarrow \check{H}_n(T, B)$.

10.3 Proposition If $f : (S, A) \rightarrow (S, A)$ is the identity, then $\check{f}_n : \check{H}_n(S, A) \rightarrow \check{H}_n(S, A)$ is

the identical isomorphism.

10.4 *Proposition* Let $f:(S,A) \rightarrow (T,B)$ and $g:(T,B) \rightarrow (Z,C)$ be precontinuous maps and $h=gf$. Then $\check{h}_n = \check{g}_n \check{f}_n$.

10.5 *Proposition (Excision Theorem)* Let A and U be nonempty subsets of a pretopological space S , such that $cl(U) \subseteq int(A)$. Then the canonical injection $f:(S-U, A-U) \rightarrow (S,A)$ induces an isomorphism $\check{f}_n : H_n(S-U, A-U) \rightarrow H_n(S,A)$.

Proof: In fact we have:

(i) Let $\bar{A}=\{A_i\} (i \in J, J')$ be an element of $Cov(S,A)$ such that $P(A_j)$ induces in A the pretopology $P(A_{j'})$. Then (see [2]) the injection $f:(S-U, A-U) \rightarrow (S,A)$ induces an isomorphism $\check{f}_n^* : H_n(S-U, A-U) \rightarrow H_n(S,A)$.

(ii) Let $\bar{A}=\{A_i\} (i \in J, J') \in Cov(S,A)$. Then $\bar{A}=\{A_i\} (i \in J, J^*)$ (where $J^*=\{i \in J/A_i \cap A \neq \emptyset\}$) is such that $P(\bar{A}_{j^*})$ induces in A the pretopology $P(\bar{A}_{j'})$. Moreover $\bar{A} \subset \bar{A}_{j^*}$.

11. The homomorphisms $\check{\partial}_n : \check{H}_n(S,A) \rightarrow \check{H}_{n-1}(A)$, $\check{y}_n : \check{H}_n(A) \rightarrow \check{H}_n(S)$, $\check{y}_n : \check{H}_n(S) \rightarrow \check{H}_n(S,A)$.

Now consider a subspace A of a pretopological space S and the functions $\psi:Cov(A) \rightarrow Cov(S,A)$, $\bar{\psi}:Cov(S) \rightarrow Cov(A)$, $\check{\psi}:Cov(S,A) \rightarrow Cov(S)$ we mentioned in 7. For any $R \in Cov(A)$ we can define a boundary homomorphism $\partial_*^{R,n}$ from $H_n(S,A)_{\psi(R)}$ to $H_{n-1}(A_R)$ in the usual way (see [2]), and $(\partial_*^{R,n}, \psi)$ is a morphism from $(H_n(S,A)_A, P_A^{AA'}, Cov(S,A))$ to $(H_{n-1}(A_R), P_A^{RR'}, Cov(A))$.

Afterwards, considering the usual homomorphisms $i_*^{X,n} : H_n(A_{\bar{\psi}(X)}) \rightarrow H_n(S_X)$ and $j_*^{A,n} : H_n(S_{\bar{\psi}(A)}) \rightarrow H_n(S,A)$, we obtain the morphisms $(i_*^{X,n}, \bar{\psi})$ from $(H_n(A_R), P_A^{RR'}, Cov(A))$ to $(H_n(S_X), P_A^{XX'}, Cov(S))$ and $(j_*^{A,n}, \bar{\psi})$ from $(H_n(S_X), P_A^{XX'}, Cov(S))$ to $(H_n(S,A)_A, P_A^{AA'}, Cov(S,A))$.

11.1 *Definition* We put $\check{\partial}_n = \varprojlim \partial_*^{R,n}$, $\check{y}_n = \varprojlim i_*^{X,n}$, $\check{y}_n = \varprojlim j_*^{A,n}$.

With a standard proof, we obtain:

11.2 *Proposition* Let $f:(S,A) \rightarrow (T,B)$ be a precontinuous map and $g:A \rightarrow B$ the restriction of f to A . For each dimension n the following diagram commutes:

$$\begin{array}{ccc} \check{H}_n(S,A) & \xrightarrow{\check{f}_n} & \check{H}_n(T,B) \\ \check{\partial}_n \downarrow & & \downarrow \check{\partial}_n \\ \check{H}_{n-1}(A) & \xrightarrow{\check{g}_n} & \check{H}_{n-1}(A) \end{array}$$

11.3 *Proposition* Let S be a pretopological space and $A \subseteq S$. We obtain the following 0-sequence:

$$\dots \xrightarrow{\check{\partial}_{n+1}} \check{H}_n(A) \xrightarrow{\check{y}_n} \check{H}_n(S) \xrightarrow{\check{y}_n} \check{H}_n(S,A) \xrightarrow{\check{\partial}_n} \check{H}_{n-1}(A) \xrightarrow{\check{y}_{n-1}} \dots$$

12. The homotopy condition for Čech homology groups.

To prove the homotopy condition (i.e. Theorem 12.5), we need some previous statements.

Let $\Delta_p = [a_0 a_1 \dots a_p]$ be the standard p -simplex, and let i_1, i_2, \dots, i_n be

integers such that $0 < i_1 < i_2 < \dots < i_n < p$ where $1 \leq n \leq p$. Given a singular p -simplex σ^λ on a pretopological space X , we denote by $\sigma_{i_1 \dots i_n}^\lambda$ the singular $(n-1)$ -simplex $\sigma^\lambda(a_{i_1} \dots a_{i_n})$ product of $\sigma^\lambda: \Delta_p \rightarrow X$ and of the singular $(n-1)$ -simplex $(a_{i_1} \dots a_{i_n})$ on Δ_p . Moreover, given a function $F^\lambda: \Delta_p \times I \rightarrow X$, we will denote by $F_{i_1 \dots i_n}^\lambda$ the function $F^\lambda((a_{i_1} \dots a_{i_n}) \times 1_I): \Delta_p \times I \rightarrow X$. We will write $\sigma_{\hat{i}}^\lambda$ and $F_{\hat{i}}^\lambda$ instead of $\sigma_{0 \dots \hat{i} \dots p}^\lambda$ and $F_{0 \dots \hat{i} \dots p}^\lambda$.

12.1 *Definition* Let $\alpha = \Sigma \alpha_\lambda \sigma^\lambda$ and $\beta = \Sigma \alpha_\lambda \tau^\lambda$ be singular p -chains on a pretopological space X . We say that α is homotopic to β , if:

- (1) for each λ with $\alpha_\lambda \neq 0$, there is a prehomotopy F^λ of σ^λ to τ^λ ;
- (2) if $\sigma_{\hat{i}}^\lambda = \tau_{\hat{j}}^\lambda$, then $F_{\hat{i}}^\lambda = F_{\hat{j}}^\mu$.

With a process which is similar to the one of the classical case (see for example [5]), it is possible to prove the following:

12.2 *Proposition* Let X be a pretopological space, $\alpha = \Sigma \alpha_\lambda \sigma^\lambda$ a p -cycle on X and $\beta = \Sigma \alpha_\lambda \tau^\lambda$ a p -chain on X . If β is homotopic to α , then also β is a p -cycle on X ; moreover α and β are homologous.

12.3 *Lemma* Let X be a symmetrical pf-space and $\alpha = \Sigma \alpha_\lambda \sigma^\lambda$ a singular p -chain on X . For each λ such that $\alpha_\lambda \neq 0$, we find a finite partition \mathcal{C}_λ of Δ_p in open cells and a singular p -simplex τ^λ on X such that:

- (i) τ^λ is quasicontant with respect to \mathcal{C}_λ ;
- (ii) $\Sigma \alpha_\lambda \tau^\lambda$ is homotopic to $\Sigma \alpha_\lambda \sigma^\lambda$.


Proof: Let us consider successively the faces of all simplices σ^λ of dimensions $0, 1, \dots, p$ and let us define the corresponding faces of the simplices τ^λ . For any 0 -dimensional face $\sigma_{i_1}^\lambda$ we take $\tau_{i_1}^\lambda = \sigma_{i_1}^\lambda$.

Now let $\sigma_{i_1 \dots i_n}^\lambda$ be a n -dimensional face of a simplex σ^λ . Assume that all simplices $\tau_{i_1 \dots i_m}^\lambda$ ($0 < m < n$) and the prehomotopies $F_{i_1 \dots i_m}^\lambda$ of $\tau_{i_1 \dots i_m}^\lambda$ to $\sigma_{i_1 \dots i_m}^\lambda$ has been defined, in a way such that: if $\sigma_{i_1 \dots i_m}^\lambda = \sigma_{j_1 \dots j_m}^\mu$ for some λ, μ , then $\tau_{i_1 \dots i_m}^\lambda = \tau_{j_1 \dots j_m}^\mu$ and $F_{i_1 \dots i_m}^\lambda = F_{j_1 \dots j_m}^\mu$.

We observe that we can consider Δ_n as the n th cone $C^n(a_0)$ on $\{a_0\}$, and we denote by π^n the projection from I^n to $\Delta_n = C^n(a_0)$. The product function $\sigma_{i_1 \dots i_n}^\lambda \pi^n$ is a precontinuous map $f_{i_1 \dots i_n}^\lambda: I^n \rightarrow X$.

So we have to construct a precontinuous map $g_{i_1 \dots i_n}^\lambda: I^n \rightarrow X$ which must be homotopic to $f_{i_1 \dots i_n}^\lambda$ and quasicontant with respect to a suitable finite partition $\mathcal{C}_{i_1 \dots i_n}^\lambda$ of I^n in open cells. To obtain the map $g_{i_1 \dots i_n}^\lambda$ and the prehomotopy $H_{i_1 \dots i_n}^\lambda$ of $g_{i_1 \dots i_n}^\lambda$ to $f_{i_1 \dots i_n}^\lambda$, we follow a process similar to the one of Lemma 8.2; but now we have to recall that $H_{i_1 \dots i_n}^\lambda$ is already determined in $I^n \times I$ by the inductive hypothesis.

Finally we observe that the function $H_{i_1 \dots i_n}^\lambda$ and its restriction $g_{i_1 \dots i_n}^\lambda$ to $I^n \times \{0\}$ are relation preserving. So the functions $F_{i_1 \dots i_n}^\lambda$ and $\tau_{i_1 \dots i_n}^\lambda$ are given by the following commutative diagrams:




12.4 Remark. Now let (X,A) be a pair of pretopological spaces. Since generally A carries a pretopology finer than the one induced by X , we have to add to Definition 12.1 the following condition:

(3) if $\sigma_{i_1 \dots i_n}^\lambda$ is a singular $(n-1)$ -simplex on A , then $\tau_{i_1 \dots i_n}^\lambda$ and $F_{i_1 \dots i_n}^\lambda$ must be precontinuous maps into A .

12.5 Theorem Let S and T be pretopological spaces, $A \subseteq S$, $B \subseteq T$, and let $f:(S,A) \rightarrow (T,B)$ and $g:(S,A) \rightarrow (T,B)$ be precontinuous maps. If f and g are homotopic, then $\check{f}_n = \check{g}_n$ for each dimension n .

Proof: Let $H:(S \times I, A \times I) \rightarrow (T,B)$ be a prehomotopy of f to g . Given $B \in \text{Cov}(T,B)$, consider the elements $\phi(B) \in \text{Cov}(S \times I, A \times I)$ and $\phi(B) \in \text{Cov}(S,A)$ from Theorem 4.8, and recall that $\phi(B)$ refines both $f^{-1}(B)$ and $g^{-1}(B)$.

Then consider the following diagram:



Let $[\alpha] \in H_n(S,A)_{\phi(B)}$ and $\alpha = \sum \alpha_\lambda \sigma^\lambda$. By Lemma 12.3 and Remark 12.4, we construct a n -chain $\beta = \sum \alpha_\lambda \tau^\lambda$ such that:

- i) β is a linear combination of a finite number of n -simplices that are quasi-constant with respect to a suitable finite partition of Δ_n ;
- ii) β is homotopic to α .

Therefore $\beta \in [\alpha]$ since, by Proposition 12.2 and Remark 12.4, β is a relative cycle homologous to α . Now consider the chains $f\beta = \sum \alpha_\lambda (f\tau^\lambda)$ and $g\beta = \sum \alpha_\lambda (g\tau^\lambda)$, and observe that $f_*^B f^{-1}(B)\phi(B) ([\alpha]) = [f\beta]$ and $g_*^B g^{-1}(B)\phi(B) ([\alpha]) = [g\beta]$.

With a proof analogous to the one of Theorem 8.3, we see that $K^\lambda = H(\tau^\lambda \times 1_I)$ is a prehomotopy of $f\tau^\lambda$ to $g\tau^\lambda$; moreover $f\beta$ and $g\beta$ are homotopic chains. Hence $[f\beta] = [g\beta]$.

REFERENCES

- [1] ČECH E. "Topological spaces", Interscience, London (1966)
- [2] DEMARIA D.C. - GARBACCIO BOGIN R. "Homotopy and homology in pretopological spaces", Proc. 11th Winter School, Suppl. Rend. Circ. Matem. Palermo (to appear)
- [3] EILEMBERG S. - STEENROD N. "Foundations of Algebraic Topology", Princeton University Press (1952)
- [4] MARDEŠIĆ S. - SEGAL J. "Shape Theory", North Holland Publishing Company, Amsterdam (1982)
- [5] WALLACE A.H. "An Introduction to Algebraic Topology", Pergamon Press, London (1957)

DAVIDE CARLO DEMARIA, DIPARTIMENTO DI MATEMATICA, UNIVERSITA' DI TORINO,
VIA PRINCIPE AMEDEO 8, 10123 TORINO

ROSANNA GARBACCIO BOGIN, DIPARTIMENTO DI MATEMATICA, UNIVERSITA' DI TORINO,
VIA PRINCIPE AMEDEO 8, 10123 TORINO